


Name _____

Units?
Units?

Phys 2020, Section 1
Quiz #3 — Spring 2002

1. A very long straight wire in the plane of the page carries a current of 0.800 A in the direction shown.


a) Find the magnitude of the magnetic field at point A (located 2.00 cm from the wire).



At a distance of 2.00 cm, the magnetic field has magnitude


$$B = \frac{\mu_0 I}{2\pi r} = \frac{(4\pi \times 10^{-7} \frac{T \cdot m}{A})(0.800 A)}{2\pi (2.00 \times 10^{-2} m)} = 8.00 \times 10^{-6} T$$

b) What is the direction of the magnetic field at point A?

Using RHR-2,  field goes out of the page at point A

2. A magnet has its North pole pointing toward the middle of a circular loop of wire as shown at the right. The magnet is moved away from the loop, generating a current in the loop.

Use Lenz's Law to find the direction of the current induced in the loop. Show its direction clearly on the diagram and explain in words how you used Lenz's Law to arrive at your answer.



The magnet gives a flux directed to the right, but it is decreasing. To oppose this change the induced current in the loop must produce a flux directed to the right. From RHR-2 we see that if the induced current goes in the direction shown it will produce the needed flux.

3. A circular loop of wire having radius 12.0 cm has a resistance 4.00Ω . It is located in a magnetic field of 0.200 T directed at right angles to the plane of the coil. The coil is removed from the field in 0.350 s .

a) What is the area of the loop?

$$A = \pi r^2 = \pi (12.0 \times 10^{-2} \text{ m})^2 = 4.52 \times 10^{-2} \text{ m}^2$$

b) What is (magnitude of) the change in magnetic flux in the loop?

Magnitude of initial flux is

$$\Phi = BA \cos \phi = (0.200 \text{ T})(4.52 \times 10^{-2} \text{ m}^2)(1) = 9.05 \times 10^{-3} \text{ T} \cdot \text{m}^2$$

Since the final flux is zero this is the same as the change in flux

← weber!

c) What is the (average) induced emf in the loop?


magnitude of \mathcal{E} :

$$\mathcal{E} = N \frac{\Delta \Phi}{\Delta t} = 1 \cdot \frac{(9.05 \times 10^{-3} \text{ weber})}{(0.350 \text{ s})} = 2.59 \times 10^{-2} \text{ V}$$


d) What is the (average) induced current in the loop?

$$I = \frac{\mathcal{E}}{R} = \frac{(2.59 \times 10^{-2} \text{ V})}{(4.00 \Omega)} = 6.5 \times 10^{-3} \text{ A}$$

(Ohm's Law for loop)



You must show all your work and include the right units with your answers!



$$V = IR \quad \mu_0 = 4\pi \times 10^{-7} \frac{\text{T} \cdot \text{m}}{\text{A}} \quad B_{\text{wire}} = \frac{\mu_0 I}{2\pi r} \quad B_{\text{sol}} = \mu_0 n I$$

$$\mathcal{E} = vBL \quad \Phi = BA \cos \phi \quad \mathcal{E} = -N \frac{\Delta \Phi}{\Delta t} \quad \mathcal{E} = -L \frac{\Delta I}{\Delta t}$$

$$\lambda f = c \quad \vec{S}_{\text{pol}} = \frac{1}{2} \vec{S}_{\text{unpol}} \quad \vec{S} = \vec{S}_0 \cos^2 \theta$$