


**Phys 4610, Fall 2004**  
**Exam #2**


1. A spherical capacitor is made from two concentric spherical shells of radii  $a$  and  $b$  (with  $a < b$ ).

Show that the capacitance of this system is  $C = 4\pi\epsilon_0 \frac{ab}{(b-a)}$ .




2. A metal sphere of radius  $R$ , carrying a charge  $q$  is surrounded by a thick concentric metal shell (inner radius  $a$ , outer radius  $b$ , as shown in the picture). The shell also carries a net charge of  $q$ .

- a) Find the surface charge density  $\sigma$  at  $R$ , at  $a$  and at  $b$ .  
b) Find the potential at the center, using infinity as the reference point.




3. Point charges of  $-q$ ,  $+2q$  and  $-q$  are located on the  $z$  axis at  $z = -a$ ,  $z = 0$  and  $z = +a$  respectively.

- a) At a point given by spherical coordinates  $(r, \theta)$  what is the electrical potential?  
b) Using the binomial theorem on the terms in the answer to (a), find an approximate expression for the potential at large distances  $r$ . Your answer just needs to have the first non-zero term proportional to some power of  $r$ .




4. Two point charges,  $\pm q$ , are held above an infinite grounded conducting plane (the  $xy$  plane). Both charges are on the  $z$  axis; the  $-q$  charge is at  $z = a$  and the  $+q$  charge is at  $z = a + d$ .

Find the potential everywhere in the region  $z > 0$ .



5. We would like to solve the electrostatics problem diagrammed at the right; we have an infinite rectangular pipe which runs along the  $z$  direction, where the rectangular cross-section goes from  $x = 0$  to  $x = b$  and  $y = 0$  to  $y = a$ . The side at  $x = b$  is held at a constant potential of  $V_0$ , but the other sides are at zero potential.



I will start the problem and you finish it. The potential is independent of  $z$ , so we are solving for  $V(x, y)$ , and first looking for suitable solutions of the form  $X(x)Y(y)$ . We find that the choices

$$X(x) = \sinh\left(\frac{n\pi x}{a}\right) \quad Y(y) = \sin\left(\frac{n\pi y}{a}\right)$$

will do the trick, i.e. they satisfy the Laplace equation and the “zero” boundary conditions. The solution is then given by


$$V(x, y) = \sum_{n=1}^{\infty} C_n \sinh\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi y}{a}\right)$$

Apply the last boundary condition to get the  $C_n$ 's

6. The potential on the surface of a sphere (radius  $R$ , centered on the origin) is given by

$$V_0 = k \cos^2 2\theta$$

where  $k$  is a constant. Find the potential outside the sphere. (There is no charge outside the sphere.)




7. Explain the difference between a “bound” charge density and a “free” charge density. (Agreed, there is a little bit of arbitrariness in the distinction, but explain why we make it for practical situations involving dielectrics.)

8. A right circular cylinder of dielectric material has a frozen-in polarization (but no free charge on it). The cylinder has radius  $R$ , length  $2a$  and a uniform polarization of magnitude  $P$  directed along the axis of the cylinder. The cylinder's axis lies along the  $z$  axis and it is centered on the origin.

a) What is the distribution of bound charge in this system?

b) At a location  $z = b$  (with  $b > a$ ) on the  $z$  axis, what is the value of the electric field?



## Useful Equations

$$\int_{\mathbf{a}}^{\mathbf{b}} (\nabla T) \cdot d\mathbf{l} = T(\mathbf{b}) - T(\mathbf{a}) \quad \int_V (\nabla \cdot \mathbf{v}) d\tau = \oint_S \mathbf{v} \cdot d\mathbf{a} \quad \int_S (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = \oint_P \mathbf{v} \cdot d\mathbf{l}$$

### Spherical:

$$d\mathbf{l} = dl_r \hat{\mathbf{r}} + dl_\theta \hat{\boldsymbol{\theta}} + dl_\phi \hat{\boldsymbol{\phi}} \quad d\tau = r^2 \sin \theta dr d\theta d\phi$$

Gradient:

$$\nabla T = \frac{\partial T}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial T}{\partial \theta} \hat{\boldsymbol{\theta}} + \frac{1}{r \sin \theta} \frac{\partial T}{\partial \phi} \hat{\boldsymbol{\phi}} \quad (1)$$

Divergence:

$$\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{1}{r \sin \theta} \frac{\partial v_\phi}{\partial \phi} \quad (2)$$

Curl:

$$\nabla \times \mathbf{v} = \frac{1}{r \sin \theta} \left[ \frac{\partial}{\partial \theta} (\sin \theta v_\phi) - \frac{\partial v_\theta}{\partial \phi} \right] \hat{\mathbf{r}} + \frac{1}{r} \left[ \frac{1}{\sin \theta} \frac{\partial v_r}{\partial \phi} - \frac{\partial}{\partial r} (r v_\phi) \right] \hat{\boldsymbol{\theta}} + \frac{1}{r} \left[ \frac{\partial}{\partial r} (r v_\theta) - \frac{\partial v_r}{\partial \theta} \right] \hat{\boldsymbol{\phi}} \quad (3)$$

Laplacian:

$$\nabla^2 T = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial T}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial T}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2} \quad (4)$$

### Cylindrical:

$$d\mathbf{l} = dl_s \hat{\mathbf{s}} + dl_\phi \hat{\boldsymbol{\phi}} + dl_z \hat{\mathbf{z}} \quad d\tau = s ds d\phi dz$$

Gradient:

$$\nabla T = \frac{\partial T}{\partial s} \hat{\mathbf{s}} + \frac{1}{s} \frac{\partial T}{\partial \phi} \hat{\boldsymbol{\phi}} + \frac{\partial T}{\partial z} \hat{\mathbf{z}} \quad (5)$$

Divergence:

$$\nabla \cdot \mathbf{v} = \frac{1}{s} \frac{\partial}{\partial s} (s v_s) + \frac{1}{s} \frac{\partial v_\phi}{\partial \phi} + \frac{\partial v_z}{\partial z} \quad (6)$$

Curl:

$$\nabla \times \mathbf{v} = \left( \frac{1}{s} \frac{\partial v_z}{\partial \phi} - \frac{\partial v_\phi}{\partial z} \right) \hat{\mathbf{s}} + \left( \frac{\partial v_s}{\partial z} - \frac{\partial v_z}{\partial s} \right) \hat{\boldsymbol{\phi}} + \frac{1}{s} \left[ \frac{\partial}{\partial s} (s v_\phi) - \frac{\partial v_s}{\partial \phi} \right] \hat{\mathbf{z}} \quad (7)$$

Laplacian:

$$\nabla^2 T = \frac{1}{s} \frac{\partial}{\partial s} \left( s \frac{\partial T}{\partial s} \right) + \frac{1}{s^2} \frac{\partial^2 T}{\partial \phi^2} + \frac{\partial^2 T}{\partial z^2} \quad (8)$$

## More Math

In the figure at the right,

$$r = \sqrt{r^2 + z'^2 - 2rz' \cos \theta}$$

If  $x < 1$  then

$$(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!}x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}x^3 + \dots$$

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\cos 2\theta = 2 \cos^2 \theta - 1$$

$$\int \sin^2 x \, dx = \frac{1}{2}x - \frac{1}{4} \sin 2x$$


$$\int \cos^2 x \, dx = \frac{1}{2}x + \frac{1}{4} \sin 2x$$

$$\int_0^a \sin(n\pi y/a) \sin(n'\pi y/a) \, dy = \begin{cases} 0, & \text{if } n' \neq n \\ \frac{a}{2} & \text{if } n' = n \end{cases}$$

$$\frac{1}{r} = \frac{1}{r} \sum_{n=0}^{\infty} \left( \frac{r'}{r} \right)^n P_n(\cos \theta') \quad V(r, \theta) = \sum_{l=0}^{\infty} \left( A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta)$$

$$P_0(x) = 1 \quad P_1(x) = x \quad P_2(x) = (3x^2 - 1)/2 \quad P_3(x) = (5x^3 - 3x)/2$$

$$\int_{-1}^1 P_l(x) P_{l'}(x) \, dx = \int_0^\pi P_l(\cos \theta) P_{l'}(\cos \theta) \sin \theta \, d\theta = \begin{cases} 0 & \text{if } l' \neq l \\ \frac{2}{2l+1} & \text{if } l' = l \end{cases}$$



## Physics:

$$\mathbf{F} = \frac{1}{4\pi\epsilon_0} \frac{qQ}{r^2} \hat{\mathbf{r}} \quad \mathbf{F} = Q\mathbf{E} \quad \mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}} \quad V(\mathbf{r}) = - \int_{\mathcal{O}}^{\mathbf{r}} \mathbf{E} \cdot d\mathbf{l}$$

$$\nabla \times \mathbf{E} = 0 \quad \mathbf{E} = -\nabla V \quad -\nabla^2 V = \nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0} \quad V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\mathbf{r}')}{r} d\tau'$$

$$E_{\text{above}}^\perp - E_{\text{below}}^\perp = \frac{\sigma}{\epsilon_0} \quad \mathbf{E}_{\text{above}}^\parallel = \mathbf{E}_{\text{below}}^\parallel \quad W = \frac{1}{8\pi\epsilon_0} \sum_{\substack{i,j \\ i \neq j}}^n \frac{q_i q_j}{r_{ij}}$$

$$W = \frac{1}{2} \int \rho V \, d\tau = \frac{\epsilon_0}{2} \int E^2 \, d\tau \quad \mathbf{f} = \frac{1}{2\epsilon_0} \sigma^2 \hat{\mathbf{n}} \quad P = \frac{\epsilon_0}{2} E^2 \quad C \equiv \frac{Q}{V}$$

$$\mathbf{p} \equiv \int \mathbf{r}' \rho(\mathbf{r}') \, d\tau' \quad V_{\text{dip}}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2} \quad \mathbf{E}_{\text{dip}}(r, \theta) = \frac{p}{4\pi\epsilon_0 r^3} (2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\boldsymbol{\theta}})$$


$$\mathbf{p} = \alpha \mathbf{E} \quad \mathbf{N} = \mathbf{p} \times \mathbf{E} \quad \mathbf{F} = (\mathbf{p} \cdot \nabla) \mathbf{E} \quad U = -\mathbf{p} \cdot \mathbf{E}$$

$$\sigma_b = \mathbf{P} \cdot \hat{\mathbf{n}} \quad \rho_b = -\nabla \cdot \mathbf{P} \quad \mathbf{P} = \epsilon_0 \chi_e \mathbf{E}$$

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P} = \epsilon \mathbf{E} \quad \nabla \cdot \mathbf{D} = \rho_f \quad \oint \mathbf{D} \cdot d\mathbf{a} = Q_{f,\text{enc}}$$

## Specific Results:

$$E_z = \frac{1}{4\pi\epsilon_0} \frac{2\lambda L}{z\sqrt{z^2 + L^2}}$$



$$E_z = \frac{\sigma}{2\epsilon_0} \left[ 1 - \frac{z}{\sqrt{z^2 + R^2}} \right]$$

