

Phys 3810, Spring 2011
Hints, Problem Set #7

1. Griffiths, 4.52 Follow the example of spin given out in class (G's problem 4.31). The eigenvectors of S_z are

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

Using (4.136), find the action of the raising and lowering operators S_+ and S_- on all the eigenstates $|\frac{3}{2} m\rangle$ and then construct the matrices for these operators. Get S_x from $S_x = \frac{1}{2}(S_+ + S_-)$. You should get

$$S_x = \frac{\hbar}{2} \begin{pmatrix} 0 & \sqrt{3} & 0 & 0 \\ \sqrt{3} & 0 & 2 & 0 \\ 0 & 2 & 0 & \sqrt{3} \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}$$

but *show* this!

You just need to get eigenvalues of S_x but it should be clear what they ought to be! For this you need to take the determinant of a 4×4 matrix which needs to be done by an expansion (not by zipping along all the diagonals as you can for 3×3).

2. Griffiths, 5.2 (a) Show that the fractional difference between m_e and μ_H is 5.4×10^{-4} . This is the same as the fractional change in the binding energy. (Show all of this!)

(b) It's same fractional correction to R ; one finds that for the H atom

$$R_H = 1.096 \times 10^7 \text{ m}^{-1}$$

The fractional difference between μ_H and μ_D (reduced masses for the H and D atoms) is 2.7×10^{-4} . Take differentials to get the fractional change in the Balmer wavelength; it comes out to about 17.9 nm.

(c) The reduced mass for positronium is half the electron mass!

(d) The reduced mass for muonium is 185.9 times the electron mass. That's the factor by which you need to fix R from the value given in the book. With this new value of R , get Lyman- α . It comes out to about 6.54×10^{-10} m.

3. Griffiths, 5.3 The energy of the photon emitted in the transition (always between adjacent HO states) is $\hbar\omega$. The *frequency* of the radiation is

$$\nu = \frac{\omega}{2\pi} = \frac{1}{2\pi} \sqrt{\frac{k}{\mu}}$$

where μ is the reduced mass of the oscillator system. Show that if μ changes, the change in frequency is related to the change in μ by

$$d\nu = -\frac{1}{2}\nu \frac{d\mu}{\mu}$$

What is the fractional difference in reduced mass between the two molecules?

4. *Griffiths*, **5.19** I get (with the help of Maple's `fsolve` function) a root of $z = 2.628$), leading to an energy of

$$E = 0.345 \text{ eV}$$

But *show all of this*.

5. *Griffiths*, **5.35**

6. *Griffiths*, **5.36**