12/5/2012

Di Maction.

$$S = S_0 \left\{ \frac{\sin(\phi h)}{\frac{1}{2}} \right\}$$

32.29 Second secondary maxinum, of intensity

2 nd mm: sin0 = 2 1 3rd mm: 5M0 = 32 Maxin question in between those,  $\sin \theta = \frac{5\lambda}{2\alpha}$ Than  $\phi = 5\pi$ , plug in Sin(57/2) 72 = 1.6 × 10-2

Three-slit system first min occurs at angle 5° where is next minyoum? M-slit pattern

N-slit pattern

N=H sytty

M=H sytty

M= mteger not minit of W

Minim

S m=1 d sin50 = 1.2

3 2 = 3.829

Next maximun? m= maximu d sin 0 = m > m=integw. Sln 0 = 7 0 = 15.16°

A thin film of toluene (n=1.49) floats on water. Fmd minimum fllm thickness if most strongly rull'd light

otc. 
$$d = \frac{460 \text{ nm}}{2 (1.49)2} = 77.2 \text{ nm}$$

$$\frac{2}{32.45} = \frac{460 \text{ nm}}{2 (1.49)2} = 77.2 \text{ nm}$$

$$\frac{2}{32.45} = \frac{460 \text{ nm}}{2 (1.49)2} = \frac{77.2 \text{ nm}}{2 (1.49)2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2}$$

$$\frac{32.45}{-513e^2} = \frac{460 \text{ nm}}{2 (1.49)^2} = \frac{77.2 \text{ nm}}{2 (1.49)^2} =$$

$$= O_{nin} = 1.22 \frac{\lambda}{D} = 2.54 \times 10^{10}$$

$$= D = 2.6 \times 10^{3} \text{m} \quad (Bi5!)$$



Two perfectly flat glass plates sep'd one end by paper, 0.065 mm. 550 nm. How many bright bands appear observer above? 2d = (m+2) > Glass m = 235.8 => 235 frings Glas

32.62 On growd Onn = larcsecond

If no an, what

= 1 2600 degree apartne size brresjon is 2 = 550 x10 m Omn = 1.22 2 to this = 360° ldeg, (T) = 4.85×10° D= 0.138m