

Class 8: PCA Mini Project

Keilyn Duarte (PID A16881868)

Today we will do a complete analysis of some breast cancer biopsy data. But first, let's revisit the main PCA function in R `prcomp()` and see what `scale=TRUE/FALSE` does.

```
head(mtcars)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1
Hornet 4 Drive	21.4	6	258	110	3.08	3.215	19.44	1	0	3	1
Hornet Sportabout	18.7	8	360	175	3.15	3.440	17.02	0	0	3	2
Valiant	18.1	6	225	105	2.76	3.460	20.22	1	0	3	1

Find the mean value per column of this dataset.

```
apply(mtcars, 2, mean)
```

mpg	cyl	disp	hp	drat	wt	qsec
20.090625	6.187500	230.721875	146.687500	3.596563	3.217250	17.848750
vs	am	gear	carb			
0.437500	0.406250	3.687500	2.812500			

Find the standard deviation per column of this data set.


```
apply(mtcars, 2, sd)
```

mpg	cyl	disp	hp	drat	wt
6.0269481	1.7859216	123.9386938	68.5628685	0.5346787	0.9784574
qsec	vs	am	gear	carb	
1.7869432	0.5040161	0.4989909	0.7378041	1.6152000	

It is clear “disp” and “hp” have the highest mean values and standard deviation. They will likely dominate any analysis I do on this dataset. Let’s see:

```
pc.noscale <- prcomp(mtcars, scale = F)
pc.scale <- prcomp(mtcars, scale = T)
```

```
# Bi plot of mtcars dataset with no scaling
biplot(pc.noscale)
```



```
# How much each column contributes to the car's position on the plot PC1 vs. PC2
pc.noscale$rotation[,1]
```


mpg	cyl	disp	hp	drat	wt
-0.038118199	0.012035150	0.899568146	0.434784387	-0.002660077	0.006239405
qsec	vs	am	gear	carb	
-0.006671270	-0.002729474	-0.001962644	-0.002604768	0.005766010	

Plot the loadings

```
library(ggplot2)


r1 <- as.data.frame(pc.noscale$rotation)
r1$names <- rownames(pc.noscale$rotation)

ggplot(r1) +
  aes(PC1, names) +
  geom_col()
```




```
# ggplot with scaling, gives a fairer distribution
r2 <- as.data.frame(pc.scale$rotation)
r2$names <- rownames(pc.scale$rotation)

ggplot(r2) +
  aes(PC1, names) +
  geom_col()
```



```
# Bi plot of mtcars with scaling makes it easier to see patterns and similarities in the cars
biplot(pc.scale)
```



Take-home: Generally, we always want to set `scale=TRUE` when we do this type of analysis to avoid our analysis being dominated by individual variables with the largest variance, just due to their unit of measurement.

FNA Breast Cancer Data

Load the data into R.

```
wisc.df <- read.csv("WisconsinCancer.csv", row.names = 1)
head(wisc.df)
```

	diagnosis	radius_mean	texture_mean	perimeter_mean	area_mean
842302	M	17.99	10.38	122.80	1001.0
842517	M	20.57	17.77	132.90	1326.0
84300903	M	19.69	21.25	130.00	1203.0
84348301	M	11.42	20.38	77.58	386.1
84358402	M	20.29	14.34	135.10	1297.0
843786	M	12.45	15.70	82.57	477.1

	smoothness_mean	compactness_mean	concavity_mean	concave.points_mean
842302	0.11840	0.27760	0.3001	0.14710
842517	0.08474	0.07864	0.0869	0.07017
84300903	0.10960	0.15990	0.1974	0.12790
84348301	0.14250	0.28390	0.2414	0.10520
84358402	0.10030	0.13280	0.1980	0.10430
843786	0.12780	0.17000	0.1578	0.08089

	symmetry_mean	fractal_dimension_mean	radius_se	texture_se	perimeter_se
842302	0.2419	0.07871	1.0950	0.9053	8.589
842517	0.1812	0.05667	0.5435	0.7339	3.398
84300903	0.2069	0.05999	0.7456	0.7869	4.585
84348301	0.2597	0.09744	0.4956	1.1560	3.445
84358402	0.1809	0.05883	0.7572	0.7813	5.438
843786	0.2087	0.07613	0.3345	0.8902	2.217

	area_se	smoothness_se	compactness_se	concavity_se	concave.points_se
842302	153.40	0.006399	0.04904	0.05373	0.01587
842517	74.08	0.005225	0.01308	0.01860	0.01340
84300903	94.03	0.006150	0.04006	0.03832	0.02058
84348301	27.23	0.009110	0.07458	0.05661	0.01867
84358402	94.44	0.011490	0.02461	0.05688	0.01885
843786	27.19	0.007510	0.03345	0.03672	0.01137

	symmetry_se	fractal_dimension_se	radius_worst	texture_worst
842302	0.03003	0.006193	25.38	17.33

842517	0.01389	0.003532	24.99	23.41
84300903	0.02250	0.004571	23.57	25.53
84348301	0.05963	0.009208	14.91	26.50
84358402	0.01756	0.005115	22.54	16.67
843786	0.02165	0.005082	15.47	23.75
	perimeter_worst	area_worst	smoothness_worst	compactness_worst
842302	184.60	2019.0	0.1622	0.6656
842517	158.80	1956.0	0.1238	0.1866
84300903	152.50	1709.0	0.1444	0.4245
84348301	98.87	567.7	0.2098	0.8663
84358402	152.20	1575.0	0.1374	0.2050
843786	103.40	741.6	0.1791	0.5249
	concavity_worst	concave.points_worst	symmetry_worst	
842302	0.7119	0.2654	0.4601	
842517	0.2416	0.1860	0.2750	
84300903	0.4504	0.2430	0.3613	
84348301	0.6869	0.2575	0.6638	
84358402	0.4000	0.1625	0.2364	
843786	0.5355	0.1741	0.3985	
	fractal_dimension_worst			
842302	0.11890			
842517	0.08902			
84300903	0.08758			
84348301	0.17300			
84358402	0.07678			
843786	0.12440			

Q1. How many observations are in this dataset?

```
nrow(wisc.df)
```

```
[1] 569
```

Q2. How many of the observations have a malignant diagnosis?

```
sum(wisc.df$diagnosis == "M")
```

```
[1] 212
```

The `table()` function is useful here as well.

```
table(wisc.df$diagnosis)
```

```
   B    M  
357 212
```

Q3. How many variables/features in the data are suffixed with `_mean`?

```
ncol(wisc.df)
```

```
[1] 31
```

```
colnames(wisc.df)
```

```
[1] "diagnosis"           "radius_mean"  
[3] "texture_mean"        "perimeter_mean"  
[5] "area_mean"           "smoothness_mean"  
[7] "compactness_mean"    "concavity_mean"  
[9] "concave.points_mean" "symmetry_mean"  
[11] "fractal_dimension_mean" "radius_se"  
[13] "texture_se"          "perimeter_se"  
[15] "area_se"             "smoothness_se"  
[17] "compactness_se"      "concavity_se"  
[19] "concave.points_se"   "symmetry_se"  
[21] "fractal_dimension_se" "radius_worst"  
[23] "texture_worst"       "perimeter_worst"  
[25] "area_worst"          "smoothness_worst"  
[27] "compactness_worst"   "concavity_worst"  
[29] "concave.points_worst" "symmetry_worst"  
[31] "fractal_dimension_worst"
```

A useful function for this is `grep()`

```
length( grep("_mean" , colnames(wisc.df)) )
```

```
[1] 10
```

Before going any further, we need to exclude the diagnoses column for any future analysis - this tells us whether a sample is cancer or non-cancer.


```
diagnosis <- as.factor(wisc.df$diagnosis)
head(diagnosis)
```

```
[1] M M M M M M
Levels: B M
```

```
wisc.data <- wisc.df[, -1]
```

Let's see if we can cluster the `wisc.data` to find some structure in the dataset.

```
# The cluster dendrogram for this data is very messy
hc <- hclust( dist(wisc.data))
plot(hc)
```



```
dist(wisc.data)
hclust (*, "complete")
```

Principal Component Anaylsis (PCA)

```
wisc.pr <- prcomp (wisc.data, scale = T)
summary(wisc.pr)
```


Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

Q4. From your results, what proportion of the original variance is captured by the first principal components (PC1)?

Based on the summary, PC1 captures 44% of the original variance.


Q5. How many principal components (PCs) are required to describe at least 70% of the original variance in the data?

The cumulative proportion shows that up to PC3 is required to describe at least 70% of the original variance.

Q6. How many principal components (PCs) are required to describe at least 90% of the original variance in the data?

The cumulative proportion shows that up to PC6 is required to describe at least 90% of the original variance.

```
biplot(wisc.pr)
```



Q7. What stands out to you about this plot? Is it easy or difficult to understand? Why?

It is really difficult to understand because it is difficult to see trends when there is a big mess on the plot.

This biplot sucks ! We need to build our own PCA score plot of PC1 vs. PC2

```
head(wisc.pr$x)
```


	PC1	PC2	PC3	PC4	PC5	PC6
842302	-9.184755	-1.946870	-1.1221788	3.6305364	1.1940595	1.41018364
842517	-2.385703	3.764859	-0.5288274	1.1172808	-0.6212284	0.02863116
84300903	-5.728855	1.074229	-0.5512625	0.9112808	0.1769302	0.54097615
84348301	-7.116691	-10.266556	-3.2299475	0.1524129	2.9582754	3.05073750
84358402	-3.931842	1.946359	1.3885450	2.9380542	-0.5462667	-1.22541641
843786	-2.378155	-3.946456	-2.9322967	0.9402096	1.0551135	-0.45064213

	PC7	PC8	PC9	PC10	PC11	PC12
842302	2.15747152	0.39805698	-0.15698023	-0.8766305	-0.2627243	-0.8582593
842517	0.01334635	-0.24077660	-0.71127897	1.1060218	-0.8124048	0.1577838
84300903	-0.66757908	-0.09728813	0.02404449	0.4538760	0.6050715	0.1242777
84348301	1.42865363	-1.05863376	-1.40420412	-1.1159933	1.1505012	1.0104267
84358402	-0.93538950	-0.63581661	-0.26357355	0.3773724	-0.6507870	-0.1104183

843786	0.49001396	0.16529843	-0.13335576	-0.5299649	-0.1096698	0.0813699
	PC13	PC14	PC15	PC16	PC17	
842302	0.10329677	-0.690196797	0.601264078	0.74446075	-0.26523740	
842517	-0.94269981	-0.652900844	-0.008966977	-0.64823831	-0.01719707	
84300903	-0.41026561	0.016665095	-0.482994760	0.32482472	0.19075064	
84348301	-0.93245070	-0.486988399	0.168699395	0.05132509	0.48220960	
84358402	0.38760691	-0.538706543	-0.310046684	-0.15247165	0.13302526	
843786	-0.02625135	0.003133944	-0.178447576	-0.01270566	0.19671335	
	PC18	PC19	PC20	PC21	PC22	
842302	-0.54907956	0.1336499	0.34526111	0.096430045	-0.06878939	
842517	0.31801756	-0.2473470	-0.11403274	-0.077259494	0.09449530	
84300903	-0.08789759	-0.3922812	-0.20435242	0.310793246	0.06025601	
84348301	-0.03584323	-0.0267241	-0.46432511	0.433811661	0.20308706	
84358402	-0.01869779	0.4610302	0.06543782	-0.116442469	0.01763433	
843786	-0.29727706	-0.1297265	-0.07117453	-0.002400178	0.10108043	
	PC23	PC24	PC25	PC26	PC27	
842302	0.08444429	0.175102213	0.150887294	-0.201326305	-0.25236294	
842517	-0.21752666	-0.011280193	0.170360355	-0.041092627	0.18111081	
84300903	-0.07422581	-0.102671419	-0.171007656	0.004731249	0.04952586	
84348301	-0.12399554	-0.153294780	-0.077427574	-0.274982822	0.18330078	
84358402	0.13933105	0.005327110	-0.003059371	0.039219780	0.03213957	
843786	0.03344819	-0.002837749	-0.122282765	-0.030272333	-0.08438081	
	PC28	PC29	PC30			
842302	-0.0338846387	0.045607590	0.0471277407			
842517	0.0325955021	-0.005682424	0.0018662342			
84300903	0.0469844833	0.003143131	-0.0007498749			
84348301	0.0424469831	-0.069233868	0.0199198881			
84358402	-0.0347556386	0.005033481	-0.0211951203			
843786	0.0007296587	-0.019703996	-0.0034564331			


Plot of PC1 vs. PC2, the first two columns

```
plot(wisc.pr$x[,1], wisc.pr$x[,2], col=diagnosis,
     xlab = "PC1", ylab = "PC2")
```



Q8. Generate a similar plot for principal components 1 and 3. What do you notice about these plots?

```
plot(wisc.pr$x[,1], wisc.pr$x[,3], col=diagnosis,  
     xlab = "PC1", ylab = "PC3")
```




The separation between the two groups are not as clean as the plot vs. PC2, since PC3 represents much less variance in the data. There is some overlap with the red and black points.

Make a ggplot version of this score plot

```
pc <- as.data.frame(wisc.pr$x)

ggplot(pc) +
  aes(PC1, PC2, col = diagnosis) +
  geom_point()
```




Variance

```
pr.var <- wisc.pr$sdev^2
head(pr.var)
```


```
[1] 13.281608  5.691355  2.817949  1.980640  1.648731  1.207357
```

```
# Variance explained by each principal component: pve
pve <- pr.var / sum(pr.var)

# Plot variance explained for each principal component
plot(pve, xlab = "Principal Component",
     ylab = "Proportion of Variance Explained",
     ylim = c(0, 1), type = "o")
```



```
barplot(pve, ylab = "Precent of Variance Explained",
        names.arg=paste0("PC",1:length(pve)), las=2, axes = FALSE)
axis(2, at=pve, labels=round(pve,2)*100 )
```



Q9. For the first principal component, what is the component of the loading vector (i.e. `wisc.pr$rotation[,1]`) for the feature `concave.points_mean`?

```
wisc.pr$rotation["concave.points_mean", 1]
```

```
[1] -0.2608538
```

Q10. What is the minimum number of principal components required to explain 80% of the variance of the data?

```
summary(wisc.pr)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

Based on the cumulative proportion, up to PC5 represents at least 80% of the data's variance.


Hierarchical clustering


```
data.scaled <- scale(wisc.data)
data.dist <- dist(data.scaled)
wisc.hclust <- hclust(data.dist, method = "complete")
```

Q11. Using the `plot()` and `abline()` functions, what is the height at which the clustering model has 4 clusters?

```
plot(wisc.hclust)
abline(h=19, col="red", lty=2)
```

Cluster Dendrogram



```
data.dist
hclust (*, "complete")
```

Selecting number of clusters

```
wisc.hclust.clusters <- cutree(wisc.hclust, h=19)
# Compare the cluster membership to the actual diagnoses.
table(wisc.hclust.clusters, diagnosis)
```

```
          diagnosis
wisc.hclust.clusters  B  M
1      12 165
```

2	2	5
3	343	40
4	0	2

Q12. Can you find a better cluster vs diagnoses match by cutting into a different number of clusters between 2 and 10?

```
table(cutree(wisc.hclust, h=4), diagnosis)
```

	diagnosis	
	B	M
1	0	1
2	0	6
3	0	7
4	0	1
5	0	3
6	0	4
7	0	6
8	0	3
9	0	1
10	4	9
11	0	4
12	0	1
13	0	3
14	0	2
15	0	3
16	0	3
17	0	4
18	21	0
19	7	1
20	7	0
21	0	1
22	0	2
23	0	1
24	0	5
25	1	4
26	0	6
27	0	2
28	3	1
29	5	0
30	0	1
31	1	3

32	5	2
33	3	1
34	0	1
35	1	8
36	1	4
37	6	0
38	15	0
39	10	0
40	15	0
41	12	0
42	0	3
43	12	0
44	3	0
45	4	0
46	4	0
47	0	2
48	1	0
49	7	0
50	1	0
51	28	0
52	1	0
53	1	0
54	0	1
55	0	1
56	0	1
57	0	1
58	0	3
59	12	0
60	3	0
61	0	9
62	0	4
63	8	2
64	2	0
65	6	0
66	0	2
67	0	1
68	3	0
69	6	0
70	1	0
71	1	0
72	0	2
73	0	1
74	0	1

75	4	0
76	0	3
77	8	0
78	7	1
79	3	1
80	3	1
81	0	1
82	3	0
83	7	0
84	2	0
85	0	1
86	5	0
87	2	0
88	1	0
89	1	1
90	20	0
91	5	0
92	0	1
93	0	2
94	0	2
95	0	1
96	2	0
97	1	0
98	4	0
99	0	2
100	0	2
101	2	0
102	0	5
103	0	1
104	1	0
105	1	0
106	0	2
107	0	2
108	0	1
109	0	1
110	4	0
111	0	2
112	0	1
113	0	1
114	0	1
115	0	1
116	2	0
117	0	3

118	0	4
119	2	0
120	0	1
121	2	0
122	3	0
123	0	3
124	0	1
125	0	1
126	0	1
127	0	1
128	0	2
129	3	0
130	1	0
131	1	0
132	3	0
133	3	1
134	8	0
135	1	0
136	2	0
137	0	1
138	0	1
139	0	1
140	0	1
141	0	2
142	0	1
143	1	0
144	1	0
145	0	1
146	1	0
147	0	1
148	0	2
149	1	0
150	0	1
151	3	0
152	0	1
153	0	1
154	2	0
155	0	1
156	1	0
157	2	0
158	1	0
159	1	0
160	0	1

161	1	0
162	1	0
163	1	0
164	1	0
165	0	1
166	0	1
167	0	1
168	1	0


Cutting at around $h=4$ creates a better separation. The goal is to have smaller clusters where most of the samples are of one type, with minimal amounts of the other type.

Q13. Which method gives your favorite results for the same data.dist dataset?
Explain your reasoning.

ward.D2 clustering seems to give the cleanest cutoff on the dendrogram.


```
complete <- hclust(data.dist, method = "complete")
single <- hclust(data.dist, method = "single")
average <- hclust(data.dist, method = "average")
ward.D2 <- hclust(data.dist, method = "ward.D2")

plot(complete)
```



```
plot(single)
```


Cluster Dendrogram



```
data.dist  
hclust (*, "single")
```


```
plot(average)
```

Cluster Dendrogram



```
data.dist  
hclust (*, "average")
```

```
plot(ward.D2)
```




```
data.dist  
hclust (*, "ward.D2")
```

Clustering in PC space

```
hc <- hclust(dist(wisc.pr$x[,1:2]), method = "ward.D2")  
  
plot(hc)  
abline(h = 70, col = "red")
```


Cluster Dendrogram



```
dist(wisc.pr$x[, 1:2])
hclust (*, "ward.D2")
```

Cluster membership vector

```
grps <- cutree(hc, h=70)
```

```
table(diagnosis)
```

```
diagnosis
  B   M
357 212
```

Cross-table to see how my clustering groups correspond to the expert diagnosis vector of M and B values

```
table(grps, diagnosis)
```

```
diagnosis
grps  B   M
  1  18 177
  2 339  35
```

Positive => Cancer M Negative => Non-cancer B

True = Cluster 1 False = Cluster 2

True positive 177 False positive 18 True negative 339 False negative 35

Q17. Which of your analysis procedures resulted in a clustering model with the best specificity? How about sensitivity?


I think that the hierarchical clustering with the ward.D2 method gave best sensitivity and the PCA gave best specificity.

We can use our PCA results `wisc.pr` to make predictions on new unseen data.

```
#url <- "new_samples.csv"
url <- "https://tinyurl.com/new-samples-CSV"
new <- read.csv(url)
npc <- predict(wisc.pr, newdata=new)
npc
```

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
[1,]	2.576616	-3.135913	1.3990492	-0.7631950	2.781648	-0.8150185	-0.3959098
[2,]	-4.754928	-3.009033	-0.1660946	-0.6052952	-1.140698	-1.2189945	0.8193031
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
[1,]	-0.2307350	0.1029569	-0.9272861	0.3411457	0.375921	0.1610764	1.187882
[2,]	-0.3307423	0.5281896	-0.4855301	0.7173233	-1.185917	0.5893856	0.303029
	PC15	PC16	PC17	PC18	PC19	PC20	
[1,]	0.3216974	-0.1743616	-0.07875393	-0.11207028	-0.08802955	-0.2495216	
[2,]	0.1299153	0.1448061	-0.40509706	0.06565549	0.25591230	-0.4289500	
	PC21	PC22	PC23	PC24	PC25	PC26	
[1,]	0.1228233	0.09358453	0.08347651	0.1223396	0.02124121	0.078884581	
[2,]	-0.1224776	0.01732146	0.06316631	-0.2338618	-0.20755948	-0.009833238	
	PC27	PC28	PC29	PC30			
[1,]	0.220199544	-0.02946023	-0.015620933	0.005269029			
[2,]	-0.001134152	0.09638361	0.002795349	-0.019015820			

```
plot(wisc.pr$x[,1:2], col=grps)
points(npc[,1], npc[,2], col="blue", pch=16, cex=3)
text(npc[,1], npc[,2], c(1,2), col="white")
```



Q18. Which of these new patients should we prioritize for follow up based on your results?

Patient 2, because their sample is similar to previous malignant samples colored red.