MAT 3004 - Abstract Algebra I

Daniel Wong

Contents

Chapter 1

Preliminaries

In this Chapter, we will recall some knowledge which will be assumed throughout the course.

1.1 Divisibility of Integers

We begin our studies on the integers $\mathbb{Z}$. We say a nonzero integer $t \in \mathbb{Z}$ **divides** (or **is a factor of**) an integer $s \in \mathbb{Z}$ if there exists $u \in \mathbb{Z}$ such that $s = tu$. In this case, we write

$$t \mid s$$

If not, write $t \nmid s$. A prime $p \in \mathbb{Z}$ is a positive integer greater than 1 whose only positive divisors are 1 and p.

Theorem 1.1 (Division Algorithm). *Let $a, b \in \mathbb{Z}$ with $b > 0$. Then there exists unique integers q and r with the property that $a = bq + r$, where $0 \leq r < b$.*

Proof. We begin with the existence portion of the theorem. Consider the set

$$S = \{a - bk \mid k \in \mathbb{Z}, \ a - bk \geq 0\}.$$

Then S is obviously nonempty, since for $a \geq 0$ then $a = a - b \cdot 0 \in S$, while for $a < 0$, one has $a - b\,(2a) = a\,(1 - 2b) \in S$. By the well-ordering property of $\mathbb{Z}_{\geq 0}$, there must be a minimal element $r \in S$.

If $r = 0 \in S$, then $b \mid a$, and we may take $q = a/b$ and $r = 0$. Otherwise, $r = a - bq > 0$ and one has $a = bq + r$, and it remains prove that $r < b$: Suppose on contrary that $r \geq b$. Then

$$a - b\,(q + 1) = a - bq - b = r - b \geq 0$$

so that $a - b\,(q + 1) < a - bq = r$ is in S, contradicting the minimality of r.

The proof of uniqueness of q and r is left as an exercise. $\qquad\square$

Definition 1.2. The **greatest common divisor (gcd)** of two nonzero integers a and b is the largest of all common divisors of a and b. We denote this integer by $\gcd(a, b)$.

If $\gcd(a, b) = 1$, we say a and b are **relatively prime**.

Remark 1.3. One can also define the gcd of $a_1, a_2, \ldots, a_k$ as the largest of common divisors of all a_i's. Then one has

$$\gcd(a_1, \ldots, a_k) = \gcd(\gcd(a_1, a_2), a_3, \ldots, a_k) = \gcd(\gcd(\gcd(a_1, a_2), a_3), \ldots, a_k).$$

Similarly, if $\gcd(a_1, \ldots, a_k) = 1$, we say $a_1, \ldots, a_k$ are **relatively prime**. Note that $a_1, \ldots, a_k$ are relatively prime does not necessarily mean $\gcd(a_i, a_j) = 1$ for any $i \neq j$. For instance,

$$\gcd(6, 10, 15) = 1$$

but $\gcd(6, 10) = 2$, $\gcd(6, 15) = 3$, $\gcd(10, 15) = 5$.

1.2 Euclidean algorithm and Bézout's Theorem

To find the greatest common divisor of two integers, one can apply Euclidean algorithm:

Example 1.4. To find $\gcd(374, 221)$, one can keep dividing the larger integer by the smaller integer:

$$374 = 1 \cdot 221 + 153$$
$$221 = 1 \cdot 153 + 68$$
$$153 = 2 \cdot 68 + 17$$
$$68 = 4 \cdot 17 + 0$$

when the remainder is zero, the last remainder to it (colored in red) is the greatest common divisor. Therefore,

$$\gcd(374, 221) = 17$$

Theorem 1.5 (Bézout's Theorem)**.** *Suppose* $\gcd(p, q) = m$. *Then there exists integers* $a, b \in \mathbb{Z}$ *such that*

$$m = a \cdot p + b \cdot q \quad \text{for some integers } a, b$$

Example 1.6. By Bezout's Theorem, there exists a, b integers such that

$$374a + 221b = 17$$

To find these integers, one do backward induction from the bottom equality to the top:

From the bottom equality:

$$17 = 153 - 2 \cdot 68$$

Substitute the second equality $(68 = 221 - 1 \cdot 153)$ to the above equation:

$$17 = 153 - 2(221 - 1 \cdot 153) = 3 \cdot 153 - 2 \cdot 221$$

Now substitute the first equality $(153 = 374 - 1 \cdot 221)$ to the above equation:

$$17 = 3(374 - 221) - 2 \cdot 221$$
$$= 3 \cdot 374 - 3 \cdot 221 - 2 \cdot 221$$
$$17 = 3 \cdot 374 - 5 \cdot 221$$

Thus:

$$a = 3, \quad b = -5$$

In the special case of when a and b are relatively prime, there exist integers s and t such that

$$as + bt = 1.$$

This plays an important role in algebra. More generally, for $a_1, \ldots, a_k \in \mathbb{Z}$, one has

$$a_1 s_1 + \cdots + a_k s_k = \gcd(a_1, \ldots, a_k)$$

for some integers $s_1, \ldots, s_k$. This can be checked by using the fact that $\gcd(a_1, a_2, a_3 \ldots, a_k) = \gcd(\cdots \gcd(\gcd(a_1, a_2), a_3), \ldots, a_k)$ and apply the above algorithms repeatedly.

Finally, we state a standard yet important theorem for natural numbers, and is slightly extended to all integers:

Theorem 1.7 (Fundamental Theorem of Airthmetic)**.** *Every nonzero integer $n \in \mathbb{Z}$ can be factorized into a product of primes up to ± 1, i.e. $n = p_1 p_2 \cdots p_r$ if $n > 0$ or $n = -p_1 p_2 \cdots p_r$ if $n < 0$. Moreover,this product is unique up to ± 1 and the ordering of factors. That is, if*

$$n = p_1 p_2 \cdots p_r = q_1 q_2 \cdots q_s,$$

where the p's and q's are primes, then $r = s$ and, after renumbering the q's, we have $p_i = \pm q_i$ for all i.

1.3 Modular Arithmetic $\mathbb{Z}_n$

Sometimes, we would like to study integers that has a certain 'cycle'. For instance, days in a week has a cycle of 7, months in a year has a cycle of 12, hours in a day has a cycle of 24 and so on. We define the *congruence numbers*:

$$\mathbb{Z}_n := \{0 \ (\mathrm{mod} \ n), \ 1 \ (\mathrm{mod} \ n), \ \ldots, (n-1) \ (\mathrm{mod} \ n)\} = \{[0]_n, [1]_n, \ldots, [n-1]_n\}$$

For example, when $n = 7$, it corresponds to days in a week, and when $n = 12$ it corresponds to months in a year.

The addition and multiplication rule in $\mathbb{Z}_n$ is given by:

$$[a] + [b] = [\text{remainder of } a + b \text{ in } n]$$
$$[a] \cdot [b] = [\text{remainder of } ab \text{ in } n]$$

(we may sometimes omit the subscript $[\ ]_n$ if it is clear from the context). For instance, when $n = 7$:

$$[4]_7 \cdot [5]_7 = [4 \cdot 5]_7 = [20]_7 = [6]_7$$

There are many applications on modular arithmetic - for instance, RSA cryptography, fast Fourier transform, etc. It can also be used to verify the validity of statements about divisibility regarding all positive integers by checking only finitely many cases.

Example 1.8. To see
$$[n^3 + (n+1)^3 + (n+2)^3]_9 \equiv [0]_9$$

for all integers $n \in \mathbb{Z}$, one only needs to check this holds for $n = 0, 1, \ldots, 8$.

1.4   Equivalence Relations

One has encountered the notion of 'equivalence' in different contexts within or outside of mathematics. This allows us to relate different objects that are not necessarily identical to each other.

For instance, one studies 'similar triangles' in middle school mathematics, where two non-identical triangles $\Delta_1 \sim \Delta_2$ are similar if one can resize one triangle to get another. Another example is the congruence numbers we studied in the previous relation, where $1 \neq 8 \neq 15 \neq 22 \neq 29$ but $[1] = [8] = [15] = [22] = [29]$ in $\mathbb{Z}_7$.

Here is a precise definition of what it means for two elements to be equivalent:

Definition 1.9 (Equivalence Relation)**.** An equivalence relation on a set S is a subset $R \subseteq S \times S$ such that:

1. $(a, a) \in R$ for all $a \in S$ (reflexivity).
2. $(a, b) \in R$ implies $(b, a) \in R$ (symmetry).
3. $(a, b) \in R$ and $(b, c) \in R$ imply $(a, c) \in R$ (transitivity).

We will write $a \sim b$ instead of $(a, b) \in R$.

If $\sim$ is an equivalence relation on a set S and $a \in S$, then the **equivalence class containing/with representative** a is the subset:

$$[a] = \{x \in S \mid x \sim a\}.$$

The collection of equivalence classes of S is denoted as:

$$S/\sim := \{[a] \mid a \in S\}$$

Example 1.10.   1. Let S be the set of all triangles in a plane. If $a, b \in S$, define $a \sim b$ if a and b are similar-that is, if a and b have corresponding angles that are the same. Then $\sim$ is an equivalence relation on S.

Let $S = \mathbb{Z}$ be the set of integers. We define $a \sim b$ if $a \equiv b \bmod 7$, or equivalently $[a]_7 = [b]_7$. Then one can show that $\sim$ is an equivalence relation, and the equivalence class containing 1 is

$$[1] = \{\cdots, -13, -6, 1, 8, 15, \cdots\} = [8] = [15]$$

This justifies our notation of using $[\ ]_n$ for the elements in $\mathbb{Z}_7$.

One important aspect of equivalence relation on S is that it gives a **partition** of S. Namely, a partition of S is a disjoint union of nonempty subsets $P_i \subseteq S$ such that

$$\bigsqcup_{i \in I} P_i = S$$

Theorem 1.11. *Let $\sim$ be an equivalence relation on a set S. Then the collection of equivalence classes constitute a partition of S.*

Conversely, for any partition P_i of S, there is an equivalence relation on S whose equivalence classes are precisely the elements of P_i.

Proof. Let $\sim$ be an equivalence relation on a set S. For any $a \in S$, $a \in [a]$ since $a \sim a$. Therefore, $[a]$ is nonempty, and $\bigcup_{a \in S} [a] = S$.

There are repetitions in the union above. One therefore has to show that for $a, b \in S$, one either has $[a] = [b]$ or $[a] \cap [b] = \emptyset$ is disjoint.

To do so, suppose $c \in [a] \cap [b]$, so that $c \sim a$ and $c \sim b$. By symmetry and transitivity, one therefore has $a \sim b$. Then for any $x \in [a]$, one has $x \sim a$ (and $a \sim b$). So $x \sim b$ and hence $x \in [b]$. In other words,

$$[a] \subseteq [b].$$

The above argument can be reversed, so that one has $[b] \subseteq [a]$ as well. Thus, $[a] = [b]$.

To prove the converse, let $\{P_i \mid i \in I\}$ be a partition of S. Define $a \sim b$ if $a, b \in P_i$ for some $i \in I$. One then can easily check that $\sim$ is an equivalence relation on S. $\square$

1.5   Functions

Definition 1.12. A **function** $\phi : A \to B$ from a set A to a set B is a rule that assigns to each element $a \in A$ exactly one element $\phi(a) \in B$. The set A is called the **domain** of ϕ, and B is called the **codomain** of ϕ.

When we say ϕ is **well-defined**, we need to show the following:

- For all $a \in A$, $\phi(a) \in B$; and

- If $a = a' \in A$, then $\phi(a) = \phi(a') \in B$.

The above definition of well-definedness may look trivial at the first sight, but this may become an issue when one has two or more 'representatives' of the same element in A. For instance, a map $\phi : \mathbb{Z}_7 \to \mathbb{Z}$ given by $\phi([a]_7) = a$ is *not* well-defined, since $[1]_7 = [8]_7$ but $\phi([1]_7) = 1 \neq 8 = \phi([8]_7)$.

Definition 1.13. A function $\phi : A \to B$ is called

- injective (or one-to-one) if for every $a, a' \in A$, $\phi(a) = \phi(a')$ implies $a = a'$.

- surjective (or onto) if for any $b \in B$, there exists $a \in A$ such that $f(a) = b$.

- bijective if it is both injective and surjective.

In the case when $\phi : A \to B$ is bijective, one says that ϕ is **invertible**, and has an inverse $\psi^{-1} : B \to A$ (often denoted as ϕ^{-1}) such that

$$\psi \circ \phi = \mathrm{id}_A, \quad \phi \circ \psi = \mathrm{id}_B$$

where $\mathrm{id}_S : S \to S$ is the identity map $\mathrm{id}_S(s) := s$ for all $s \in S$.

1.6   Polynomials

In this section, we will introduce some basic aspects of polynomials over a field $\mathbb{F}$. For beginners, it is safe to assume $\mathbb{F} = \mathbb{R}$ or $\mathbb{C}$.

Definition 1.14.   1. A polynomial over $\mathbb{F}$ has the form

$$p(z) = a_m z^m + \cdots + a_1 z + a_0, \quad (a_m \neq 0).$$

Here $a_m z^m$ is called the **leading term** of $p(z)$; m is called the degree; a_m is called the **leading coefficient**; $a_m, \ldots, a_0$ are called the coefficients of this polynomial.

2. A polynomial over $\mathbb{F}$ is **monic** if its leading coefficient is $1_{\mathbb{F}}$.

3. A polynomial $p(z) \in \mathbb{F}[z]$ is **irreducible** if for any $a(z), b(z) \in \mathbb{F}[z]$,

$$p(z) = a(z)b(z) \implies \text{ either } a(z) \text{ or } b(z) \text{ is a constant polynomial.}$$

Otherwise $p(z)$ is **reducible**.

Example 1.15. The polynomial $p(x) = x^2 + 1$ is irreducible over $\mathbb{R}[x]$; but $p(x) = (x - i)(x + i)$ is **reducible** in $\mathbb{C}[x]$.

Theorem 1.16. *Division Algorithm For all $p, q \in \mathbb{F}[z]$ such that $p \neq 0$, there exists unique $s, r \in \mathbb{F}[x]$ satisfying $\deg(r) < \deg(q)$, such that*

$$p(z) = s(z) \cdot q(z) + r(z).$$

*Here $r(z)$ is called the **remainder**.*

Theorem 1.17 (Root Theorem). *For $p(x) \in \mathbb{F}[x]$, and $\lambda \in \mathbb{F}$, $x - \lambda$ divides $p(x)$ if and only if $p(\lambda) = 0$.*

Proof. 1. If $(x - \lambda)$ divides p, then $p(x) = (x - \lambda)q(x)$ for some $q(x) \in \mathbb{F}[x]$. Thus clearly $p(\lambda) = 0$.

2. For the other direction, suppose that $p(\lambda) = 0$. By division theorem, there exists $q(x), r(x) \in \mathbb{F}[x]$ such that

$$p(x) = (x - \lambda)q(x) + r(x) \quad \text{with } \deg(r(x)) < \deg(x - \lambda) = 1. \tag{1.1}$$

Therefore, $r(x) = r$ must be a constant polynomial. Substituting λ into both sides in the above equation, we have

$$0 = p(\lambda) = 0 \cdot q(x) + r \implies r = 0.$$

Therefore, $p = (x - \lambda) \cdot q(x)$, i.e., $(x - \lambda)$ divides $p(x)$. $\qquad \square$

Corollary 1.18. *A polynomial with degree n has at most n roots counting multiplicity.*

Definition 1.19 (Algebraically Closed). A field $\mathbb{F}$ is called **algebraically closed** if every non-constant polynomial $p(x) \in \mathbb{F}[x]$ has a root $\lambda \in \mathbb{F}$, or equivalently, all polynomials in $p(x) \in \mathbb{F}[x]$ can be factorized into linear terms:

$$p(x) = c(x - \lambda_1) \cdots (x - \lambda_n)$$

for $c, \lambda_1, \ldots, \lambda_n \in \mathbb{F}$.

We have seen before that $\mathbb{R}$ is not algebraically closed. Nevertheless, we have:

Theorem 1.20 (Fundamental Theorem of Algebra). $\mathbb{C}$ *is algebraically closed.*

We will skip the proof of the theorem. This can be proved using complex **analysis** (MAT 3253); or **topology** of S^1 (MAT 4002); or **algebra**ic number theory (MAT 5210).

In general, $\mathbb{F}$ may not necessarily be factorized into linear terms. But the factorization is still unique. This can be seen as an analogue of the fundamental theory of arithmetic in $\mathbb{Z}$:

Theorem 1.21 (Unique Factorization). *Every $f(x) = a_n x^n + \cdots + a_0$ in $\mathbb{F}[x]$ can be factorized as*

$$f(x) = a_n [p_1(x)]^{e_1} \cdots [p_k(x)]^{e_k}$$

*where p_i's are **monic, irreducible, distinct**. Furthermore, this expression is unique up to the permutation of factors.*

This will be proved in the chapter of Ring Theory. Assuming its validity for the moment, we can now define:

Definition 1.22 (Factor). If $p(x) = q(x)s(x)$ with $p, q, s \in \mathbb{F}[x]$, then we say

- $p(x)$ is **divisible** by $s(x)$;

- $s(x)$ is a **factor** of $p(x)$;

- $s(x)|p(x)$;

- $s(x)$ **divides** $p(x)$;

- $p(x)$ is **multiple** of $s(x)$.

Definition 1.23 (Common Factor).   1. The polynomial $g(x)$ is said to be a **common factor** of $f_1, \ldots, f_k \in \mathbb{F}[x]$ if

$$g | f_i, \quad i = 1, \ldots, k$$

2. The polynomial $g(x)$ is said to be a **greatest common divisor** of $f_1, \ldots, f_k$ if

  - g is **monic**.

  - g is common factor of $f_1, \ldots, f_k$

  - g is of largest possible (maximal) degree.

The $\gcd(f_1, f_2)$ is easy to compute for factorized polynomials. For example, let $f_1(x) = (x^2 + x + 1)^3 (x - 3)^2 x^4$ and $f_2(x) = (x^2 + 1)(x - 3)^4 x^2$ in $\mathbb{R}[x]$, then

$$\gcd(f_1, f_2) = (x - 3)^2 x^2$$

As for general polynomials, the gcd can be computed using Euclidean algorithm: For example, given $x^3 + 6x + 7$ and $x^2 + 3x + 2$, we imply

$$x^3 + 6x + 7 = (x - 3)(x^2 + 3x + 2) + (13x + 13)$$

and

$$x^2 + 3x + 2 = \frac{x + 2}{13}(13x + 13) + 0$$

Therefore, $\gcd(x^2 + 3x + 2, 13x + 13)$ is equal to a scalar multiple of $13x + 13$ such that it is monic, namely

$$\gcd(x^3 + 6x + 7, x^2 + 3x + 2) = x + 1.$$

Similarly, one has Bezout's theorem for polynomials:

Theorem 1.24 (Bezout)**.** *Let* $g = \gcd(f_1, f_2)$*, then there exists* $r_1, r_2 \in \mathbb{F}[x]$ *such that*

$$g(x) = r_1(x)f_1(x) + r_2(x)f_2(x)$$

More generally, $g = \gcd(f_1, \ldots, f_k)$ *implies there exists* $r_1, \ldots, r_k$ *such that*

$$g = r_1 f_1 + \cdots + r_k f_k$$

1.7 Introduction to Abstract Algebra

We now give a brief introduction of abstract algebra - in a nutshell, abstract algebra is about generalization of number systems we studied in kindergarten, such as:

$$\mathbb{Z} = \{\ldots, -1, 0, 1, \ldots\}$$
$$\mathbb{Q} = \left\{\frac{m}{n} \;\middle|\; m \in \mathbb{Z}, \; n \in \mathbb{Z} \setminus \{0\}\right\}$$
$$\mathbb{R} = \text{real numbers (limits of Cauchy sequences in } \mathbb{Q})$$
$$\mathbb{C} = \{x + iy \mid x, y \in \mathbb{R}\}$$

All these number systems have addition and multiplication, e.g. in $\mathbb{Q}$:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}, \quad \frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

And they all possess nice properties, e.g.,

$$(a + b) + c = a + (b + c) \tag{A1}$$
$$(ab)c = a(bc) \tag{A2}$$
$$ab = ba \tag{C1}$$
$$a + b = b + a \tag{C2}$$
$$a(b + c) = ab + ac \tag{D1}$$
$$(a + b)c = ac + bc \tag{D2}$$

One goal in abstract algebra is to study different number systems, and to find out their common features and obtain theorems that hold for all such generalized number systems.

Example 1.25.

$$M_{2\times 2}(\mathbb{R}) = \left\{\begin{pmatrix} a & b \\ c & d \end{pmatrix} \;\middle|\; a, b, c, d \in \mathbb{R}\right\}$$

We can still do $+$ and $\times$ on $M_{2\times 2}(\mathbb{R})$, but we no longer have (C2) - in other words,

$$AB \neq BA \quad \text{in general for matrices.}$$

Chapter 2

Groups

2.1 Basic Definitions

Definition 2.1. Let S be a set. A *binary operation* on S is a map:

$$* : S \times S \to S$$

Let $T \subseteq S$ be a subset. We say that the binary operation is *closed* in T if:

$$\forall a, b \in T, \quad a * b \in T$$

Example 2.2. Let $S = \mathbb{Z}$. Then the following are binary operations:

- $S = \mathbb{Z}, \quad * = +$ (addition)
- $S = \mathbb{Z}, \quad * = \times$ (multiplication)

Let

$$T = \{\text{all even integers}\} \subset \mathbb{Z}.$$

For $a, b \in T$, $a + b \in T$ (the sum of two even numbers is even). Hence $(T, +)$ is closed in $(S, +)$. Also, if $a, b \in T$, $a \cdot b \in T$ (the product of two even numbers is even). Therefore, $(T, \times)$ is also closed in $(S, \times)$.

However, let

$$T' = \{\text{all odd integers}\}$$

Then $(T', +)$ is **not** closed in $(S, +)$. But for any $(2p + 1), (2q + 1) \in T'$:

$$(2p + 1) \cdot (2q + 1) = 4pq + 2p + 2q + 1 = 2(2pq + p + q) + 1 \in T'$$

Therefore, $(T', \times)$ is closed in $(S, \times)$.

Example 2.3.   1. Let $S = \mathbb{R}^n$, and $* = $ be the vector addition operation. If $W \leq \mathbb{R}^n$ is a vector subspace, then $(W, +)$ is closed in $(\mathbb{R}^n, +)$.

2. Let $S = M_{n \times n}(\mathbb{R})$, and $* =$ multiplication of the ne matrix. Suppose $T = GL_n(\mathbb{R})$ is the subset of all invertible real matrices. To check if T is closed in $(S, \times)$: If $A, B \in T$ (invertible matrices), then $A \cdot B$ is invertible since

$$(AB)^{-1} = B^{-1}A^{-1}$$

or alternatively:

$$\det(AB) = \det(A)\det(B) \neq 0$$

Hence, T is closed in $(S, \times)$.

Definition 2.4 (Group). A **group** G is a set along with a binary operation $* : G \times G \to G$ satisfying:

1. $(a * b) * c = a * (b * c) \quad \forall a, b, c \in G \quad$ (associativity)

2. $\exists\, e \in G$ such that $e * a = a * e = a, \ \forall a \in G \quad$ (identity)

3. $\forall a \in G, \ \exists b \in G$ s.t. $a * b = b * a = e$ (b is often written as a^{-1}, called the inverse of a)

Example 2.5.   (a) $(\mathbb{Z}, +)$ is a group. To see this, note that

$$\begin{cases} (a + b) + c = a + (b + c) \\ 0 + a = a + 0 = a \quad \forall a \in \mathbb{Z} \quad (\text{i.e., } e = 0) \\ a + (-a) = (-a) + a = 0 \quad (\text{i.e., } a^{-1} = -a) \end{cases}$$

(b) Similarly $(\mathbb{Q}, +), (\mathbb{R}, +), (\mathbb{C}, +)$ as well as $(\mathbb{Z}_n, +)$ are groups.

(c) $(\mathbb{Z}, -)$ is **not** a group, since $(a - b) - c \neq a - (b - c)$ and $0 - a \neq a - 0 = a$ in general, i.e. it is not associative.

(d) $(\mathbb{Z}, \times)$ is also **not** a group. Although $(a \times b) \times c = a \times (b \times c)$ is associative, and $a \times 1 = 1 \times a = a$ (so that the identity can be taken as $e = 1$), but there is *no* $a \in \mathbb{Z}$ such that $2 \times a = 1 = e$.)

(e) $(\mathbb{Q}, \times)$ is **not** a group. Although now one can take $a = \frac{1}{2}$ such that $2 \times \frac{1}{2} = 1$, yet the element $0 \in \mathbb{Q}$ does not have a multiplicative inverse.

(f) To resolve the issue, let $\mathbb{Q}^* = \{q \in \mathbb{Q} \mid q \neq 0\}$. Then $(\mathbb{Q}^*, \times)$ is a group.

(g) Similarly, $(\mathbb{R}^*, \times)$ and $(\mathbb{C}^*, \times)$ are groups.

(h) The set of 2×2 real matrices, denoted $M_2(\mathbb{R})$, under matrix multiplication $(M_2(\mathbb{R}), \times)$ is **not** a group, because not all matrices have a multiplicative inverse (namely, singular matrices do not have a multiplicative inverse).

(i) Let $GL(2, \mathbb{R}) = \{A \in M_2(\mathbb{R}) \mid \det(A) \neq 0\}$ be the set of 2×2 invertible matrices with real entries. Then $(GL(2, \mathbb{R}), \times)$ is a group, known as the general linear group of degree 2 over $\mathbb{R}$.

Question 2.6. *Given $n \in \mathbb{N}$, for which $a \in \mathbb{Z}$ does $[a]_n$ (the congruence class of a modulo n) have a multiplicative inverse in $\mathbb{Z}_n$?*

Answer: An element $[a]_n \in \mathbb{Z}_n$ has a multiplicative inverse if and only if $\gcd(a, n) = 1$. For instance, when $n = 6$, then:

- For $a = 3$, $\gcd(3, 6) = 3 \neq 1$. Therefore, $[3]_6$ does not have a multiplicative inverse in $\mathbb{Z}_6$.

- For $a = 5$, $\gcd(5, 6) = 1$. Therefore, $[5]_6$ has a multiplicative inverse in $\mathbb{Z}_6$. Indeed, $[5]_6 \cdot [5]_6 = [25]_6 \equiv [1]_6 \pmod 6$, so $[5]_6^{-1} = [5]_6$.

Remark 2.7.   1. When there is no ambiguity, we often write ab instead of $a * b$.

2. The group operation $*$ may not be commutative.

Definition 2.8 (Abelian Group)**.** A group $(G, *)$ is called *abelian* if its operation is commutative; that is, for all $a, b \in G$, we have $a * b = b * a$.

Example 2.9.   • Examples of abelian groups include: $(\mathbb{Z}, +)$, $(\mathbb{Q}, +)$, $(\mathbb{R}, +)$, $(\mathbb{C}, +)$, and $(\mathbb{Z}_n, +)$.

- The group $(GL(2, \mathbb{R}), \times)$ is NOT abelian, as matrix multiplication is generally not commutative.

Proposition 2.10. *Let $(G, *)$ be a group.*

1. **Uniqueness of Identity:** *The identity element e in G is unique.*

2. **Uniqueness of Inverses:** *For each $a \in G$, its inverse a^{-1} is unique.*

3. **Cancellation Laws:** *For $a, b, c \in G$:*

   - *If $a * b = a * c$, then $b = c$ (Left Cancellation).*

   - *If $b * a = c * a$, then $b = c$ (Right Cancellation).*

Finite Groupsfi and Cayley Tables

Definition 2.11 (Order of a Group)**.** The *order* of a group G, denoted $|G|$, is the number of elements in the group. If $|G| < \infty$, the group is called a *finite group*.

Definition 2.12 (Cayley Table)**.** For a finite group $(G, *)$, a *Cayley table* (or group table) is a square table where the rows and columns are labeled by the elements of the group. The entry in the row corresponding to a and the column corresponding to b is $a * b$.

Proposition 2.13. *In a Cayley table for a finite group, each element of the group appears exactly once in each row and exactly once in each column.*

   Warning: If you have a table, you can check if it forms a group by verifying the group axioms.

Example 2.14. Consider a group G with $|G| = 4$. For instance, $(\mathbb{Z}_4, +)$ would have a Cayley table where each row and column contains $\{[0]_4, [1]_4, [2]_4, [3]_4\}$ exactly once.

Subgroups

Definition 2.15 (Subgroup)**.** Let $(G, *)$ be a group and H be a non-empty subset of G. We say that H is a *subgroup* of G, denoted $H \leq G$, if $(H, *)$ is itself a group under the same operation $*$ restricted to H.

Proposition 2.16 (Subgroup Test)**.** *Let $(G, *)$ be a group and H be a non-empty subset of G. Then H is a subgroup of G if and only if:*

1. *For all $a, b \in H$, $a * b \in H$ (Closure).*

2. *For all $a \in H$, $a^{-1} \in H$ (Existence of Inverses).*

An equivalent, more concise test (the one-step subgroup test) is:

- *For all $a, b \in H$, $a * b^{-1} \in H$.*

Example 2.17. • $(\mathbb{Z}, +)$ is a subgroup of $(\mathbb{Q}, +)$.

- If W is a vector subspace of $\mathbb{R}^n$, then W is a subgroup of $(\mathbb{R}^n, +)$.

- $(G, *) = (\mathbb{Z}, +)$. $H = $ all even integers $= 2\mathbb{Z} \subseteq G$.
  Take $2p, 2q \in H$ $(p, q \in \mathbb{Z})$. Then $2p + 2q = 2(p+q) \in H$ & $-2p = 2(-p) \in H$.
  Hence, $H \leq G$ is a subgroup of G.

- $(G, *) = (\mathbb{Z}, +)$. $K = $ all odd integers $\subseteq G$.
  K is **NOT** a subgroup, since $1, 3 \in K$ but $1 + 3 \notin K$.

- $(G, *) = (\mathbb{Z}, +)$. Then $k\mathbb{Z} := \{$multiples of $k\} \leq G$.

- Let $(G, *)$ be any group. Then:

  (i) $(G, *) \leq (G, *)$ is a subgroup of G.
      We'll say all $H \leq G$ satisfying $H \neq G$ a <u>proper subgroup</u> of G.

  (ii) $\{e\} \leq G$ is a subgroup.
      We'll say all $H \leq G$ with $H \neq \{e\}$ <u>nontrivial subgroup</u> of G.

- $(G, *) = (GL(n, \mathbb{R}), \cdot)$. $SL(n, \mathbb{R}) := \{A \in G \mid \det(A) = 1\} \leq G = GL(n, \mathbb{R})$.
  $$\begin{cases} \forall A, B \in SL(n, \mathbb{R}), \quad \det(AB) = \det(A) \cdot \det(B) = 1 \cdot 1 = 1 \Rightarrow AB \in SL(n, \mathbb{R}) \\ \forall A \in SL(n, \mathbb{R}), \qquad \det(A^{-1}) = \frac{1}{\det(A)} = \frac{1}{1} = 1 \Rightarrow A^{-1} \in SL(n, \mathbb{R}) \end{cases}$$

- $(\mathbb{Z}, +) \leq (\mathbb{Q}, +) \leq (\mathbb{R}, +) \leq (\mathbb{C}, +)$

- $(G, *) = (\mathbb{Z}_8, +)$ $H = \{0, 4\}$ is a subgroup
  $\qquad\qquad\qquad\qquad K = \{0, 3\}$ is **NOT** a subgroup.

Cyclic

Definition 2.18 (Cyclic Subgroup)**.** Let $(G, *)$ be a group. A *cyclic subgroup* generated by $g \in G$ is the subgroup $\langle g \rangle := \{g^m \mid m \in \mathbb{Z}\}$.

(Exercise: Check $\langle g \rangle \leq G$ is a subgroup of G, i.e.: for any

$$g^a, g^b \in \langle g \rangle \quad \begin{cases} g^a * g^b \in \langle g \rangle \\ (g^a)^{-1} \in \langle g \rangle \end{cases}$$

Example 2.19. • $(G, *) = (\mathbb{Z}, +)$.

$\langle 2 \rangle = \{2, 2 + 2, 2 + 2 + 2, \ldots, 0, (-2), (-2) + (-2), (-2) + (-2) + (-2), \ldots\} = 2\mathbb{Z}$

• $(G, *) = (GL(2, \mathbb{R}), \cdot)$.

$$\left\langle \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right\rangle = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^m \middle| m \in \mathbb{Z} \right\} = \left\{ \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} \middle| m \in \mathbb{Z} \right\}.$$

(Check: $\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & a + b \\ 0 & 1 \end{pmatrix}$)

• $(G, *) = (\mathbb{Z}_8, +)$.

$\langle 2 \rangle = \{0, 2, 4, 6, \ldots\} = \{0, 2, 4, 6\}$

$\langle 3 \rangle = \{0, 3, 6, 9, 12, 15, 18, 21, 24, \ldots\} = \{0, 3, 6, 1, 4, 7, 2, 5, 0, \ldots\} = G$

Definition 2.20 (Cyclic Group). Let $(G, *)$ be a group. We say G is **cyclic** if there is $g \in G$ such that $\langle g \rangle = G$.

Example 2.21. • $(G, *) = (\mathbb{Z}, +)$ is cyclic, since $G = \langle 1 \rangle$.

• $(G, *) = (\mathbb{Z}_8, +)$ is cyclic, since $G = \langle 3 \rangle = \langle 1 \rangle$.

• $(G, *) = (GL(2, \mathbb{R}), \cdot)$ is **NOT** cyclic, since for all $H = \langle g \rangle$, H is countable but G is uncountable.

Hence, H can never be equal to G.

• $(G, *) = (\mathbb{Z}_8^*, \cdot) = \{1, 3, 5, 7\}$. Then all the cyclic subgroups:

$\langle 1 \rangle = \{1\}$, $\langle 3 \rangle = \{3^0 = 1, 3^1 = 3, 3^2 = 1, 3^3 = 3, \ldots\} = \{1, 3\}$, $\langle 5 \rangle = \{1, 5\}$,

$\langle 7 \rangle = \{1, 7\}$. None of them is equal to G. So G is **not** cyclic.

(Exercise: $(\mathbb{Z}_5^*, \cdot)$ is cyclic)

Question: $(G, *)$ group, $H = \langle g \rangle$. What's the order of H?

e.g.: $(\mathbb{Z}_8, +) : |\langle 2 \rangle| = 4$, $|\langle 3 \rangle| = 8$, $|\langle 4 \rangle| = 2$.

Definition 2.22 (order). Let $g \in G$. The *order* of g is equal to the smallest positive integer k such that $g^k = e$. If no such k exists, then we say the order of g is ∞.

Proposition 2.23. *If $\text{ord}(g) = k$, then $|\langle g \rangle| = k$.*

Example 2.24. • $(G, *) = (\mathbb{Z}_8, +)$. $\text{ord}(2) = 4 : 2^1 = 2, 2^2 = 4, 2^3 = 6, 2^4 = 8 \equiv 0$;

$\text{ord}(3) = 8 : 3^1 = 3, 3^2 = 6, 3^3 = 9 \equiv 1, 3^4 = 12 \equiv 4, 3^5 = 15 \equiv 7, 3^6 = 18 \equiv 2, 3^7 = 21 \equiv 5, 3^8 = 24 \equiv 0$.

- $(G, *) = (GL(2, \mathbb{R}), \cdot)$.

  $\text{ord}\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right) = \infty$, since $\begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^m \neq \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \forall m \in \mathbb{N}$.

Remark 2.25. If $|G| < \infty$ is finite, then $\langle g \rangle \leq G$ is a finite subgroup with $|\langle g \rangle| \leq |G|$. Indeed, all $H \leq G$ has $|H| \mid |G|$ (Lagrange's Theorem).

e.g.: If $|G| = 12$, then there is **NO** $H \leq G$ with $|H| = 8$, can only be $1, 2, 3, 4, 6, 12$.

Therefore, $\text{ord}(g) \mid |G|$ (take $H = \langle g \rangle$).

2.2 More examples of Groups

Permutation group / Symmetric group S_n

Definition 2.26. A *permutation* of $X_n = \{1, 2, \ldots, n\}$ is a bijection $\sigma : X_n \to X_n$.

(e.g. $(n = 3), \sigma(1) = 2, \sigma(2) = 3, \sigma(3) = 1$)

Definition 2.27. The **symmetric / permutation group** of n elements is the collection of all permutations $\sigma : X_n \to X_n$.

$$S_n := \{\sigma : X_n \to X_n \mid \sigma \text{ is bijective}\}$$

e.g.: $S_3 = \left\{ \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \ldots, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \right\}$. Hence $|S_3| = 6$.

More generally, $|S_n| = n \cdot (n-1) \cdots 1 = n!$.

Proposition 2.28. $(S_n, \circ)$ *is a group, where $\circ$ is the composition of functions.*

Proof:

1. $(\sigma_1 \circ \sigma_2) \circ \sigma_3 = \sigma_1 \circ (\sigma_2 \circ \sigma_3)$ *is true, since $\circ$ is associative.*

2. *Take $e = id : X_n \to X_n = \begin{pmatrix} 1 & 2 & \cdots & n \\ 1 & 2 & \cdots & n \end{pmatrix}$. Then $\sigma \circ e = e \circ \sigma = \sigma$.*

3. *Since σ is bijective, then $\sigma^{-1} : X_n \to X_n$ exists with $\sigma \circ \sigma^{-1} = \sigma^{-1} \circ \sigma = e$.*

Remark 2.29. Groups are often used to study "symmetry" of objects. For example, S_n can be used to study symmetry of n objects. As another example, to describe symmetries of $\mathbb{R}^n$, we study $\{A : \mathbb{R}^n \to \mathbb{R}^n \mid A \text{ is a bijective linear transformation}\} = GL(n, \mathbb{R})$.

Calculations on S_n

Let $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 1 & 2 \end{pmatrix}$, $\tau = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 1 & 4 \end{pmatrix}$. Then:

$$\sigma \circ \tau = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 2 & 3 \end{pmatrix} \quad \text{vs} \quad \tau \circ \sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 1 & 4 & 2 \end{pmatrix}$$

Remark 2.30. $\sigma \circ \tau \neq \tau \circ \sigma$ in general).

The inverse of σ is $\sigma^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 2 & 1 \end{pmatrix}$.

Example 2.31. Let $\sigma = (1\,2\,3)$, $\tau = (1\,3\,4\,2)(7\,6)$ in S_7. Then:

- $\sigma\tau = (1\,2\,3)(1\,3\,4\,2)(7\,6) = (1)(2)(3\,4)(7\,6) = (3\,4)(6\,7)$.

- $\tau\sigma = (1\,3\,4\,2)(7\,6)(1\,2\,3) = (1)(2\,4)(3)(6\,7) = (2\,4)(6\,7)$.

Remark 2.32.   • From now on, we'll express $\sigma \in S_n$ using (disjoint) product of k-cycles.

- If α and β are k-cycles with disjoint entries, then $\alpha\beta = \beta\alpha$.
  (e.g.: $\alpha = (1\,3\,2)$, $\beta = (4\,7) \Rightarrow \alpha\beta = (1\,3\,2)(4\,7) = (4\,7)(1\,3\,2) = \beta\alpha$).
  Otherwise, $\alpha\beta \neq \beta\alpha$ in general.

- The inverse of a k-cycle is given by:

$$(i_1\, i_2 \ldots i_k)^{-1} = (i_1\, i_k\, i_{k-1} \ldots i_2)$$

  (Exercise: $(1\,2\,3\,4\,5)(1\,5\,4\,3\,2) = $ identity)

- More generally, for any $\sigma \in S_n$, $\sigma = (i_1 \ldots i_k)(j_1 \ldots j_\ell) \ldots (m_1 \ldots m_p)$ where the cycles are disjoint, then:
$$\sigma^{-1} = (i_1\, i_k \ldots i_2)(j_1\, j_\ell \ldots j_2) \ldots (m_1\, m_p \ldots m_2)$$

Definition 2.33 (Transposition). A 2-cycle $\tau = (i\, j) \in S_n$ is called a **transposition**.

Proposition 2.34. *Every $\sigma \in S_n$ can be expressed as a product of (NOT necessarily disjoint) transpositions.*

Proof. $\sigma = (i_1 \ldots i_k)(j_1 \ldots j_\ell) \ldots (m_1 \ldots m_p)$ with each $i, j, \ldots, m$ cycles are disjoint. Then for $(i_1 \ldots i_k)$:

$$(i_1 \ldots i_k) = (i_1\, i_k)(i_1\, i_{k-1}) \ldots (i_1\, i_3)(i_1\, i_2)$$

Do the same for $(j_1 \ldots j_\ell), \ldots, (m_1 \ldots m_p)$. Then we're done. $\square$ $\square$

Remark 2.35. The expression of $\sigma \in S_n$ into product of transpositions is **NOT** unique:

e.g.: $\sigma = (2\,3) = (2\,3)(2\,3)(2\,3) = (1\,2)(2\,3)(1\,3)$.

But the number of transpositions in each expression of σ is always **even** or **odd**.

Lemma 2.36. *If $e = \tau_1 \cdots \tau_k$, then $k \equiv 0 \pmod 2$ is even.*

Proof. Induction on k, i.e.: "If $e = \tau_1 \cdots \tau_k$, then k is even"   (*)

  $k = 0$ is trivial; also, it's obvious that $k \neq 1$, so (*) holds for $k = 1$.

Suppose (*) holds for all expressions of $e = \tau'_1 \cdots \tau'_{k'}$ for $k' \leq m$.

Consider $e = \tau_1 \cdots \tau_m \tau_{m+1}$, an expression of e with $(m+1)$ transpositions, with $\tau_{m+1} = (a\,b)$.

Then:

  Case 1: $\tau_m = (a\,b)$.

$\Rightarrow e = \tau_1 \cdots \tau_{m-1}(a\,b)(a\,b) = \tau_1 \cdots \tau_{m-1}$.

By induction, $m - 1 \equiv 0 \pmod 2$. Hence, $m + 1 \equiv 0 \pmod 2$ is even.

$$
\textbf{Case 2: } \tau_m \neq (a\,b). \text{ Then } \tau_m \tau_{m+1} = \begin{cases} (a\,c)(a\,b) = (a\,b)(b\,c) \\ (b\,d)(a\,b) = (a\,d)(d\,b) \\ (i\,j)(a\,b) = (a\,b)(i\,j) \end{cases} \quad \text{(where the left side has "no } a\text{")}
$$

  $\therefore e = \tau_1 \cdots \tau_{m-1}\tau_m\tau_{m+1} = \tau_1 \cdots \tau_{m-1}(a\,*)(**)$.

Then do the same Case 1, 2 for $\tau_{m-1}(a\,*)$. If $\tau_{m-1} = (a\,*)$, then $\tau_{m-1}(a\,*)$ goes away. Then induction.

Otherwise, $\tau_{m-1} \neq (a\,*)$, move one more step to the left and get:

$e = \tau_1 \cdots \tau_{m-2}(a\,*)(**)(**)$ (with "no a" on the right).

Continue until Case 1 occurs, or:

$e = (a\,\bigstar)(**) \cdots (**)$ (with "no a" on the right).

But this **cannot** happen, since the expression on right-hand-side will permute "a" to $\bigstar$. Then contradict.

Hence, Case 1 must occur somewhere. Then use induction. $\square$ $\square$

Proposition 2.37. *Let $\sigma = \tau_1 \cdots \tau_k = \tau'_1 \cdots \tau'_\ell$ be two expressions of σ as product of transpositions. Then $k \equiv \ell \pmod 2$.*

Proof. $e = \sigma^{-1}\sigma = (\tau_1 \cdots \tau_k)^{-1}(\tau'_1 \cdots \tau'_\ell) = \tau_k^{-1} \cdots \tau_1^{-1}\tau'_1 \cdots \tau'_\ell = \tau_k \cdots \tau_1 \tau'_1 \cdots \tau'_\ell$.

Then by Lemma 2.34, $k + \ell \equiv 0 \pmod 2$. $\square$ $\square$

Definition 2.38 ((Even/Odd Permutation). $\sigma \in S_n$ is called an **even / odd permutation** if σ is a product of an even / odd number of transpositions.

Definition 2.39 (Alternating Group). The **alternating group** A_n is the subgroup of S_n given by:

$$A_n := \{\sigma \in S_n \mid \sigma \text{ is even}\}$$

(Check: $A_n \leq S_n$. Also, is the set $\{\sigma \in S_n \mid \sigma \text{ is odd}\}$ a subgroup?)

Proposition 2.40. $|A_n| = \frac{1}{2}|S_n| = \frac{n!}{2}$

Proof. Let $\tau \in S_n$ be a transposition and $f : A_n \to \{\sigma \in S_n \mid \sigma \text{ odd}\}$ be defined as $f(\sigma) := \sigma\tau$, where $\sigma = \tau_1 \cdots \tau_{2m} \in A_n$.

Note that f is bijective with $f^{-1} = f$. Hence, $|A_n| = |\{\sigma \in S_n \mid \sigma \text{ odd}\}|$.

Since $A_n \cup \{\sigma \in S_n \mid \sigma \text{ odd}\} = S_n$, therefore $|A_n| = |S_n|/2 = n!/2$. $\square$ $\square$

Example 2.41 (Applications).   1. A_4 is the group of "symmetries of a tetrahedron".

- A rotation of the tetrahedron corresponds to a permutation of its 4 vertices. For example, a 120° rotation about a vertex fixies one vertex and cyclically permutes the other three:

$$(2\,3\,4) = (2\,4)(2\,3) \in A_4$$

- A 180° rotation about the midpoints of opposite edges corresponds to:

$$(1\,3)(2\,4) \in A_4$$

1'. If we allow reflections (not just rotations), we get the full symmetric group:

$$(2\,4) \in S_4 \setminus A_4$$

2. **Galois Theory**: A_5 is a simple group, so we can't solve a general degree-5 equation using $\sqrt[k]{*}, \frac{*}{*}$.

Dihedral Groups D_n

Definition 2.42 (dihedral group). D_n describes the rotational and reflectional symmetries of a regular n-gon.

- **Example ($n = 3$)**: Symmetries of an equilateral triangle.
  - Rotations: e (identity), $\sigma_{120°} \leftrightarrow (1\,2\,3)$, $\sigma_{240°} \leftrightarrow (1\,3\,2)$.
  - Reflections: 3 axes of symmetry, each fixing one vertex and swapping the other two, e.g., $\leftrightarrow (2\,3)$, $(1\,2)$, or $(1\,3)$.
  - Hence, D_3 has 6 elements.
- **Example ($n = 4$)**: Symmetries of a square.
  - Includes 4 rotations $(0°, 90°, 180°, 270°)$ and 4 reflections.
  - Hence, $|D_4| = 8$.

More generally, $|D_n| = 2n$ for all $n \geq 3$.

Algebraic Representation of n-gon Symmetries

Let r be a rotation by $\left(\frac{360}{n}\right)^{\circ}$ anticlockwise, which corresponds to the n-cycle $(1\,2\,3\ldots n)$.
Let s be a reflection along the axis passing through vertex 1.

- Then $r^n = e$ and $s^2 = e$.

- Any other reflection in D_n can be expressed as $r^k s$. This represents a reflection along an axis obtained by rotating the "principal axis" by $\left(\frac{k180}{n}\right)$ degrees anticlockwise.

- **Example D_4:**

  - $r^3 s$ represents reflecting the square across a specific axis.

  - Reflecting about an axis at $\frac{3}{4}(180°)$ results in a specific vertex permutation.

Proposition 2.43. *The dihedral group D_n is given by the set:*

$$D_n = \{e, r, r^2, \ldots, r^{n-1}, s, rs, r^2 s, \ldots, r^{n-1} s\}$$

(Exercise: Show that $sr^l = r^{n-l}s$)

Hence, $D_n = \langle s, r \mid r^n = e, s^2 = e, sr^l = r^{n-l}s \text{ for all } l \rangle$.
eg: $s^3 r^{14} s^5 r^{-3} s^{16}$ in D_5
$= sr^{10} r^4 sr^5 r^{-3} e = sr^4 sr^2 = s(sr^1)r^2 = s^2 r^3 = r^3$

Product Groups (External Direct Product)

Definition 2.44 (Product Group). Let $G_1, \ldots, G_n$ be groups. The **product group** $G := \prod_{i=1}^{n} G_i = G_1 \times \cdots \times G_n$ (in Gallian, $G_1 \oplus G_2 \oplus \cdots \oplus G_n$) is given by $G := \{(g_1, \ldots, g_n) \mid g_i \in G_i\}$ and multiplication is given by:

$$(g_1, \ldots, g_n) * (h_1, \ldots, h_n) := (g_1 h_1, g_2 h_2, \ldots, g_n h_n)$$

In particular, $|G| = |G_1| \times \cdots \times |G_n|$ & $e_G \in G$ is $(e_1, \ldots, e_n)$ where $e_i = e_{G_i}$.

e.g.: $\mathbb{Z}_2 \times \mathbb{Z}_3 = \{(a, b) \mid a = 0, 1; b = 0, 1, 2\}$
$(1, 2) * (0, 1) = (1 + 0, 2 + 1) = (1, 3) \equiv (1, 0)$.

Exercise 2.1. *1. $\mathbb{Z}_2 \times \mathbb{Z}_3 = \langle (1, 1) \rangle$ is cyclic.*

2. If $p_1, \ldots, p_n$ are distinct primes, then $\mathbb{Z}_{p_1} \times \cdots \times \mathbb{Z}_{p_n} = \langle (1, \ldots, 1) \rangle$.

More examples: $S_3 \times D_4$; $\mathbb{Z} \times \mathbb{R}$; $GL(2, \mathbb{R}) \times S_5$; $\ldots$

2.3 Homomorphism and Isomorphism

Motivation: Recall

D_4		S_4	D_4		S_4
e	$\leftrightarrow$	e	s	$\leftrightarrow$	$(2\,4)$
r	$\leftrightarrow$	$(1\,2\,3\,4)$	rs	$\leftrightarrow$	$(1\,2)(3\,4)$
r^2	$\leftrightarrow$	$(1\,3)(2\,4)$	r^2s	$\leftrightarrow$	$(1\,3)$
r^3	$\leftrightarrow$	$(1\,4\,3\,2)$	r^3s	$\leftrightarrow$	$(1\,4)(2\,3)$

To understand D_4, it's equally good to understand 8 elements in S_4.

Definition 2.45 (Homomorphism & Isomorphism)**.** Let $(G, *), (H, \circledast)$ be groups. A *homomorphism* from G to H is a map $\phi : G \to H$ s.t.: $\phi(g_1 * g_2) = \phi(g_1) \circledast \phi(g_2)$.

If ϕ is bijective, then ϕ is an *isomorphism*. (ϕ "preserves" the multiplication rule of G & H)

Example 2.46.   • $\phi : D_4 \to S_4$ given by $\phi(r) = (1\,2\,3\,4)$, $\phi(s) = (2\,4)$, ...

is an *injective homomorphism*, eg: $\phi(r^2 s) = \phi(r^2)\phi(s)$

- $\phi : \mathbb{Z}_2 \times \mathbb{Z}_2 \to K$ given by $\phi(0,0) = e, \phi(1,0) = a, \phi(0,1) = b, \phi(1,1) = c$

  Then ϕ is an isomorphism, e.g., $\phi((1,0) * (0,1)) = \phi(1,1) = c$
$$\phi(1,0) \circledast \phi(0,1) = a \circledast b = c$$
  $\therefore$ understanding (multiplication in K) $\Leftrightarrow$ understanding $\mathbb{Z}_2 \times \mathbb{Z}_2$.

- Let $(G, *) = (\mathbb{R}^n, +)$ & $(H, \circledast) = (\mathbb{R}^m, +)$. Then any linear transformation $T : \mathbb{R}^n \to \mathbb{R}^m$ is a group homomorphism, i.e.:

  $T(x_1 + x_2) = T(x_1) + T(x_2)$

- $\phi : (\mathbb{Z}_{12}, +) \to (\mathbb{Z}_4, +)$ given by $\phi(k) := k$ is **NOT** a homom.

  $\phi(2 \cdot 4) = \phi(8) = 0 \neq 2 = 2 + 0 = \phi(2) + \phi(4)$

- $\exp : (\mathbb{R}, +) \to (\mathbb{R}^*, \cdot)$ given by $\exp(a) := e^a$ is a homom.:

  $\exp(a + b) = e^{a+b} = e^a \cdot e^b = \phi(a) \cdot \phi(b)$

- Define $\phi : A_3 \to \mathbb{Z}_3$ by $\phi(e) = 0, \phi(1\,2\,3) = 1, \phi(1\,3\,2) = 2$

  ϕ is a "dictionary", translating the "A_3 language" to "$\mathbb{Z}_3$ language"
$$\begin{pmatrix} (1\,2\,3) \text{ in } A_3 \text{ language} \\ (1\,3\,2) \text{ in } A_3 \text{ language} \end{pmatrix} \xrightarrow{\phi} \begin{pmatrix} 1 \text{ in } \mathbb{Z}_3 \text{ language} \\ 2 \text{ in } \mathbb{Z}_3 \text{ language} \end{pmatrix}$$
  The dictionary also translates multiplication in A_3 to mult. in $\mathbb{Z}_3$

  $(1\,2\,3) * (1\,2\,3) = (1\,3\,2)$ in $A_3 \leftrightarrow (1 + 1 = 2$ in $\mathbb{Z}_3)$

  In other words, $\phi((1\,2\,3) * (1\,2\,3)) = \phi(1\,2\,3) + \phi(1\,2\,3)$

  $\therefore \phi : A_3 \cong \mathbb{Z}_3$ is an isomorphism.

  (Question: S_3 and $\mathbb{Z}_6$ both have 6 elements, can $S_3 \cong \mathbb{Z}_6$?)

- $i : (\mathbb{Z}, +) \hookrightarrow (\mathbb{R}, +)$ is a homom. with $i(a) := a \ \forall a \in \mathbb{Z}$.

  More generally, if $H \leq G$, then $i : (H, *) \hookrightarrow (G, *)$ is a homom.

- $\pi : (\mathbb{Z}, +) \to (\mathbb{Z}_n, +)$ with $\pi(a) := a \pmod{n}$ is a homom.

  $(\pi(a + b) = \pi(a) + \pi(b))$

- $\det : GL(n, \mathbb{R}) \to (\mathbb{R}^*, \cdot)$ is a homom.

$$\det(A \cdot B) = \det(A) \cdot \det(B)$$

- $\phi : S_n \to (\{\pm 1\}, \cdot)$ with $\phi(\sigma) := \begin{cases} +1, & \text{if } \sigma \text{ is even} \\ -1, & \text{if } \sigma \text{ is odd} \end{cases}$

  Then $\phi(\sigma\tau) = \phi(\sigma) \cdot \phi(\tau) \quad \forall \sigma, \tau \in S_n$, since:

			$\phi(\sigma\tau)$		$\phi(\sigma)\phi(\tau)$
σ even,	τ even	$\Rightarrow \sigma\tau$ even	1	$=$	$1 \cdot 1$
σ even,	τ odd	$\Rightarrow \sigma\tau$ odd	-1	$=$	$1 \cdot (-1)$
σ odd,	τ even	$\Rightarrow \sigma\tau$ odd	-1	$=$	$(-1) \cdot 1$
σ odd,	τ odd	$\Rightarrow \sigma\tau$ even	1	$=$	$(-1) \cdot (-1)$

Proposition 2.47. *Let $(G, *), (H, \circledast)$ and $(K, \star)$ be groups. Then:*

(a) *If $\phi : G \to H, \psi : H \to K$ are homom., then $\psi \circ \phi : G \to K$ is homom.*

(b) *If $\phi : G \to H$ is homom., then $\phi(e_G) = e_H$, $\phi(a^{-1}) = (\phi(a))^{-1}$.*

(c) *If $\phi : G \cong H$ is isomorphism, then $\phi^{-1} : H \to G$ satisfies $\phi^{-1}(h_1 \circledast h_2) = \phi^{-1}(h_1) * \phi^{-1}(h_2)$.*
    (In other words, the inverse of isom. is an isom.)

Proof. (a) $(\psi \circ \phi)(g_1 * g_2) = \psi(\phi(g_1 * g_2)) = \psi(\phi(g_1) \circledast \phi(g_2)) = \psi(\phi(g_1)) \star \psi(\phi(g_2))$
$= (\psi \circ \phi)(g_1) \star (\psi \circ \phi)(g_2).$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ $\square$
    (b) $\phi(g) = \phi(e_G * g) = \phi(e_G) \circledast \phi(g) \Rightarrow \phi(g)(\phi(g))^{-1} = \phi(e_G) \circledast (\phi(g) \circledast (\phi(g))^{-1})$
$\Rightarrow e_H = \phi(e_G).$
Similarly, $\phi(e_G) = \phi(a * a^{-1}) = \phi(a) \circledast \phi(a^{-1}) \Rightarrow e_H = \phi(a) \circledast \phi(a^{-1}) \Rightarrow \phi(a^{-1}) = (\phi(a))^{-1}.$ $\quad\square$
    (c) $\phi(\phi^{-1}(h_1 \circledast h_2)) = h_1 \circledast h_2 = (\phi \circ \phi^{-1}(h_1)) \circledast (\phi \circ \phi^{-1}(h_2)) = \phi(\phi^{-1}(h_1) * \phi^{-1}(h_2))$
$\Rightarrow \phi^{-1}(h_1 \circledast h_2) = \phi^{-1}(h_1) * \phi^{-1}(h_2)$, since ϕ is bijective. $\qquad\qquad \square \qquad\qquad \square$

Definition 2.48 (Kernel and Image). Let $\phi : G \to H$ be a group homomorphism.

- **kernel** of ϕ: $\ker \phi := \{g \in G \mid \phi(g) = e_H\}$

- **image** of ϕ: $\mathrm{im}\phi := \{\phi(g) \mid g \in G\}$

Example 2.49.   • $A : (\mathbb{R}^n, +) \to (\mathbb{R}^m, +)$. Then $\ker(A) = \{\mathbf{v} \in \mathbb{R}^n \mid A(\mathbf{v}) = \mathbf{0}_{\mathbb{R}^m}\} =$ Null space of A.
    $\mathrm{im}(A) = \{A(\mathbf{v}) \in \mathbb{R}^m \mid \mathbf{v} \in \mathbb{R}^n\} =$ Column space of A.

- $\pi : (\mathbb{Z}, +) \to (\mathbb{Z}_n, +)$. Then
  $\ker(\pi) = \{\text{multiples of } n\} = \langle n \rangle, \quad \mathrm{im}(\pi) = (\mathbb{Z}_n, +)$.

- $\det : GL(n, \mathbb{R}) \to (\mathbb{R}^*, \cdot)$. Then
  $\ker(\det) = \{A \in GL(n, \mathbb{R}) \mid \det(A) = 1\} =: SL(n, \mathbb{R}), \quad \mathrm{im}(\det) = \mathbb{R}^*$.

Proposition 2.50. *(a)* $\ker \phi \leq G$, $\operatorname{im} \phi \leq H$ *are subgroups of G and H respectively.*

(b) ϕ is an isomorphism $\iff \ker \phi = \{e_G\}$ & $\operatorname{im} \phi = H$.

(c) If G is cyclic / abelian, then $\phi(G)$ is cyclic / abelian.

Proof. (a) $\forall g_1, g_2 \in \ker \phi$:

   (i) $\phi(g_1 g_2) = \phi(g_1)\phi(g_2) = e_H \cdot e_H = e_H \Rightarrow g_1 g_2 \in \ker \phi$

   (ii) $\phi(g_1^{-1}) = (\phi(g_1))^{-1} \Rightarrow g_1^{-1} \in \ker \phi$

   (b) Skipped. ($\ker \phi = \{e_G\} \iff \phi$ is injective; $\operatorname{im} \phi = H \iff \phi$ is surjective)

   (c)

- If G is abelian, i.e.: $ab = ba \ \forall a, b \in G$. Then $\forall \phi(a), \phi(b) \in \phi(G)$:

$$\phi(a)\phi(b) = \phi(ab) = \phi(ba) = \phi(b)\phi(a)$$

- If G is cyclic, i.e.: $G = \langle g \rangle$ for some $g \in G$. Then $\forall \phi(a) \in \phi(G)$:

$$\phi(a) = \phi(g \cdot g \cdots g) = \phi(g) \cdot \phi(g) \cdots \phi(g) = (\phi(g))^k \in \langle \phi(g) \rangle$$

   $\Rightarrow \phi(G) \subseteq \langle \phi(g) \rangle$. Meanwhile, $\langle \phi(g) \rangle \subseteq \phi(G)$ since $\phi(g) \in \phi(G)$.
   Therefore, $\phi(G) = \langle \phi(g) \rangle$ is cyclic. $\qquad\square$

$\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\square$

Example 2.51. • $S_3, \mathbb{Z}_6$ both have 6 elements. But $S_3 \not\cong \mathbb{Z}_6$. Why? Suppose by contrary, $\phi : \mathbb{Z}_6 \xrightarrow{\cong} S_3$. Then $\phi(\mathbb{Z}_6) = S_3$ & $\phi(\mathbb{Z}_6)$ is abelian.
$\Rightarrow S_3$ is abelian, contradicting "S_3 is NOT abelian".

- $\mathbb{Z}_2 \times \mathbb{Z}_2 \not\cong \mathbb{Z}_4$. (Hint: Suppose on contrary, $\phi : \mathbb{Z}_4 \to \mathbb{Z}_2 \times \mathbb{Z}_2$ and study $\phi(x)$)

From now on, we'll classify groups up to isomorphism, i.e.: we won't distinguish $A_3 \cong \mathbb{Z}_3$ but $\mathbb{Z}_2 \times \mathbb{Z}_2$ & $\mathbb{Z}_4$ are different!

Theorem 2.52 (Classification of cyclic groups)**.** *Let $G = \langle g \rangle$ be a cyclic group. Then:*

   (a) If $|G| = \infty$, then $G \cong (\mathbb{Z}, +)$.

   (b) If $|G| = n < \infty$, then $G \cong (\mathbb{Z}_n, +)$.

Proof. (a) Consider $\phi : \mathbb{Z} \to \langle g \rangle = \{g^i \mid i \in \mathbb{Z}\} = G$ with $\phi(a) := g^a$.
Then $\phi(a + b) = g^{a+b} = g^a g^b = \phi(a)\phi(b) \Rightarrow \phi$ is a surjective homomorphism
Suppose ϕ is **NOT** injective, i.e.: $\exists \ m \neq n$, s.t.: $\phi(m) = \phi(n)$.
WLOG, assume $m > n$. Then:
$\phi(m) = \phi(n) \Leftrightarrow g^m = g^n \Rightarrow g^{m-n} = e \Rightarrow \operatorname{ord}(g) \leq m - n$ (in fact, $\operatorname{ord}(g) \mid m - n$).
Then $\langle g \rangle = \{g^0 = e, g^1, \ldots, g^{\operatorname{ord}(g)-1}\}$.
$\Rightarrow |G| = |\langle g \rangle| = \operatorname{ord}(g) < \infty$, contradicting $|G| = \infty$. $\qquad\square$

(b) Let $G = \{g^0 = e, g^1, \ldots, g^{n-1}\}$ (i.e.: $|G| = n$). Consider $\psi : G \to \mathbb{Z}_n$ with $\psi(g^i) := i$.

Then ψ is a surjective homom., since $\psi(g^a g^b) = \psi(g^{a+b}) = a + b = \psi(g^a) + \psi(g^b)$.

Since $|G| = |\mathbb{Z}_n| = n$, then any surjective map is injective also.

Therefore, $\psi : G \xrightarrow{\cong} \mathbb{Z}_n$ is bijective. $\square$ $\square$

Question (from a student): Where is negative powers of g?

Answer: $g^n = e$, so $g^{-1} = g^{-1} g^n = g^{n-1}$, $g^{-2} = g^{-2} g^n = g^{n-2} \ldots$

2.4   Lagrange's Theorem

Recall: equivalence relation $\sim$ on S:

when $x \sim y$, then x, y are in one equivalence class (with respect to $\sim$). $S = \coprod_{\alpha \in I} C_\alpha$

e.g.:

- $G = \mathbb{Z}$, $H = 3\mathbb{Z}$ on $\mathbb{Z}$, let $a \sim b \pmod 3$.

  (i) $a \sim a \pmod 3 \Rightarrow 3 \mid (a - a)$.

  (ii) $a \sim b \pmod 3 \Rightarrow b \sim a \pmod 3$ is obvious since $3 \mid (a - b)$.

  (iii) $a \sim b \pmod 3$, $b \sim c \pmod 3 \Rightarrow a - b = 3k$, $b - c = 3l \Rightarrow a - c = 3(k + l) \in 3\mathbb{Z}$.
     $\Rightarrow a \sim c \pmod 3$. $\therefore \sim$ is an equiv. relation.

  $G = \{0 + 3\mathbb{Z}\} \cup \{1 + 3\mathbb{Z}\} \cup \{2 + 3\mathbb{Z}\} = C_0 \cup C_1 \cup C_2$.

- $G = GL(n, \mathbb{R})$, $H = SL(n, \mathbb{R})$. $A \sim B$ iff $\det(A) = \det(B)$.
  (Exercise: check $\sim$ satisfies (i)-(iii)).
  $C_\alpha = [A] = \{B \in GL(n, \mathbb{R}) \mid \det(B) = \alpha\}$.

• If $a \sim a'$ in S, then $C_a = C_{a'}$. (e.g.: in the example above, $1 \sim 4$).

Therefore, any element in an equivalence class C is a representative of C.

• If $a \nsim a'$, then $C_a \cap C_{a'} = \emptyset$. (e.g.: in the example above, $1 \nsim 2$, $C_1 \cap C_2 = \emptyset$, $C_0 \cap C_1 = \emptyset \ldots$).

Therefore, S can be partitioned into disjoint equivalence classes $S = C_1 \cup C_2 \cup \ldots$ where $\{C_\alpha\}_{\alpha \in I}$ are equivalence classes and $C_\alpha \cap C_\beta = \emptyset$ if $\alpha \neq \beta$.

e.g.: $\mathbb{Z} = C_0 \cup C_1 \cup C_2 = (0 + 3\mathbb{Z}) \cup (1 + 3\mathbb{Z}) \cup (2 + 3\mathbb{Z})$

$GL(n, \mathbb{R}) = \coprod_{\alpha \in \mathbb{R}^*} \{ \begin{pmatrix} \alpha & 0 \\ 0 & I \end{pmatrix} SL(n, \mathbb{R}) \} = \coprod_{\alpha \in \mathbb{R}^*} \{A \mid \det(A) = \alpha\}$.

Definition 2.53 (Equivalence Relation on Groups). Let G be a group and $H \leq G$. Define an equivalence relation on G by $a \sim b$ iff $a^{-1} b \in H$.

(Check):

  (i) $a \sim a$, since $a^{-1} a = e \in H$.

  (ii) $a \sim b \Rightarrow a^{-1} b \in H \Rightarrow (a^{-1} b)^{-1} \in H \Rightarrow b^{-1} a \in H \Rightarrow b \sim a$.

(iii) $a \sim b,\ b \sim c \Rightarrow a^{-1}b \in H,\ b^{-1}c \in H \Rightarrow (a^{-1}b)(b^{-1}c) \in H$
$\Rightarrow a^{-1}c \in H \Rightarrow a \sim c.$

Definition 2.54 (Left Coset). Let G be a group and $H \leq G$. Define the **left coset** of H with representative a as $aH := C_a = \{b \in G \mid a \sim b\} = \{b \in G \mid a^{-1}b \in H,\ \text{for } b \in G\}$
$= \{b \in G \mid b = ah,\ h \in H\}.$

Example 2.55.   • $G = \mathbb{Z},\ H = 3\mathbb{Z} = \langle 3 \rangle$. The cosets of $3\mathbb{Z}$ in $\mathbb{Z}$ are (under addition):
The left cosets are: $0 + 3\mathbb{Z} = 3\mathbb{Z},\ 1 + 3\mathbb{Z},\ 2 + 3\mathbb{Z}.$

- $G = GL(n, \mathbb{R}),\ H = SL(n, \mathbb{R})$. $a \sim b$ iff $a^{-1}b \in SL(n, \mathbb{R}) \Leftrightarrow \det(a^{-1}b) = 1$
$\Leftrightarrow \det(a) = \det(b).$
The left cosets are: $C_{\begin{pmatrix} \alpha & 0 \\ 0 & I \end{pmatrix}} = \begin{pmatrix} \alpha & 0 \\ 0 & I \end{pmatrix} SL(n, \mathbb{R}) = \{B \mid \det(B) = \alpha\} = \{B \mid \det(B) = \det(\alpha)\}.$

- $G = S_3,\ H = \langle (1\,2) \rangle = \{e, (1\,2)\}.$
Left cosets: $C_e = eH = \{e, (1\,2)\}.$
$C_{(1\,2\,3)} = (1\,2\,3)H = \{(1\,2\,3), (1\,2\,3)(1\,2)\} = \{(1\,2\,3), (1\,3)\}.$
$C_{(1\,3\,2)} = (1\,3\,2)H = \{(1\,3\,2), (1\,3\,2)(1\,2)\} = \{(1\,3\,2), (2\,3)\}.$

- $G = D_4,\ H = \langle s \rangle = \{e, s\}.$
$eH = \{e, s\} = sH.$
$rH = \{r, rs\} = rsH.$
$r^2 H = \{r^2, r^2 s\} = r^2 sH.$
$r^3 H = \{r^3, r^3 s\} = r^3 sH.$

Remark 2.56. We can also define $a \sim_R b$ by $ba^{-1} \in H$. Then the equivalence class of a is called **right coset** $Ha = \{ha \mid h \in H\}.$

Theorem 2.57 (Lagrange's Theorem). *Let G be a finite group and $H \leq G$. Then $|H| \mid |G|$.*
(e.g.: all subgroups of $G = S_3$ have order 1, 2, 3, 6 only).
More precisely, let $m = [G : H] :=$ the number of disjoint left cosets of H.
Then $|H| = |G|/[G : H]$.
e.g.: • $G = S_3,\ H = \langle (1\,2) \rangle$. *Then $m = 3 \Rightarrow |H| = |S_3|/3 = 6/3 = 2$.*
    • $G = D_n,\ H = \langle r \rangle$. *Then $m = 2 \Rightarrow |H| = |D_n|/2 = 2n/2 = n$.*

Proof. Since (left cosets of H) = (equivalence classes of G), we can partition $G = eH \amalg a_2 H \amalg \cdots \amalg a_m H$ (let $a_1 := e$) into disjoint union of equivalence classes.
$(m = [G : H] < \infty$, since $|G| < \infty)$.
So we just need to check each $a_i H$ have the same number of elements $|H| = r$ (*)

$a_i H = \{a_i h_1, a_i h_2, \ldots, a_i h_r\}$ where $\{h_1, \ldots, h_r\} = H$. Then for each $a_i h_x, a_i h_y \in a_i H$:
$a_i h_x = a_i h_y \Leftrightarrow a_i^{-1} a_i h_x = a_i^{-1} a_i h_y \Leftrightarrow h_x = h_y \Leftrightarrow x = y$.
So the elements in $a_i H$ are distinct, and hence, $|a_i H| = |H| = r$.
$\therefore$ By (*), $|G| = |H| + |H| + \cdots + |H| = m|H|$. $\square$ $\square$

Corollary 2.58. *Let $|G| < \infty$. Then for all $g \in G$, $\operatorname{ord}(g) \mid |G|$.*
e.g.: all elements in $G = D_5$ must have order 1, 2, 5, 10 only.

Proof. Take $H := \langle g \rangle \leq G$. Then $\operatorname{ord}(g) = |\langle g \rangle| = |\{e, g, \ldots, g^{n-1}\}|$.
Then by Lagrange's Thm, $\operatorname{ord}(g) = |\langle g \rangle| \mid |G|$. $\square$ $\square$

Example 2.59 (Fermat's Little Theorem)**.** Let $a \in G = \mathbb{Z}_p^*$. Then $a^{p-1} = e = 1$ in $\mathbb{Z}_p^*$.
(e.g.: In $\mathbb{Z}_7$, $2^6 \equiv 3^6 \equiv \cdots \equiv 6^6 \equiv 1 \pmod{7}$)

(Proof): By corollary 2.55, for all $a \in \mathbb{Z}_p^* = G$, $d = \operatorname{ord}(a) \mid p - 1 = |\mathbb{Z}_p^*|$.
Hence, $a^d = 1$ and $d \cdot \alpha = p - 1$ for some integer α. Then
$a^{p-1} = a^{d\alpha} = (a^d)^\alpha = 1^\alpha = 1$.

More generally, $G = \mathbb{Z}_n^*$, $|G| = \phi(n) = \#\{r \mid 1 \leq r \leq n, \ \gcd(r, n) = 1\}$
(ϕ Euler's totient function). Then $a^{\phi(n)} \equiv 1 \pmod{n}$.

2.5 Normal Subgroups

Motivations: We have $G = (g_1 H) \amalg (g_2 H) \amalg \ldots$ (left cosets). Also,
$G = (Hb_1) \amalg (Hb_2) \amalg \ldots$ (right cosets).

Question: Do we have $\{gh \mid h \in H\} = gH = Hg = \{hg \mid h \in H\}$ in general?

Answer: No. Let's try $G = S_3$, $H = \langle (1\,2) \rangle$. Then
$(1\,3)H = \{(1\,3), (1\,3)(1\,2)\} = \{(1\,3), (1\,2\,3)\}$
$H(1\,3) = \{(1\,3), (1\,2)(1\,3)\} = \{(1\,3), (1\,3\,2)\} \neq (1\,3)H$.

Question: What kind of $H \leq G$ gives $gH = Hg$?

Definition 2.60 (Normal Subgroup)**.** $H \leq G$ is a **normal subgroup** if $gH = Hg \quad \forall g \in G$. (And we'll write $H \triangleleft G$ in such cases).

Example 2.61.   • $G = S_n$, $H = A_n$. Then $H \triangleleft G$.

   • If G is abelian, then any subgroup $H \leq G$ is normal.
     Proof: $\forall g \in G, \forall h \in H$, $gh = hg$ (since G is abelian). So $gH = Hg$ for all g.

- Similarly, $SL(n, \mathbb{R}) \lhd GL(n, \mathbb{R})$ is normal.
  Proof: Let $A \in GL(n, \mathbb{R})$ and $H = SL(n, \mathbb{R})$. For any $h \in H$,

$$\det(AhA^{-1}) = \det(A)\det(h)\det(A)^{-1} = 1 \cdot \det(A) \cdot \frac{1}{\det(A)} = 1$$

  $\Rightarrow AhA^{-1} \in SL(n, \mathbb{R})$, so $AH = HA$.

- If $H \leq G$ and $[G : H] = 2$, then $H \lhd G$.
  Proof: In this case, $G = H \amalg (gH)$ and $G = H \amalg (Hg)$.
  Since $H \cap gH = \emptyset$ and $H \cap Hg = \emptyset$, it must be $gH = Hg$.

- $\{e\} \lhd G$ and $G \lhd G$ are always normal subgroups.

Theorem 2.62. *Let $H \leq G$ be a subgroup. The followings are equivalent:*
(i) $H \lhd G$;   (ii) $\forall h \in H, g \in G,\ ghg^{-1} \in H$;   (iii) $gHg^{-1} = H \quad \forall g \in G$.

(Proof):
((i) $\Rightarrow$ (ii)): By assumption, $gH = Hg \quad \forall g \in G$. $\Rightarrow \forall h \in H,\ gh \in gH = Hg$
$\Rightarrow gh = h'g$ for some $h' \in H$. $\Rightarrow ghg^{-1} = h' \in H$.

((ii) $\Rightarrow$ (iii)): By (ii), $gHg^{-1} = \{ghg^{-1} \mid h \in H\} \subseteq H$. $\bullet\ g^{-1}gHg^{-1}g \subseteq g^{-1}Hg \Rightarrow H \subseteq g^{-1}Hg$.
for all $g \in G$. Then $H \subseteq (g^{-1})H(g^{-1})^{-1}$ (take "$g = g^{-1}$"). $\Rightarrow H \subseteq gHg^{-1}$.
Therefore, $H = gHg^{-1}$.

((iii) $\Rightarrow$ (i)): $gHg^{-1}g = Hg \quad \forall g \in G \Rightarrow gH = Hg \quad \forall g \in G \Rightarrow H \lhd G$. $\square$

Corollary 2.63. *Let $\phi : G \to H$ be a homomorphism. Then $\ker \phi \lhd G$.*

(Proof): By (ii) in theorem 2.59, we only need to show $\forall k \in \ker \phi,\ g \in G,\ gkg^{-1} \in \ker \phi$.
Indeed, $\phi(gkg^{-1}) = \phi(g)\phi(k)\phi(g^{-1}) = \phi(g)e_H(\phi(g))^{-1} = \phi(g)(\phi(g))^{-1} = e_H$. $\square$

Example 2.64.   - $\phi : GL(n, \mathbb{R}) \to \mathbb{R}^*$ with $\phi(A) = \det(A)$. Then $\ker \phi = \{A \mid \det(A) = 1\} = SL(n, \mathbb{R})$.
     $\therefore SL(n, \mathbb{R}) \lhd GL(n, \mathbb{R})$.

- $\psi : S_n \to (\{\pm 1\}, \times)$ with $\psi(\sigma) :=$ parity of σ. Then $\ker \psi = A_n \lhd S_n$.

2.6   Quotient Groups

Motivation: Let $H \lhd G$. We want to define a group structure on the set of left cosets $G/H :=$ $\{gH \mid g \in G\}$.

Definition 2.65 (Quotient Group). Let G be a group and $H \triangleleft G$. Define the **quotient group** (or factor group) G/H as the set of left cosets $\{gH \mid g \in G\}$ with the multiplication:

$$(aH) * (bH) := (ab)H$$

(WARNING):
There are lots of repetitions in this set, i.e.: there can be $a_1 \neq a_2$, s.t.: $a_1 H = a_2 H$

(Check):

- $*$ is well-defined: Suppose $aH = a'H$ & $bH = b'H$ — (*).
  By (*), $h_a := a^{-1}a' \in H$, $h_b := b^{-1}b' \in H$. Then $a'b' = (ah_a)(bh_b) = abh_a^h h_b$
  $\Rightarrow a'b' \in (ab)H \Rightarrow (a'b')H = (ab)H$

- $(G/H, *)$ is a group:

  1. $((aH) * (bH)) * (cH) = (ab)H * (cH) = (abc)H = (aH) * (bc)H$
     $$= (aH) * ((bH) * (cH))$$

  2. $(aH) * (eH) = aH = (eH) * (aH)$, i.e.: $e_{G/H} = eH$.

  3. $(aH) * (a^{-1}H) = eH = e_{G/H} = (a^{-1}H) * (aH)$, i.e.: $(aH)^{-1} = a^{-1}H$.

Example 2.66.   • $G = \mathbb{Z}$, $H = \langle n \rangle = n\mathbb{Z} \triangleleft G$. (Exercise : All subgroups of abelian group G are normal)
  $G/H = \{0 + n\mathbb{Z} , 1 + n\mathbb{Z} , 2 + n\mathbb{Z} , \ldots , (n-1) + n\mathbb{Z}\}$
  For $a + n\mathbb{Z}$, $b + n\mathbb{Z} \in G/H$, $(a + n\mathbb{Z}) * (b + n\mathbb{Z}) := (a + b) + n\mathbb{Z} = c + n\mathbb{Z}$
  where $c \equiv a + b \pmod{n}$ & $0 \leq c < n$
  $\therefore$ There's an isomorphism $\phi : G/H = \mathbb{Z}/n\mathbb{Z} \overset{\cong}{\to} \mathbb{Z}_n$ with $\phi(a + n\mathbb{Z}) := a \pmod{n}$
  Remark: From now on, I may interchange $\mathbb{Z}/n\mathbb{Z}$ and $\mathbb{Z}_n$.

- $A_n \triangleleft S_n$, then $S_n/A_n = \{A_n , \tau A_n\}$ (τ is any transposition $\tau = (a\ b)$).
  (Recall $|S_n| = n!$, $|A_n| = \frac{n!}{2}$, $\therefore |S_n/A_n| = 2$)
  $\therefore S_n/A_n \cong \mathbb{Z}_2 \ (\cong \mathbb{Z}/2\mathbb{Z})$

- $SL(n, \mathbb{R}) \trianglelefteq GL(n, \mathbb{R})$.

$$GL(n, \mathbb{R})/SL(n, \mathbb{R}) = \left\{ \begin{pmatrix} x & \ldots & 0 \\ \vdots & 1 & \ldots \\ 0 & \ldots & 1 \end{pmatrix} SL(n, \mathbb{R}) \mid x \in \mathbb{R}^* \right\}$$
$$= \{A \in GL(n, \mathbb{R}) \mid \det(A) = x\}$$

$$\left(\begin{pmatrix} x & \dots & 0 \\ \vdots & 1 & \dots \\ 0 & \dots & 1 \end{pmatrix} SL(n,\mathbb{R}) \right) \left(\begin{pmatrix} y & \dots & 0 \\ \vdots & 1 & \dots \\ 0 & \dots & 1 \end{pmatrix} SL(n,\mathbb{R}) \right)$$

$$= \left(\begin{pmatrix} x & \dots & 0 \\ \vdots & 1 & \dots \\ 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} y & \dots & 0 \\ \vdots & 1 & \dots \\ 0 & \dots & 1 \end{pmatrix} \right) SL(n,\mathbb{R})$$

$$= \begin{pmatrix} xy & \dots & 0 \\ \vdots & 1 & \dots \\ 0 & \dots & 1 \end{pmatrix} SL(n,\mathbb{R})$$

$\therefore GL(nd,\mathbb{R})/SL(n,\mathbb{R}) \cong (\mathbb{R}^*, \times)$

- $K = \{e, (12)(34), (13)(24), (14)(23)\} \triangleleft S_4$
  Then $|S_4/K| = |S_4|/|K| = 24/4 = 6$ Indeed, $S_4/K \cong S_3$

2.7   First Isomorphism Theorem

Theorem 2.67 (First Isomorphism Theorem). *Let $\phi : G \to H$ be a group homomorphism. Then $G/\ker\phi \cong im\ \phi$.*

Example 2.68.   • $\phi : \mathbb{Z} \to \mathbb{Z}_n$ with $\phi(a) := a \pmod{n}$ is a homomorphism, with
    $\ker\phi = n\mathbb{Z}$, $im\phi = \mathbb{Z}_n$. Then by 1st isomorphism, $\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}_n$.

- $\det : GL(n,\mathbb{R}) \to (\mathbb{R}^*, \times)$ with $\ker(\det) = \{A \mid \det(A) = 1\} = SL(n,\mathbb{R})$,
  $im(\det) = \mathbb{R}^*$. Hence, $GL(n,\mathbb{R})/SL(n,\mathbb{R}) \cong \mathbb{R}^*$.

- $\phi : \mathbb{R} \to GL(2,\mathbb{R})$ with $\phi(x) := \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$. Then

  $\ker\phi = 2\pi\mathbb{Z}$, and $im\phi = \left\{ \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix} \middle| 0 \le x < 2\pi \right\} = SO(2)$.
  $\therefore \mathbb{R}/2\pi\mathbb{Z} \cong SO(2)$.

Proof of 1st isomorphism theorem: Define $\Psi : G/\ker\phi \to im\phi$ by
$\Psi(g\ker\phi) := \phi(g)$.

(1) Ψ is well-defined: For $g\ker\phi = g'\ker\phi$, $g^{-1}g' \in \ker\phi \Leftrightarrow e_H = \phi(g^{-1}g')$
    $= \phi(g)^{-1}\phi(g') \Leftrightarrow \phi(g)e_H = \phi(g)\phi(g)^{-1}\phi(g') = \phi(g') \Leftrightarrow \phi(g) = \phi(g')$.
    $\Leftrightarrow \Psi(g\ker\phi) = \Psi(g'\ker\phi)$.

(2) Ψ is a homomorphism: $\Psi((g\ker\phi)(g'\ker\phi)) = \Psi(gg'\ker\phi) := \phi(gg') = \phi(g)\phi(g')$
    $= \Psi(g\ker\phi)\Psi(g'\ker\phi)$.

(3) Ψ is surjective: Obvious from definition of Ψ.

(4) Ψ is injective: $\forall\, g \ker \phi \in \ker \Psi$, $\phi(g) = e_H \Rightarrow g \in \ker \phi$
$\Rightarrow g \ker \phi = e_G \ker \phi = e_{G/\ker \phi} \Rightarrow \ker \Psi = \{e_{G/\ker \phi}\}$. $\qquad\square$

Applications of quotient groups

- We can construct new groups from old ones.

- Cauchy's Theorem: Suppose $|G| < \infty$ is finite and $p \mid |G|$ for some prime p. Then G must have an element g of order p.

- Classification of finite groups.

2.8 Simple Groups

Definition 2.69 (Simple group)**.** A group G is called **simple** if G has no normal subgroups other than $\{e\}$ and G itself.

Reason for this definition:
If G is **NOT** simple, then we have $N \triangleleft G$. So we can "decompose" G into G/N and N, and study them individually.

Remark/WARNING: If $|G| < \infty$ is NOT simple, then $|G/N| = |G|/|N|$.
So $|G/N \times N| = |G/N| \times |N| = |G|/|N| \times |N| = |G|$. But $G \not\cong G/N \times N$ in general.

Example 2.70. • S_n is **NOT** simple, since $A_n \triangleleft S_n$. So we can "understand" S_n by understanding A_n and $S_n/A_n \cong \mathbb{Z}_2$ individually.
But $S_n \not\cong A_n \times (S_n/A_n)$.

- A_4 is **NOT** simple, since $K = \{e, (12)(34), (13)(24), (14)(23)\} \triangleleft A_4$.
However, A_n is simple for $n \geq 5$ (very important in Galois theory: there're no radical solutions for quintic polynomial $x^5 + a_4 x^4 + a_3 x^3 + \cdots + a_0 = 0$).

2.9 Fundamental Theorem of Finite Abelian Groups

Observations:

(1) $\mathbb{Z}_n$ $(\cong \mathbb{Z}/n\mathbb{Z} = \mathbb{Z}/\langle n \rangle)$ is abelian

(2) G and H are abelian $\iff$ $G \times H$ is abelian

(3) If $\gcd(m, n) = 1$, then $\mathbb{Z}_{mn} \cong \mathbb{Z}_m \times \mathbb{Z}_n$

Theorem 2.71. *All finite abelian groups are of the form $G \cong \mathbb{Z}_{p_1^{a_1}} \times \cdots \times \mathbb{Z}_{p_k^{a_k}}$ where $p_1 \leq \cdots \leq p_k$ are primes, $a_1, \ldots, a_k \in \mathbb{N} \setminus \{0\}$ (p_i NOT necessarily distinct).*

(Proof): There are 2 main steps to prove theorem 2.68:

Step 1: If $|G| = q_1^{a_1} \ldots q_r^{a_r}$, q_i distinct primes, then $G \cong G_1 \times \cdots \times G_r, |G_i| = q_i^{a_i}$

Step 2: If $|G| = q^b$, then $G \cong \mathbb{Z}_{q^{b_1}} \times \cdots \times \mathbb{Z}_{q^{b_r}}, \sum_{j=1}^r b_j = b$

Proof of step 1: Suppose $|G| = p^a q^b$ (p, q distinct primes). We want to show $G \cong G_1 \times G_2$, $|G_1| = p^a =: m$, $|G_2| = q^b =: n$, and the general case follows from induction on # of prime powers.

Let $G^m := \{g^m \mid g \in G\} \leq G$ & $G^n := \{g^n \mid g \in G\} \leq G$.

<u>**Claim 1:**</u> $G^m \cap G^n = \{e\}$. Indeed, suppose $x \in G^m \cap G^n$. Then $x = g^m = h^n$ for some $g, h \in G$. Then $x^n = g^{mn} = e$ & $x^m = h^{nm} = e$ (by Lagrange's Theorem, $|G| = mn$). Note that $\gcd(m, n) = \gcd(p^a, q^b) = 1$

$\Rightarrow \alpha m + \beta n = \gcd(m, n) = 1$ for some $\alpha, \beta \in \mathbb{Z}$. $\Rightarrow x^{\alpha m + \beta n} = (x^m)^\alpha (x^n)^\beta = x^1$

$\Rightarrow e \cdot e = x^1 \Rightarrow x = e$. Then <u>claim 1 holds</u>.

<u>**Claim 2:**</u> $\phi : G^m \times G^n \to G$ with $\phi(g^m, h^n) := g^m (h^n)^{-1}$ is an isomorphism:

- ϕ is a homomorphism. (exercise, need G is abelian)

- ϕ is injective: Suppose $\phi(g^m, h^n) = e$. Then $g^m (h^n)^{-1} = e \Rightarrow g^m = h^n$
  $\Rightarrow g^m, h^n \in G^m \cap G^n = \{e\} \Rightarrow g^m = h^n = e \Rightarrow \ker \phi = \{(e, e)\}$.

- ϕ is surjective: Take $y \in G$.
  $y = y^1 = y^{\alpha m + \beta n} = (y^\alpha)^m (y^\beta)^n = \phi((y^\alpha)^m, (y^{-\beta})^n)$
  Hence, ϕ is an isomorphism.

<u>**Claim 3:**</u> $|G^m| = n = q^b$, $|G^n| = m = p^a$

Begin by showing $\gcd(|G^n|, q) = 1$. Suppose $q \mid |G^n|$. Then by Cauchy's Theorem, there exists $x = g^n$ in G^n of order q.

By Bezout, $\gcd(q, m) = 1$. So one has $\alpha q + \beta m = 1$, and $x^q = e$. Then $x^{\alpha q} = e \Rightarrow x^{1 - \beta m} = e \Rightarrow x \cdot (x^m)^{-\beta} = e \Rightarrow x(g^{nm})^{-\beta} = e$

$\xrightarrow{\text{Lagrange's}} x \cdot e^{-1} = e \Rightarrow x = e$, contradicting order of x is q.

So $\gcd(|G^n|, q) = 1 \Rightarrow |G^n| = p^r$ & $\gcd(|G^m|, p) = 1 \Rightarrow |G^m| = q^s$

But $G^m \times G^n \cong G$, so $p^r \cdot q^s = |G| = p^a q^b$. Then $a = r, b = s$.

Therefore, <u>claim 3 holds</u>.

Proof of step 2: Suppose $|G| = p^a$. Then for all $x \in G$, $\text{ord}(x) = p^i$ for some $0 \leq i \leq a$ by Lagrange's Theorem.

Let $m \in G$ be an element with maximum order $\text{ord}(m) = p^l$ in G.

Proposition: Let G be an abelian group with $|G| = p^a$. Suppose $\mu \in G$ has maximum order $\text{ord}(\mu) = p^l$ in G, then $G \cong \langle \mu \rangle \times K \cong \mathbb{Z}_{p^l} \times K$.

Proof of Proposition: Induction on $p^a = |G|$. If $a = 1$, then $|G| = p \Rightarrow G \cong \mathbb{Z}_p$.

Then suppose the Proposition holds for all abelian G with $|G| = p^r, r < a$.

Now let $|G| = p^a$. Suppose $\text{ord}(\mu) = p^m$ is maximal:

- If $m = a$, then $\langle \mu \rangle = G$ and then we're done.

- If $m < a$, then let $\nu \in G$ be such that ν is non-identity element NOT in $\langle \mu \rangle$ with smallest possible order p^k.

<u>Claim 4:</u> $\text{ord}(\nu) = p$. $\because \text{ord}(\nu^p) < p^k$, so $\nu^p \in \langle \mu \rangle$ and $\nu^p = \mu^i$ for some $i \in \mathbb{Z}$. Then $(\mu^i)^{p^{m-1}} = (\nu^p)^{p^{m-1}} = \nu^{p^m} = e$, since every $g \in G$ has order $\leq p^m$.

$\Rightarrow \text{ord}(\mu^i) \mid p^{m-1} \Rightarrow |\langle \mu^i \rangle| < p^m \Rightarrow \langle \mu^i \rangle \neq \langle \mu \rangle \Rightarrow \gcd(i, p^m) > 1$

$\therefore i = pj$ for some $j \in \mathbb{Z}$. Hence, $\nu^p = \mu^{pj}$ for some $j \in \mathbb{Z}$.

Let $c := \nu \mu^{-j}$. Then $c \notin \langle \mu \rangle$ and $c^p = e$, since $\nu \notin \langle \mu \rangle$ is chosen to have smallest order. Therefore, $\text{ord}(\nu) = p$. $\Rightarrow$ <u>claim 4 holds</u>.

<u>Claim 5:</u> $\langle \mu \rangle \cap \langle \nu \rangle = \{e\}$. Let $\nu^l \in \langle \mu \rangle \cap \langle \nu \rangle$. Then $0 \leq l < p$, since $\text{ord}(\nu) = p$. Suppose $l \neq 0$, then $\gcd(l, p) = 1 \Rightarrow \alpha l + \beta p = 1$

$\Rightarrow \langle \mu \rangle \cap \langle \nu \rangle \ni (\nu^l)^\alpha = \nu \cdot \nu^{-\beta p} = \nu \Rightarrow \nu \in \langle \mu \rangle$. Contradiction.

Therefore, $\langle \mu \rangle \cap \langle \nu \rangle = \{e\}$, i.e.: <u>claim 5 holds</u>.

Let $\bar{G} := G/\langle \nu \rangle$, s.t.: $|\bar{G}| = \frac{|G|}{|\langle \nu \rangle|} = \frac{p^a}{p} = p^{a-1}$ & write $\bar{g} := g\langle \nu \rangle \in \bar{G}$.

<u>Claim 6:</u> $\bar{\mu} \in \bar{G}$ has order p^m (maximal order). Obviously, $\bar{\mu}^{p^m} = (\mu\langle \nu \rangle)^{p^m} = \mu^{p^m}\langle \nu \rangle = e\langle \nu \rangle = e_{\bar{G}} \Rightarrow \text{ord}(\bar{\mu}) \mid p^m$

Suppose by contrary, $\text{ord}(\bar{\mu}) = p^u$ for $u < m$. Then $\mu^{p^u}\langle \nu \rangle = e_{\bar{G}} = e\langle \nu \rangle$

$\Rightarrow \mu^{p^u} \in \langle \nu \rangle \Rightarrow \mu^{p^u} \in \langle \mu \rangle \cap \langle \nu \rangle \xrightarrow{\text{claim 5}} \{e\} \Rightarrow \mu^{p^u} = e$, contradicting $\text{ord}(\mu) = p^m$. Therefore, <u>claim 6 holds</u>.

By induction, there is another subgroup $\bar{K} \leq \bar{G}$ s.t.: $\langle \bar{\mu} \rangle \times \bar{K} \cong \bar{G}$ $(*)$.

That is, every element in $\bar{G}$ can be written <u>uniquely</u> as $\bar{\mu}^i \bar{k} \in \bar{G}$ for some $\bar{\mu}^i \in \langle \bar{\mu} \rangle, \bar{k} \in \bar{K}$. Let $\pi : G \to \bar{G}$ be given by $\pi(g) := \bar{g}$. Then $K := \pi^{-1}(\bar{K}) = \{k \in G \mid \pi(k) \in \bar{K}\} \leq G$.

<u>Claim 7:</u> $\pi|_K : K \to \bar{K}$ has $\ker(\pi|_K) = \langle \nu \rangle$. $x \in \ker(\pi|_K) \iff \pi(x) = x\langle \nu \rangle = e_{\bar{G}} = \langle \nu \rangle \iff x \in \langle \nu \rangle$. Therefore, <u>claim 7 holds</u>.

Hence, by 1st isomorphism theorem, $|K/\langle \nu \rangle| = |\bar{K}|$. $\Rightarrow |K|/p \overset{(*)}{=} \frac{|G|}{|\langle \mu \rangle|} \Rightarrow |K| = \frac{p \cdot p^{a-1}}{p^m} \Rightarrow |K| = p^{a-m}$

<u>Claim 8:</u> $\langle \mu \rangle \cap K = \{e\}$ in G. Let $\mu^i \in \langle \mu \rangle \cap K$, then $\pi(\mu^i) = \bar{\mu}^i \cdot e_{\bar{K}} = e_{\langle \bar{\mu} \rangle} \cdot \bar{k}$. By uniqueness, $\bar{\mu}^i = e_{\langle \bar{\mu} \rangle}$ & $\bar{k} = e_{\bar{K}}$

$\Rightarrow \mu^i \in \langle \nu \rangle \Rightarrow \mu^i \in \langle \mu \rangle \cap \langle \nu \rangle = e$. Then <u>claim 8 holds</u>.

Finally, consider the homomorphism $\theta : \langle \mu \rangle \times K \to G$ with $\theta(\mu^i, k) := \mu^i k$

Then by claim 8, θ is injective. By claim 7, $|\langle \mu \rangle \times K| = |\langle \mu \rangle||K| = p^m \cdot p^{a-m} = p^a = |G|$. Hence, θ is surjective as well, and hence:

$\langle \mu \rangle \times K \cong G$ $\hfill \square$

Example 2.72. $|G| = 4$. Then all elements of G have order $1, 2, 4$.

   (i) If G has an element m of order 4, then $G = \langle m \rangle \cong \mathbb{Z}_4$.

  (ii) Suppose all elements of G has order 1 or 2. Take any $m \in G$ of order 2, $G \cong \langle m \rangle \times K \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

Corollary 2.73 (of proposition in the proof of Theorem 2.69). *If $|G| = p^a$, abelian, then $G \cong$*
$\mathbb{Z}_{p^{b_1}} \times \mathbb{Z}_{p^{b_2}} \times \cdots \times \mathbb{Z}_{p^{b_\ell}}$, *where $\sum_{j=1}^{\ell} b_j = a$.*
(Proof): Induction on $|G| = p^a$.

Example 2.74. How does an abelian G with $|G| = 360 = 2^3 \times 3^2 \times 5$ look like?
By Step 1: $G \cong G_8 \times G_9 \times G_5$ where $|G_i| = i$ for $i = 5, 8, 9$.
By Step 2: G_8 can be $\mathbb{Z}_8$ or $\mathbb{Z}_2 \times \mathbb{Z}_4 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$ or $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$.
                G_9 can be $\mathbb{Z}_9$ or $\mathbb{Z}_3 \times \mathbb{Z}_3$.
                G_5 can be $\mathbb{Z}_5$ only.

$$\therefore G \cong \begin{pmatrix} \mathbb{Z}_8 \\ \mathbb{Z}_4 \times \mathbb{Z}_2 \\ \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \end{pmatrix} \times \begin{pmatrix} \mathbb{Z}_9 \\ \mathbb{Z}_3 \times \mathbb{Z}_3 \end{pmatrix} \times \mathbb{Z}_5 \quad \text{(with 6 possibilities)}$$

Corollary 2.75 (Smith Normal Form). *All finite abelian group G is isomorphic to $G \cong \mathbb{Z}_{d_1} \times \cdots \times$*
$\mathbb{Z}_{d_k}$ *where $d_i > 1$ integers, and $d_i \mid d_{i+1}$ $\forall i = 1, \ldots, k-1$.*

Moreover, if $H \cong \mathbb{Z}_{e_1} \times \cdots \times \mathbb{Z}_{e_\ell}$ for $e_i > 1$ integers, $e_i \mid e_{i+1}$ $\forall i = 1, \ldots, \ell - 1$. Then:
$(G \cong H) \iff (\ell = k \text{ and } d_i = e_i \ \forall i)$

Chapter 3

Rings

3.1 Basic Definition

Rings

Definition 3.1 (Ring). A ring $(R, +, \cdot)$ is a set equipped with 2 binary operations $+, \cdot : R \times R \to R$ such that

1. $(R, +)$ is an abelian group with additive identity $0_R \in R$.

2. $(R, \cdot)$ is associative: $(a \cdot b) \cdot c = a \cdot (b \cdot c) \quad \forall a, b, c \in R$.

3. $(R, +, \cdot)$ is distributive:

$$\begin{cases} a \cdot (b + c) = a \cdot b + a \cdot c \\ (a + b) \cdot c = a \cdot c + b \cdot c \end{cases} \quad \forall a, b, c \in R.$$

Example 3.2.
- $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ are rings.

- $\mathbb{Z}[i] = \{a + bi \mid a, b \in \mathbb{Z}\}$ where $i = \sqrt{-1}$ (Gaussian integers)
  $(a + bi) \cdot (c + di) := (ac - bd) + (ad + bc)i$
  $(a + bi) + (c + di) := (a + c) + (b + d)i$

- More generally, $\mathbb{Z}[e^{\frac{2\pi i}{n}}] = \{a_0 + a_1 e^{\frac{2\pi i}{n}} + \cdots + a_k e^{\frac{2\pi k i}{n}} + \cdots + a_{n-1} e^{\frac{2\pi (n-1) i}{n}} \mid a^i \in \mathbb{Z}\}$
  Since $e^{\frac{2\pi i}{n}} := cos\frac{2\pi i}{n} + i(sin\frac{2\pi i}{n})$, then $n = 4$, $e^{\frac{2\pi i}{4}} = i$.

- $2\mathbb{Z} = \{2a \mid a \in \mathbb{Z}\}$ is a ring,
  but $(\{\cdots, -3, -1, 1, 3, \cdots\}, +, \cdot)$ is **NOT** a ring.

- $n\mathbb{Z} = \{na \mid a \in \mathbb{Z}\}$ is a ring

- $M_{n \times n}(\mathbb{Z})$ is a ring

- $\mathbb{Z}[x] = \{$all polynomials with integer coefficients$\}$
  For $f(x) = a_m x^m + \cdots + a_1 x + a_0$ and $g(x) = b_n x^n + \cdots + b_1 x + b_0 \in \mathbb{Z}[x]$,

$$(f \cdot g)(x) := a_m b_n x^{n+m} + \cdots + \sum_{\substack{p \geq 0 \\ q \geq 0 \\ p+q=i}} a_p b_q x^i + \cdots + a_0 b_0.$$

Mativation

Solve $x^2 + y^2 = z^2$ for integers x, y, z. $\Rightarrow (x + iy)(x - iy) = z^2$ in $\mathbb{Z}[i]$,

so we wish to study more about properties of the Gaussian integers $\mathbb{Z}[i]$, e.g.: does the **fundamental theorem of arithmetic** hold for $\mathbb{Z}[i]$, i.e.: can every element in $\mathbb{Z}[i]$ be factorized into product of prime numbers uniquely?

Fermat's Last Theorem: $x^n + y^n = z^n$ for $n \geq 3$, $x^n + y^n = (x - e^{\frac{2\pi i}{n}} y) \cdots (x - e^{\frac{2\pi (n-1)i}{n}} y)$ in $\mathbb{Z}[e^{\frac{2\pi i}{n}}]$

Definition 3.3 (Unital Rings and Commutative Rings)**.** Let $(R, +, \cdot)$ be a ring. We say:

1. R is **unital** if $\exists\, 1_R \in R$ such that

   $1_R \cdot r = r \cdot 1_R = r \; \forall\, r \in R$

   (1_R is the multiplication identity of R)

   (non-example: $R = 2\mathbb{Z}$ or $n\mathbb{Z}$)

2. If R is unital, then the **units** of R are the elements

   $U(R) = \{a \in R \mid \exists a^{-1} \in R \text{ s.t. } aa^{-1} = a^{-1}a = 1_R\}$

   (e.g.: $U(\mathbb{Z}) = \{\pm 1\}$, $U(M_{n \times n}(R)) = GL(n, \mathbb{R})$), and $U(\mathbb{Z}[i]) = \{\pm 1, \pm i\}$, why they are the only units?)

3. R is **commutative** if $a \cdot b = b \cdot a \; \forall a, b \in R$

   (non-example: $R = M_{n \times n}(\mathbb{Z})$)

Example 3.4. $(\mathbb{Z}_n, +, \cdot)$ is a commutative and unital ring with the multiplication identity $1_R = [1]$. The units of $\mathbb{Z}_n$ are $U(\mathbb{Z}) = \mathbb{Z}_n^* = \{a \mid gcd(a, n) = 1\}$

Remark 3.5. • The additive identity $0_R \in R$ is unique in R. If R is unital, then the multiplication ifentity $1_R \in R$ is also unique

• We write $-r \in R$ to be the additive inverse of $r \in R$, i.e.: $(-r) + r = r + (-r) = 0_R$

• If $r \in U(R)$ is a unit in a (commutative) unital ring, then we write r^{-1} to be the multiplication inverse of r, i.e.: $r^{-1}r = rr^{-1} = 1_R$

• We write $nr := r + \cdots + r$ (n terms of r) for $n \in \mathbb{N}$ and $r \in R$.

• For $a, b \in R$, we write $a \mid b$ if $\exists c \in R$ s.t.: $ac = b$

Proposition 3.6.   *1.* $0_R \cdot r = r \cdot 0_R = 0_R$

   2. $(-1_R) \cdot r = r \cdot (-1_R) = -r$

   3. $(-1_R) \cdot (-r) = (-r) \cdot (-1_R) = r$

Proof:

1. For any $a \in R$, $a \cdot r = (0_R + a) \cdot r = 0_R \cdot r + a \cdot r$

   By the uniqueness of 0_R, $0_R \cdot r = 0_R$

2. $0_R = 0_R \cdot r = (1_R + (-1_R)) \cdot r = 1_R \cdot r + (-1_R) \cdot r = r + (-1_R) \cdot r$
$\Rightarrow -r = (-r) + 0_R = (-r) + (r + (-1_R) \cdot r) = (-1_R) \cdot r$
Similarly for $r \cdot (-1_R) = -r$, by taking $0_R \cdot r = ((-1_R) + 1_R) \cdot r$

3. Directly proved by proposition 3.5(2). $\square$

Definition 3.7 (Product Rings)**.** Let $(R_1, +_1, \cdot_1)$ and $(R_2, +_2, \cdot_2)$ be two rings. Then $R_1 \times R_2$ is called a **product ring** with
$(r_1, r_2) +_{R_1 \times R_2} (r_1' + r_2') := (r_1 +_1 r_1', r_2 +_2 r_2')$
$(r_1, r_2) \cdot_{R_1 \times R_2} (r_1 \cdot_1 r_1', r_2 \cdot_2 r_2')$

Definition 3.8 (Subrings)**.** Let $(R, +, \cdot)$ be a ring. A subset $S \subset R$ is a subring if
$+ \mid_{S \times S} \colon S \times S \to S$ and $\cdot \mid_{S \times S} \colon S \times S \to S$ give a ring substructure of S.

Example 3.9.   • $\mathbb{Z}[i] \subset \mathbb{C}$ is a subring

- $n\mathbb{Z} \subset \mathbb{Z}$ is a subring

- $\{\cdots, -3, -1, 1, 3, \cdots\} \subset \mathbb{Z}$ is **NOT** a subring

Proposition 3.10. *$S \subset R$ is a subring $\Leftrightarrow \forall a, b \in S \colon a + b \in S, \ -a \in S, \ and \ a \cdot b \in S$*

Definition 3.11 (Field)**.** Let R be a commutative unital ring. We say R is a **field** if all nonzero elemens of R are in the units U(R) of R.
(Field is a very special kind of ring)

Example 3.12.   • $\mathbb{Q}$ is a field, since all nonzero elements $\frac{a}{b}$ of $\mathbb{Q}$ has a multiplication inverse $\left(\frac{a}{b}\right)^{-1} = \frac{b}{a}$

- $\mathbb{R}$ and $\mathbb{C}$ are fields

- $\mathbb{Z}$ is **NOT** a field: $u(\mathbb{Z}) = \{\pm 1\} \neq \mathbb{Z} \setminus \{0\}$

- $\mathbb{Z}_p$ is a field for all p prime

More examples:

- $n\mathbb{Z}(\textbf{NOT} \text{ unital}) \subset \mathbb{Z}(\textbf{NOT} \text{ field}) \subset \mathbb{Q}(\text{field}) \subset \mathbb{R}(\text{field}) \subset \mathbb{C}(\text{field})$

- $R[x] = \{\sum_i a_i x^i \mid a_i \in R\}$ is commutative if R is commutative, and is unital if R is unital

- $M_{n \times n}(R) = \{(a_{ij}) \mid a_{ij} \in R\}$ is **NOT** commutative for $n \geq 2$

3.2   Ring Homomorphism

Definition 3.13 (Ring Homomorphism)**.** Let R and S be rings. A map $\phi \colon R \to S$ is a **homomorphism of rings** if
$\phi(a +_R b) = \phi(a) +_S \phi(b)$ and $\phi(a \cdot_R b) = \phi(a) \cdot_S \phi(b)$

- If R and S are unital, we say a homomorphism $\phi \colon R \to S$ is **unital** if $\phi(1_R) = 1_S$

- If ϕ is bijective,then ϕ is called a **ring isomorphism**

Example 3.14.   • $\phi : \mathbb{Z} \to \mathbb{Z}$, $\phi(n) := 2n$. Then:

    1. $\phi : (\mathbb{Z}, +) \to (\mathbb{Z}, +)$ is a group homomorphism, **BUT**

    2. $\phi : (\mathbb{Z}, +, \cdot) \to (\mathbb{Z}, +, \cdot)$ is **NOT** a ring homomorphism, since $\phi(a \cdot b) = 2ab \neq 4ab = (2a) \cdot (2b) = \phi(a) \cdot \phi(b)$ for $a, b \neq 0$

- Similarly, $\phi : \mathbb{Z}_{10} \to \mathbb{Z}_{10}$ with $\phi(n) = 2n$ is a group homomorphism, but **NOT** a ring homomorphism.
  But $\phi(n) := 5n$ is a ring homomorphism.

- $\phi : \mathbb{Z} \to \mathbb{Z}_m$ with $\phi(n):=$n(mod m) is a ring homomorphism

- Let R be a commutative unital ring. $\phi : R[x] \to R$ with $\phi(p(x)) := p(1_R)$
  Check $\phi(p + q) = (p + q)(1_R) = p(1_R) + q(1_R) = \phi(p) + \phi(q)$ and $\phi(p \cdot_{R[x]} q) = \phi(p) \cdot_R \phi(q)$

- $\mathbb{Z} \mapsto \mathbb{Q}, \mathbb{Q} \mapsto \mathbb{R}, \mathbb{R} \mapsto \mathbb{C}$

Proposition 3.15. *Let $\phi : R \to S$ be a ring homomorphism.*

    1. $\phi(0_R) = 0_S$

    2. $\phi(-a) = -\phi(a)$

    3. If ϕ is unital and $a \in U(R)$, then $\phi(a) \in U(S)$ with $\phi(a)^{-1} = \phi(a^{-1})$

    4. If $\phi : R \xrightarrow{\cong} S$ is an isomorphism, then $\phi^{-1} : S \to R$ is a ring isomorphism as well

  Proof:

    1. $\phi(r) = \phi(0_R + r) = \phi(0_R) + \phi(r)$. By uniqueness of additive identity $\phi(0_r) = 0_S$

    2. Same as group homomorphism

    3. $\phi(a^{-1}) \cdot \phi(a) = \phi(a^{-1}a) = \phi(1_R) = 1_S$ and similarly we have $\phi(a)\phi(a^{-1}) = 1_S$. Hence, $\phi(a)$ is a unit with $\phi(a)^{-1} = \phi(a^{-1})$

    4. Let α and $\beta \in S$. Then $\exists a, b \in R, s.t. : \phi(a) = \alpha$ and $\phi(b) = \beta$——(*)
    (WTS: $\phi^{-1}(\alpha\beta) = \phi^{-1}(\alpha)\phi^{-1}(\beta)$)
    $\phi^{-1}(\alpha) \cdot \phi^{-1}(\beta) = a \cdot b = \phi^{-1}(\phi(ab)) = \phi^{-1}(\phi(a) \cdot \phi(b)) = \phi^{-1}(\alpha \cdot \beta)$
    (HW: Check $\phi^{-1}(\alpha + \beta) = \phi^{-1}(\alpha) + \phi^{-1}(\beta)$) $\square$

Proposition 3.16. *Let $\phi : R \to S$ be a ring homomorphism. Then $ker\phi := \{r \in R \mid \phi(r) = 0_S\} \leq R$ and $im\phi := \{\phi(r) \mid r \in R\} \leq S$ are sunrings.*

  Proof: Let a,b$\in ker\phi$, i.e.: $\phi(a) = \phi(b) = 0_S$. (WTS: a+b, -a, $a \cdot b \in ker\phi$)

- $\phi(a + b) = \phi(a) + \phi(b) = 0_S + 0_S = 0_S$

- $\phi(-a) = -\phi(a) = -0_S = 0_S$

- $\phi(a \cdot b) = \phi(a) \cdot \phi(b) = 0_S \cdot 0_S = 0_S$ (the last equality is left to check in the last lecture)

Remark 3.17. In groups, we know $ker\phi$ is a **normal** subroup of R. How about rings, are there any notion of "normal subring"?

Answer: We'll study **ideals**, which is the ring analogue of normal subgroups.

3.3   Integral Domain

Definition 3.18 (Zero-divisor)**.** Let R be a ring. A nonzero element r$\in R$ is a **zero-divisor** if $\exists 0 \neq s \in R, s.t. : r \cdot s = 0$ or $s \cdot r = 0$.

Example 3.19. R=$\mathbb{Z}$. Then $2 \in R$ is a zerodivisor, since $2 \cdot 3 = 6 = 0$ in R.

Definition 3.20 (Integral Domain)**.** If R has no zerodivisors, then R is a **domain**.
Moreover, if R is commutative ring with no zerodivisors, then R is an **integral domain (ID)**.

Example 3.21.   • $\mathbb{Z}_6$ is **NOT** ID. More generally, $\mathbb{Z}_m$ is ID $\Leftrightarrow$ m is prime.

   • $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}, \mathbb{Z}[i]$ are IDs.

   • If R is ID, then so is R[x].

Proposition 3.22 (Cancellation Property)**.** *Let R be commutative ring. Then R is ID $\Leftrightarrow$ whenever ca=cb for some $c \neq 0$, then a=b.*

Proof: ca=cb $\Leftrightarrow$ ca+c(-b)=0 $\Leftrightarrow$ c(a+(-b))=0 $\Leftrightarrow$ c(a-b)=0 $\Leftarrow$(R is ID)$\Rightarrow$ c=0 OR a-b=0 $\Leftrightarrow$ a-b=0 $\Leftrightarrow$ a=b. $\qquad\square$

Remark 3.23. One can cancel the common factor on both sides of equation **exactly** when R is an ID.

Overview Fields $\subsetneq$ Euclidean Domain (ED) $\subsetneq$ Principal Ideal Domain (PID) $\subsetneq$ Unique Factorization Domain (UFD) $\subsetneq$ Itegral Domain (ID)
(HW: If R is a field, then R is integral domain.)

Lemma 3.24. *Let R be ID with $\mid R \mid < \infty$. Then R is unital.*

  Proof: Let a $\in R$ be nonzero. Then $\{a, a^2, \cdots\}(\subset R)$ must repeat, since $\mid R \mid < \infty$. Say $a^m = a^n$ for some m>n.
Claim: $1_R = a^{m-n}$.
Proof of claim: $\forall x \in R$, let $xa^{m-n} = y$. Then $xa^m = xa^{m-n}a^n = ya^n = ya^m$. Then, by cancellation property, we have: x=y. Again by cancellation property and $xa^{m-n} = y$, $a^{m-n} = 1_R$. Hence, by the claim, $1_R = a^{m-n} \in R$. Therefore, R is unital. $\qquad\square$

Proposition 3.25. *Let R be an ID with $\mid R \mid < \infty$. Then R is a field.*

  Proof: Let $R \setminus \{0_R\} = \{r_1 = 1_R, r_2, \cdots, r_m\}$. Then for all a$\neq 0$ in R, consider

$$S := \{ar_1, ar_2, \cdots, ar_m\}$$

Suppose this set has repetitions, i.e.: $\exists i \neq j, s.t. : ar_i = ar_j$. Then by cancellation property, $r_i = r_j$. Then i=j.

Hence, S has no repetitions. Then S=$R \setminus \{0_R\}$. In particular, $\exists r_l, s.t. : ar_l = r_1 = 1_R$, and hence, a$\in U(R)$. $\qquad\qquad\qquad\square$

3.4 ideal

Motivation A ring analogue of normal subgroup.

Definition 3.26 (Ideal)**.** Let R be a ring. We say I $\subset R$ an **ideal** if

- I is an additive subgroup of (R,+)

- For all x $\in R$, i $\in I$, we have xi,ix $\in I$.

Remark 3.27.   • We write I ◁ R if I is an ideal

- If I ◁ R, then I $\leq$ R automatically. (REASON: Take x $\in$ I, then ax=xa$\in I$)

Example 3.28.   • (Nonexample:)$\mathbb{Z} \leq \mathbb{Q}$ is **NOT** an ideal, since 2$\in \mathbb{Z}$ and $\frac{1}{3} \in \mathbb{Q}$, **BUT** $2 \cdot \frac{1}{3} = \frac{2}{3} \notin \mathbb{Z}$

- R=$\mathbb{Z}$, I=$n\mathbb{Z}$. Take na$\in I$, m$\in R$. Then $(na)m = n(am) \in I$

- R=$\mathbb{Z}[x]$, I=$\{p(x) \in R \mid p(0) = 0\}$(=\{polynomials with no constant term\})
  (HW: Check I is subring.)
  Now take $p(x) = a_n x^n + \cdots + a_1 x \in I$ and $q(x) = b_m x^m + \cdots + b_1 x + b_0 \in R$. Then $p(x)q(x) = a_n b_m x^{n+m} + \cdots + a_1 b_0 x \in I$.
  (Alternatingly, $p(0)q(0) = 0 \cdot q(0) = 0$, and hence p(x)q(x)$\in I$)

- $R = \mathbb{Z}[x]$, $I = \{p(x) \in R \mid$constant term is an even integer$\}$
  e.g.: $1 + x \notin I, 10002 + x^2 \notin I, (0+)x^3 + 15x^5 \in I$
  (HW: Check that this is an ideal)

Question: For any R, how to construct I?

Definition 3.29. Let R be a ring, V $\subset$ R subset. Then the **ideal generated by V** is <V>, which is the smallest ideal in R containing all elements in V.

Example 3.30.   • R=$\mathbb{Z}$, V=n. What's <V>=<n>?
  For any ideal I s.t.: $n \in I$, then $n + \cdots + n$ and $(-n) + \cdots + (-n) \in S$, since I is a subgroup of (R,+). Then $n\mathbb{Z} \subset I$. But $n\mathbb{Z}$ itself is an ideal. So <n>=$n\mathbb{Z}$

- How about general case? Let R be a unital commutative ring, and let $V = \{a_1, \cdots, a_n\}$ is a finite set. Then <V>=<$a_1, \cdots, a_n$>=$\{r_1 a_1 + \cdots + r_n a_n \mid r_1, \cdots, r_n \in R\}$=:S

  1. Check S contains $a_1, \cdots, a_n$, e.g.: $a_2 = 0a_1 + 1a_2 + 0a_3 + \cdots + 0a_n \in S$

  2. Check S ◁ R. (Check $s \in S$, $r \in R$, then $s \cdot r \in S$)

3. For any ideal I satisfying: $a_1, \cdots, a_n \in I$, check $S \subset I$
   (**Proof:** $a_1, \cdots, a_n \in S$. Then by the definition of ideal, $a_1 r_1, \cdots, a_n r_n \in I$. Since I is an additive subgroup, then $a_1 r_1 + \cdots + a_n r_n \in I$)

- R= $\mathbb{Z}[x]$, $I = <2, x> = \{2p(x) + xq(x) \mid p(x), q(x) \in R\} = \{$all polynomials with even constant term$\}$.

Lemma 3.31. *Suppose $I_1, \cdots, I_k \triangleleft R$, then*

1. $I_1 + \cdots + I_k := \{i_1 + \cdots + i_k \mid i_r \in I_r\} \triangleleft R$

2. $\bigcap_{i=1}^{k} I_i \triangleleft R$

Proof:

1. Let $(i_1 + \cdots + i_k)$ and $(i_1' + \cdots + i_k') \in I_1 + \cdots + I_k$. Then $(-i_r) \in I_r \ \forall r = 1, \cdots, k$.
   $\Rightarrow -(i_1 + \cdots + i_k) = (-i_1) + \cdots + (-i_k) \in I_1 + \cdots + I_k$
   Also, $(i_1 + \cdots + i_k) + (i_1' + \cdots + i_k') = (i_1 + i_1') + \cdots + (i_k + i_k') \in I_1 + \cdots + I_k$
   Hence, $I_1 + \cdots + I_k$ is an additive subgroup.
   Now, take any $r \in R$, then $r \cdot (i_1 + \cdots + i_k) = ri_1 + \cdots + ri_k \in I_1 + \cdots + I_k$. Similarly, $(i_1 + \cdots + i_k) \cdot r \in I_1 + \cdots + I_k$

2. Left to the readers. $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad$ $\square$

Example 3.32. $R = \mathbb{Z}$, $I_r = a_r \mathbb{Z}$ where $a_r \in \mathbb{N}$
Then $I_1 + \cdots + I_k = gcd(a_1, \cdots, a_k)\mathbb{Z}$ and $I_1 \bigcap \cdots \bigcap I_k = lcm(a_1, \cdots, a_k)\mathbb{Z}$

Definition 3.33 (Principle Ideal Domain).   1. Let R be a commutative ring. An ideal $I \triangleleft R$ is called **principal** if I=<a> for some a$\in R$.

2. Let R be an ID. We say r is a **principal ideal domain(PID)** if all ideals in R are principal.

Proposition 3.34. *Let $\phi : R \to S$ be a ring homomorphism. Then $ker\phi \triangleleft R$.*

Proof: We only need to show that $\forall r \in R, i \in ker\phi \Rightarrow ri, ir \in ker\phi$, since we know $ker\phi \leq R$ already.
Indeed, $\phi(ri) = \phi(r)\phi(i) = \phi(r) \cdot 0_S = 0_S$. Hence, $ri \in ker\phi$. Similarly for ir. $\qquad\qquad$ $\square$

3.5 Quotient Ring

Definition 3.35 (Quotient Ring). Let R be a ring, and $I \triangleleft R$ ideal. Consider the collection of left cosets of (R,+), $R/I := \{r + I \mid r \in R\}$ (Then (R/I,+) has a group structure) with the operations:
$(r_1 + I) +_{R/I} (r_2 + I) := (r_1 + r_2) + I$
$(r_1 + I) \cdot_{R/I} (r_2 + I) := (r_1 \cdot r_1) + I$.
Then $(R/I, +_{R/I}, \cdot_{R/I})$ is a ring, and it's called the **quotient ring of R** by I.

Check:

- $\cdot_{R/I}$ is well-defined:

  Take $r_1 + I = r_1' + I$ and $r_2 + I = r_2' + I$ (*).

  (*) $\Leftrightarrow r_1 - r_1' \in I$ and $r_2 - r_2' \in I \Rightarrow r_1' = r_1 + i_1$ and $r_2' = r_2 + i_2$ for some $i_1, i_2 \in I \Rightarrow$
  $r_1'r_2' = r_1r_2 + r_1i_2 + i_1r_2 + i_1i_2 \Rightarrow r_1'r_2' - r_1r_2 \in I \Leftrightarrow r_1'r_2' + I = r_1r_2 + I \Leftrightarrow (r_1' + I) \cdot_{R/I} (r_2' + I) =$
  $(r_1 + I) \cdot_{R/I} (r_2 + I)$

- $+_{R/I}$ is well-defined by quotient theory

- Check $(R/I, +_{R/I}, \cdot_{R/I})$ is associative:

  $(r_1 + I) \cdot_{R/I} ((r_2 + I) \cdot_{R/I} (r_3 + I)) = ((r_1 + I) \cdot_{R/I} (r_2 + I)) \cdot_{R/I} (r_3 + I)$

- Check $(R/I, +_{R/I}, \cdot_{R/I})$ is distributive, e.g.:

  $(r_1 + I) \cdot_{R/I} ((r_2 + I) +_{R/I} (r_3 + I)) = (r_1 + I) \cdot_{R/I} (r_2 + I) +_{R/I} (r_1 + I) \cdot_{R/I} (r_3 + I)$ $\square$

Easy Exercise

- If R is commutative, then R/I is commutative.

- If $1_R \in R$ unital, then $1_{R/I} := 1_R + I$ unital.

Example 3.36.   - $\mathbb{R}[x]/\langle x^2 - 1 \rangle = \{p(x) + \langle x^2 - 1 \rangle \mid p(x) \in \mathbb{R}[x]\} = \{(ax + b) + \langle x^2 - 1 \rangle \mid a, b \in \mathbb{R}\}$
  (shorthand: $\overline{p(x)} := p(x) + \langle x^2 - 1 \rangle$)

$$\overline{x - 1}, \overline{x + 1} \in \mathbb{R}[x]/\langle x^2 - 1 \rangle \quad and \quad (x - 1) \cdot (x + 1) = x^2 - 1 \in \langle x^2 - 1 \rangle$$

  $\Rightarrow$

$$\overline{x - 1} \cdot \overline{x + 1} = \overline{0} \in \mathbb{R}[x]/\langle x^2 - 1 \rangle$$

  $\therefore \overline{x - 1}, \overline{x + 1}$ are zerodivisors of $\mathbb{R}[x]/\langle x^2 - 1 \rangle$ and hence it is **NOT integral domain (ID)**
  (issue: $x^2 - 1 = (x + 1)(x - 1)$ is not irreducible in $\mathbb{R}[x]$)

- $\mathbb{R}[x]/\langle x^2 + 1 \rangle$. Then

$$\overline{x} \cdot \overline{x} = \overline{x^2} = \overline{x^2 - (x^2 + 1)} = \overline{-1} \Rightarrow (\overline{x})^2 = \overline{-1}$$

- $\mathbb{Z}/\langle 2, x \rangle = \{(a_n x^n + \cdots + a_1 x + a_0) + \langle 2, x \rangle\} = \{a_0 + \langle 2, x \rangle\} = \{0 + \langle 2, x \rangle, 1 + \langle 2, x \rangle\} \cong \mathbb{Z}_2$

Theorem 3.37 (First Isomorphism Theorem of Rings). *Let $\Phi : R \to S$ be a ring homomorphism. Then the map $\phi : R/ker\Phi \to im\Phi$ defined by $\phi(r + ker\Phi) := \Phi(r)$ is a well-defined ring isomorphism.*

  Proof:

- ϕ is well-defined, i.e.: if (*)"$r + ker\Phi = r' + ker\Phi$", then $\Phi(r) = \Phi(r')$.
  (Exercise: recall (*) $\Leftrightarrow r - r' \in ker\Phi$)

- ϕ is a ring homomorphism: only need to check $\phi(rr') = \phi(r)\phi(r')$, since we know ϕ is group homomorphism already by the 1_{st} isomorphism theorem of groups.

- ϕ is bijective as in the 1_{st} isomorphism theorem of groups.   □

Example 3.38.   • Let $\Phi : \mathbb{R}[x] \to \mathbb{C}$ be defined by $\Phi(p(x)) := p(i)$ where $(i = \sqrt{-1})$. Then ϕ
   is a homomorphism. (Exercise: $\phi(pq) = \phi(p)\phi(q)$)
   Now, $im\phi = \mathbb{C}$, e.g.: $\Phi(bx + a) := a + bi \in \mathbb{C}$ and

$$p(x) \in ker\Phi$$

$$\Leftrightarrow p(i) = 0$$

$$\Leftrightarrow \overline{p(i)} = p(i) = 0$$

$$\Leftrightarrow p(-i) = p(i) = 0$$

$$\Leftarrow (Factor\ theorem\ on\ \mathbb{C}[x]) \Rightarrow (x - i), (x - (-i)) \mid p(x)$$

$$\Leftrightarrow (x - i)(x + i) = x^2 + 1 \mid p(x)\ in\ \mathbb{R}[x]$$

   $\therefore\ ker\Phi = \{(x^2 + 1)q(x) \mid q(x) \in \mathbb{R}[x]\} = \langle x^2 + 1\rangle$.
   $\therefore 1_{st}$ isomorphism theorem says
$$\mathbb{R}[x]/\langle x^2 + 1\rangle \cong \mathbb{C}$$

- Let $\Phi : \mathbb{Z}[x] \to \mathbb{Z}_2$ with $\Phi := p(0)(\mathrm{mod}\ 2)$.
   Check that $im\Phi = \mathbb{Z}_2$, $ker\Phi = \langle 2, x\rangle$. (Exercise)

$$\Rightarrow\ \mathbb{Z}[x]/\langle 2, x\rangle \cong \mathbb{Z}_2$$

3.6   Chinese Remainder Theorem

Definition 3.39 (Product Ring). Let R_i be rings. Then the **product ring** $\Pi_i R_i (= R_1 \times \cdots \times R_k)$
has a ring structure given by

$$(r_1, \cdots, r_k) +_{\Pi_i R_i} (r'_1, \cdots, r'_k) := (r_1 + r'_1, \cdots, r_k + r'_k)$$

$$(r_1, \cdots, r_k) \cdot_{\Pi_i R_i} (r'_1, \cdots, r'_k) := (r_1 r'_1, \cdots, r_k r'_k)$$

Remark 3.40. If R_1, R_2 are ID, then $R_1 \times R_2$ is **NOT** an ID:

$$(r_1, 0) \cdot (0, r_2) = (0, 0) = 0_{R_1 \times R_2}$$

Definition 3.41. Let R be a commutative ring. We say $I_1, I_2 \triangleleft R$ **coprime** if $I_1 + I_2 = R$.

Example 3.42. Let $R = \mathbb{Z}$ and $I_1 = \langle m\rangle = m\mathbb{Z}$, $I_2 = \langle n\rangle = n\mathbb{Z}$.
Then $I_1 + I_2 := \{mp + nq \mid p, q \in \mathbb{Z}\} = \langle gcd(m, n)\rangle$. (Exercise: prove the second equality.)
$\therefore\ I_1$ & I_2 are coprime $\Leftrightarrow I_1 + I_2 = \mathbb{Z} \Leftrightarrow \langle gcd(m, n)\rangle = \mathbb{Z} \Leftrightarrow gcd(m, n) = 1 \Leftrightarrow$ m,n coprime as
integers.

The above example generalizes our understanding of "coprime" frome $\mathbb{Z}$ to any commutative ring R: Two elements r_1, r_2 are coprime in R means $\langle r_1 \rangle + \langle r_2 \rangle = R$.

Theorem 3.43. *Let R be commutative unital, and $I_1, \cdots, I_k \lhd R, s.t. : I_i, I_j$ are pairwise coprime. Then we have a ring isomorphism*

$$\phi : R/I_1 \cap \cdots \cap I_k \to R/I_1 \times \cdots \times R/I_k \quad defined \ by \quad \phi(r + I_1 \cap \cdots \cap I_k) := (r + I_1, \cdots, r + I_k).$$

Proof: Let $\Phi : R \to R/I_1 \times \cdots \times R/I_k$ be a ring homomorphism with $\Phi(r) := (r + I_1, \cdots, r + I_k)$. By the first isomorphism theorem, we need to show:

1. $ker\Phi = I_1 \cap \cdots \cap I_k$ and

2. $im\Phi = R/I_1 \times \cdots \times R/I_k$

1. $r \in ker\Phi \Leftrightarrow r + I_l = 0 + I_l \ \forall l \Leftrightarrow r - 0 \in I_l \ \forall l = 1, \cdots, k \Leftrightarrow r \in I_1 \cap \cdots \cap I_k$

2. Fix $j \in \{1, \cdots, k\}$. Since R is unital and I_j, I_i coprime for all $i \neq j$, so we have $z_i \in I_j \quad w_i \in I_i$, s.t.:

$$(*) : z_i + w_i = 1 \ (1 \in I_i + I_j = R) \ \forall i \neq j$$

Consider $1 = (1 - \Pi_{i \neq j} w_i) + (\Pi_{i \neq j} w_i)$.

$$Set \quad x_j := 1 - \Pi_{i \neq j} w_i = 1 - \Pi_{i \neq j}(1 - z_i) = sum \ of \ products \ of \ z_i's \ with \ no \ constants \in I_j$$

$$Set \quad y_j := \Pi_{i \neq j} w_i \in \bigcap_{i \neq j} I_i, \ since \ I_i \ ideals$$

$\therefore$ For each fixed j, we have

$$(**) : \quad 1 = x_j (\in I_j) + y_j (\in \bigcap_{i \neq j} I_i)$$

We already to check Φ is surjective:

For each $(u_1 + I_1, \cdots, u_k + I_k) \in R/I_1 \times \cdots \times R/I_k$, we **CLAIM** that $\Phi(u_1 y_1 + \cdots + u_k y_k) = (u_1 + I_1, \cdots, u_k + I_k)$:

REASON: $\Phi(u_1 y_1 + \cdots + u_k y_k) = (\cdots, u_1 y_1 + \cdots + u_k y_k + I_l, \cdots) = (u_1 + I_1, \cdots, u_k + I_k)$, since

$$(u_1 y_1 + \cdots + u_l y_l + \cdots + u_k y_k + I_l) = u_l y_l + I_l \stackrel{(**)}{=} u_l(1 - x_l) + I_l = u_l - u_l x_l + I_l = u_l + I_l$$

$\square$

Corollary 3.44. *Let $p_1, \cdots, p_n$ be distinct prime numbers. Then*

$$\mathbb{Z}/\langle p_1^{a_1} \cdots p_n^{a_n} \rangle \cong \mathbb{Z}/\langle p_1^{a_1} \rangle \times \cdots \times \mathbb{Z}/\langle p_n^{a_n} \rangle.$$

Therefore, for each $b_i \in \mathbb{Z}/\langle p_i^{a_i} \rangle \cong \mathbb{Z}_{p_i^{a_i}}$, there exists $x \in \mathbb{Z}, s.t. : \Phi(x) = (b_1 + \langle p_1^{a_1} \rangle, \cdots, b_n +$

$\langle p_n^{a_n} \rangle)$ $x \equiv b_i \pmod{\mathbb{Z}_{p_i^{a_i}}}$ for all i.

3.7   Prime and Maximal Ideals

Motivation: Define "prime" in any commutative unital R.

(Kummer, mid 1800's): Rather thaan studying $r \in R$, study ideals $I \lhd R$.

Basic case: $R = \mathbb{Z}$

All ideals $I \lhd \mathbb{Z}$ are of the form $I = \langle n \rangle$. So study $\langle n \rangle$ instead of n.

e.g.: in $R = \mathbb{Z}$,

$$(\langle p \rangle, \langle q \rangle \ coprime \ (as \ ideals)) \Leftrightarrow (p, q \ are \ coprime \ (as \ integers))$$

Definition 3.45. Let R be a commutative, unital ring. We say a proper $(I \neq R)$ ideal $I \lhd R$ is

1. **prime** if for all a,b $\in$ R such that ab $\in$ R, then we must have $a \in I \ or \ b \in R$;

2. **maximal** if for all ideals $J \lhd R$ s.t. $I \subset J \subset R$, then J=I or J=R.

Example 3.46.   • R=$\mathbb{Z}$. Let's check

$$(p \in \mathbb{Z}, \ prime) \Leftrightarrow (I = \langle p \rangle \ is \ a \ prime \ ideal)$$

Proof: Let $a, b \in \mathbb{Z}$ s.t.

$$ab \in \langle p \rangle \Leftrightarrow ab = pk \ for \ some \ k \in \mathbb{Z} \Leftrightarrow p \mid ab \Leftrightarrow p \mid a \ or \ p \mid b \Leftrightarrow a \in \mathbb{Z} \ or \ b \in \mathbb{Z}$$

(**Exercise:** $\langle n \rangle$ is a maximal ideal $\Leftrightarrow$ n is prime $\Leftrightarrow$ $\langle n \rangle$ is prime)

(**Nonexample:** $\langle 6 \rangle$ is **NOT** maximal since $\langle 6 \rangle \subsetneq \langle 2 \rangle \subsetneq \mathbb{Z}$)

• R=$\mathbb{Z}_{12}$. All the ideals of R are:

$I_0 = \{0\}, \ I_1 = \{0, 2, 4, 6, 8, 10\} (maximal \ and \ prime), \ I_2 = \{0, 3, 6, 9\} (maximal \ and \ prime),$

$I_3 = \{0, 4, 8\} (\textbf{NOT} \ prime, \ since \ 2 \cdot 2 = 4), \ I_4 = \{0, 6\} (\textbf{NOT} \ prime, \ since \ 2 \cdot 3 = 6), \ I_5 = R$

• $R = \mathbb{Z}[x]$. $I = \langle x \rangle$=polynomials with 0 constant term is a prime ideal:

Take $p(x) = a_0 + a_1 x + \cdots + a_m x^m$, $q(x) = b_0 + b_1 x + \cdots + b_n x^n \in I$. Then

$$pq \in I \Leftrightarrow a_0 b_0 = 0 \Leftrightarrow a_0 = 0 \ or \ b_0 = 0 \Leftrightarrow p(x) \in I \ or \ q(x) \in I$$

But I is **NOT** maximal:
$$I \subsetneq \langle 2, x \rangle \subsetneq R$$

Proposition 3.47. *Let R be commutative unital, and $I \lhd R$. Then*

1. I is prime $\Leftrightarrow$ R/I is ID

2. I is maximal $\Leftrightarrow$ R/I is a field

Example 3.46.(3) revisited: Consider $\langle x \rangle \subset \mathbb{Z}[x]$. Then

$$R/I = \mathbb{Z}[x]/\langle x \rangle \cong \mathbb{Z}.$$

$\mathbb{Z}$ is ID $\Leftrightarrow \langle x \rangle$ is prime

$\mathbb{Z}$ is **NOT** a field $\Leftrightarrow \langle x \rangle$ is **NOT** maximal

Proof:

1. Let (a+I), (b+I) $\in R/I$. Then (*): $(a + I) \cdot (b + I) = 0 + I \Leftrightarrow ab \in I$.
   If I is prime, then (*) says

   $$(a+I)(b+I) = 0_{R/I} \Leftrightarrow ab \in I \Leftrightarrow a \in I \text{ or } b \in I \Leftrightarrow a+I = 0_{R/I} \text{ or } b+I = 0_{R/I} \Leftrightarrow R/I \text{ is ID}$$

2. Suppose $I \subset J \subset R$ for some $J \triangleleft R$. Then $I/I \subset J/I \subset R/I$. (HW9: J/I $\triangleleft$ R/I)
   Recall: $\mathbb{F}$ is a field $\Leftrightarrow$ the only ideals of $\mathbb{F}$ are 0 and $\mathbb{F}$
   (**Proof** of "Recall": Let $I \triangleleft \mathbb{F}$ be nonzero. Take $0 \neq a \in I$. Then $1 = a \cdot a^{-1} \in I \Rightarrow \forall x \in \mathbb{F}$,
   $x = x \cdot 1 \in I$)
   Therefore,

   $$\mathbb{F} = R/I \text{ is a field} \Leftrightarrow J/I \triangleleft R/I \text{ must be } \{0\} \text{ or } R/I \Leftrightarrow J = \{0\} \text{ or } J = R$$

   $\square$

Corollary 3.48. *Let R be commutative unital. Then $I \triangleleft R$ is maximal $\Rightarrow I \triangleleft R$ is prime.*

Proof. All fields are IDs. $\square$

Definition 3.49. Let R be commutative unital. We say $a \in R$ is **prime** if $\langle a \rangle$ is a prime ideal.
(Generalization of "prime number" in $R = \mathbb{Z}$)

Definition 3.50. Let $a, b \in R$ (commutative unital). We say a **divides** b (or $a \mid b$ in short) if
$\langle b \rangle \subseteq \langle a \rangle$. (or equivalently, $b \in \langle a \rangle$, or $\exists x \in R$, s.t.: $ax = b$.)

We say a and b are **associates** ($a \sim b$) if ($a \mid b$ & $b \mid a$), (or $\langle a \rangle = \langle b \rangle$).

Lemma 3.51. *Suppose R is an integral domain. Then $(a \sim b) \Leftrightarrow \exists$ unit $x \in U(R)$ s.t.: $a = xb$.*

Proof:
($\Leftarrow$): $a = xb \Rightarrow a \in \langle b \rangle \Rightarrow \langle a \rangle \subseteq \langle b \rangle$,
AND $x^{-1}a = b \Rightarrow \langle b \rangle \subseteq \langle a \rangle$ Similarly.
$\therefore \langle a \rangle = \langle b \rangle$
($\Rightarrow$): Suppose $a \sim b$, i.e.: $\langle a \rangle = \langle b \rangle$. Then we have $a \in \langle b \rangle$.
$\Rightarrow a = p \cdot b$ for $p \in R$. Similarly, $b \in \langle a \rangle \Rightarrow b = q \cdot a$ ($q \in R$)
$\Rightarrow a = p \cdot q \cdot a \xrightarrow{\text{cancellation property}} 1_R = p \cdot q \Rightarrow p, q \in U(R)$. $\square$

3.8 Principal Ideal Domain

Recall: I maximal $\Rightarrow I$ prime for R unital commutative.

Converse NOT holds in general (e.g. $\mathbb{Z}[x] = R$).

Goal: Study R s.t.: (I maximal) $\Leftrightarrow$ (I prime).

Definition 3.52 (Principal Integral Domain)**.** Let R be ID. We call R a **principal ideal domain (PID)** if all ideals $I \triangleleft R$ are principal, i.e.: all I are of the form $I = \langle a \rangle$.

Example 3.53. • $\mathbb{Z}$ is PID.

    Proof: Let I be a nonzero ideal of $\mathbb{Z}$. (Otherwise $I = \{0\} = \langle 0 \rangle$)

    Let $\mu > 0$ be the smallest positive integer in I.

    <u>Claim:</u> $I = \langle \mu \rangle$.

    Suppose on contrary that $I \setminus \langle \mu \rangle$ is nonempty. Take $\lambda > 0$ be the smallest integer in $I \setminus \langle \mu \rangle$. By minimality of μ, we have $\lambda > \mu$. Then $\lambda - \mu \in I$ and $\lambda - \mu \notin \langle \mu \rangle$, since otherwise $(\lambda - \mu) + \mu = \lambda \in \langle \mu \rangle$, contradicting the choice of λ.

    Hence, $(\lambda - \mu)$ is an element in $I \setminus \langle \mu \rangle$, which is smaller than λ, contradicting the minimality of λ in $I \setminus \langle \mu \rangle$.

    $\therefore I = \langle \mu \rangle$ and hence $\mathbb{Z}$ is PID.

• $\mathbb{Z}[x]$ is <u>NOT</u> PID, since $I = \langle 2, x \rangle \neq \langle p \rangle$ for any $p \in \mathbb{Z}[x]$ which could be proven in HW9.

• For any field $\mathbb{F}$ (e.g. $\mathbb{F} = \mathbb{Q}, \mathbb{R}, \mathbb{C}$), $\mathbb{F}[x]$ is PID.

    Proof: Let I be an ideal of $\mathbb{F}$ which is nonzero. For any elements $p(x) \in I$, we can multiply unit $u \in \mathbb{F}$ st.: $u \cdot p(x) \in I$ is a **monic** polynomial (leading power coefficient is 1). Take a monic polynomial $p(x) \in I$ of smallest possible degree in I.

    Claim: $I = \langle p(x) \rangle$.

    Proof of claim: Suppose on contrary. Take monic $q(x) \in I \setminus \langle p(x) \rangle$ such that q is of smallest positive degree in $I \setminus \langle p \rangle$. Then $\deg(p) \leq \deg(q)$ by the minimality of $\deg(p)$ in I.

    Let $h(x) := q(x) - x^{\deg(q) - \deg(p)} \cdot p(x)$. Then:

        – $h(x) \in I$

        – $h(x) \notin \langle p(x) \rangle$

        – $\deg(h) < \deg(q)$

    Then $h \in I \setminus \langle p \rangle$ with degree $< \deg(q)$. Contradict! $\qquad\qquad\square$

Note: We have division algorithm for $\mathbb{F}[x]$ just like the case of $\mathbb{Z}$:

$$p(x) = q(x)a(x) + r(x) \quad (\deg(r) < \deg(q))$$

and so we have Euclidean algorithm to find $\gcd(p(x), q(x))$ in $\mathbb{F}[x]$

3.9   Irreducible Elements and Unique Factorization Domain

Definition 3.54 (Irreducibility). Let R be unital commutative ring. An element $a \in R$ is **irreducible** if the followings hold:

Whenever $\langle a \rangle \subseteq \langle b \rangle \subseteq R$ for some $b \in R$, we have $\langle a \rangle = \langle b \rangle$ or $\langle b \rangle = R$, i.e.: $\langle a \rangle$ is maximal among all principal ideals of R.

To understand what it means for $a \in R$ irreducible:

Lemma 3.55. *Let R be ID. Then:*

($a \in R$ irreducible) $\Leftrightarrow$ (whenever $a = xy$, we have $a \sim x$ or $a \sim y$)

(Recall $a \sim x \Leftrightarrow a = ux$ for unit u)

Proof: ($a \in R$ is irreducible $\Rightarrow (a = xy \Rightarrow a \sim x$ or $a \sim y)$) :

Suppose $a = xy$. Then $a \in \langle x \rangle$ and $a \in \langle y \rangle$. $\Rightarrow \langle a \rangle \subseteq \langle x \rangle$ and $\langle a \rangle \subseteq \langle y \rangle$. By definition of "a irreducible", $\langle a \rangle = \langle x \rangle$ or $\langle x \rangle = R$.

- $\langle a \rangle = \langle x \rangle \Leftrightarrow a \sim x$, then we're done

- $\langle x \rangle = R \Rightarrow \exists x' \in R$, s.t.: $xx' = 1 \in \langle x \rangle = R \Rightarrow x$ is a unit $\Rightarrow a \sim y$.

$((a = xy \Rightarrow a \sim x$ or $a \sim y) \Rightarrow (a \in R$ is irreducible)) :

Suppose $\langle a \rangle \subseteq \langle b \rangle$ for some b. (WTS: $\langle b \rangle = \langle a \rangle$ or $\langle b \rangle = R$).

$\Rightarrow a = b \cdot b'$ for some $b' \in R \Rightarrow a \sim b$ or $a \sim b'$. Then

- $\langle a \rangle = \langle b \rangle$, then we're done

- $a = ub'$ for some unit $u \Rightarrow u = b$ by cancellation $\Rightarrow \langle b \rangle = R$, then we're done.   □

Remark 3.56. As a corollary of lemma, a is irreducible $\Leftrightarrow \nexists b, c$ such that $a = bc$, and $b, c \notin U(R)$ (non-units).

$\therefore$ we can "factorize" a into "smaller" elements b & c. So we can factorize all $a \in R$ into product of irreducibles.

$\therefore$ Irreducible elements are "building blocks" of R.

Proposition 3.57. *Let R be ID. If $0 \neq a \in R$ is prime, then a is irreducible.*

Proof: Suppose $a = xy$ (WTS: $a \sim x$ or $a \sim y$)

Then $a \in \langle x \rangle$ and $a \in \langle y \rangle \Rightarrow \langle a \rangle \subseteq \langle x \rangle$ and $\langle a \rangle \subseteq \langle y \rangle$

On the other hand, we have $a \mid xy \xrightarrow{(a \text{ prime})} a \mid x$ or $a \mid y$

WLOG, assume $a \mid x$. Then $aa' = x \Rightarrow \langle x \rangle \subseteq \langle a \rangle \Rightarrow \langle a \rangle = \langle x \rangle$.   □

CONCLUSION: R PID.

$\langle a \rangle$ maximal $\Leftrightarrow \langle a \rangle$ prime $\Leftrightarrow a$ is irreducible

(e.g.: $\langle x \rangle$ in $\mathbb{Z}[x]$ is prime but <u>NOT</u> maximal ;

3 in $\mathbb{Z}[\sqrt{-5}]$ is irreducible but $\overset{\langle 3 \rangle}{\text{NOT}}$ prime)

WRAP-UP:

- All $a \in R$ can be factorized into irreducibles

- Primes are irreducibles

If we want $a \in R$ to be factorized into primes, we need "all irreducibles are primes" instead.

Proposition 3.58. *Let R be PID. Then all irreducibles are primes.*

Proof: Let $x \in R$ be irreducible. Then $\langle x \rangle$ is maximal among all ideals of the form $\langle b \rangle$.
Since all ideals are of the form $\langle b \rangle$ in PID, then $\langle x \rangle$ is a maximal ideal.
$\Leftrightarrow \langle x \rangle$ is prime ideal $\Rightarrow x$ is prime.   □

Definition 3.59. Let R be ID. We say R is a **factorization domain** if for all $0 \neq x \in R$, there exists irreducible elements $x_1, \ldots, x_r$ such that $x \sim x_1 x_2 \ldots x_r$ (or $x = u x_1 x_2 \ldots x_r$ for some unit $u \in U(R)$), i.e.: one can factorize any x into a <u>finite</u> product of irreducibles.

Theorem 3.60. *R is PID $\Rightarrow R$ is factorization domain.*

Definition 3.61. Let R be unital commutative. We say R has the **ascending chain condition** on **principal ideals** (ACCP) if all $I_1 \subseteq I_2 \subseteq I_3 \subseteq \ldots$ ($I_n \triangleleft R$ principal ideals), there exists N such that $I_N = I_{N+1} = I_{N+2} = \ldots$

Lemma 3.62. *All PIDs satisfies (ACCP).*

Proof: Suppose $I_1 \subseteq I_2 \subseteq I_3 \subseteq \ldots$ principal ideals in R. Consider $I := \bigcup_{n=1}^{\infty} I_n$ (HW 10: $I \triangleleft R$).
So $I = \langle r \rangle$ for some $r \in R$.
Since R is PID, then $r \in I_M$ for $M \in \mathbb{N} \Rightarrow r \in I_n \quad \forall n \geq M$.
$\Rightarrow I = \langle r \rangle \subseteq I_n \quad \forall n \geq M$.
Since $I_n \subseteq I$ by defn of I, then $I_n = I \quad \forall n \geq M$.   □

Lemma 3.63. *If R satisfies (ACCP), then R is a factorization domain.*
(Consequently, all PIDs are factorization domains)

Proof: Let $F := \{x \in R \mid x \sim x_1 \ldots x_r, x_i \text{ irreducibles}\}$ (W.T.S.: $R = F$).
Then: (a) $1 \in F$ (by convention); (b) all irreducibles $x \in R$ are in F ($r = 1$); (c) F is closed under multiplication.
Suppose on contrary, $\exists x_0 \in R \setminus F$. Then by (b), x_0 is reducible.
Then $x_0 = y_0 z_0$ where y_0, z_0 are <u>NOT</u> units. ($x_0 \nsim y_0$ & $x_0 \nsim z_0$)
$\Leftrightarrow \langle x_0 \rangle \neq \langle y_0 \rangle$
By (c), either $y_0 \in R \setminus F$ or $z_0 \in R \setminus F$ (or both). WLOG let $x_1 := y_0 \in R \setminus F$. Hence, $x_0 = x_1 z_0 \Rightarrow x_0 \in \langle x_1 \rangle \Rightarrow \langle x_0 \rangle \subsetneq \langle x_1 \rangle$ (and $\langle x_1 \rangle \neq R$, since x_1 is NOT unit).
Continue same argument on x_1, we have $\langle x_0 \rangle \subsetneq \langle x_1 \rangle \subsetneq \langle x_2 \rangle \subsetneq \ldots$ contradicting (ACCP) which says $\exists N$ s.t.: $\langle x_N \rangle = \langle x_{N+1} \rangle = \ldots$
$\therefore$ There's no such $x_0 \in R \setminus F$ and hence $R = F$.   □

CONCLUSION: In PID, one can factorize any $x \in R$ into finite product of irreducibles.
Q: Is the factorization unique?

Proposition 3.64. *Let R be an ID. Then any factorizations of $x \in R$ into PRIME factors are unique (up to permutation and units), i.e.: if $x_1 \ldots x_r \sim y_1 \ldots y_s$ (x_i, y_j nonunit PRIMES). Then: (1) $r = s$, and (2) $\exists \sigma \in S_r$ s.t.: $x_i \sim y_{\sigma(i)}$*

(Consequently, if R is PID, then we can factorize any $x \in R$ uniquely into (irreducibles $\Leftrightarrow$ primes).)

Proof: We'll prove the following stronger statement:

Let $x_1, \ldots, x_r$ primes, $y_1, \ldots, y_n$ irreducibles (e.g.: y_j primes) s.t.:

$(*) : x_1 x_2 \ldots x_r \sim y_1 y_2 \ldots y_n$, then $r = n$ & $\exists \sigma \in S_r$ s.t.: $x_i \sim y_{\sigma(i)}$

Proof by induction on r.

Basic Case: $r = 0$, then the left-hand-side $= 1$. $\therefore y_1 y_2 \ldots y_n \sim 1$

$\therefore y_j$ units $\forall j$. Since we assume x_i, y_j nonunits, must have $n = 0$.

Inductive Step: Suppose $(*)$ holds for $0 \le r \le s-1$. Consider $r = s$ and $u x_1 \ldots x_{s-1} x_s = y_1 y_2 \ldots y_n$

with u unit. (x_i primes, y_j irreducible, nonunits)

$\Rightarrow x_s \mid y_1 y_2 \ldots y_n \Rightarrow x_s \mid y_j$ for some j. $\Rightarrow y_j = x_s \cdot a$ for some $a \in R$.

$\Rightarrow y_j \sim x_s$ or $y_j \sim a \xleftarrow{\;y_j \text{ irreducible}\;} y_j \sim x_s \Rightarrow x_1 \ldots x_{s-1} \sim y_1 \ldots y_{j-1} y_{j+1} \ldots y_n$

Then by inductive hypothesis, $s - 1 = n - 1$ and $\exists \sigma \in S_{s-1}$ s.t.: $x_i \sim y_{\sigma(i)} \forall i$

$\therefore$ Let $\sigma' \in S_s$ by $\sigma' := \begin{cases} \sigma(i), \text{if } i = 1, \ldots, s-1 \\ j, \text{if } i = s \end{cases}$

$\therefore x_1 \ldots x_{s-1} x_s \sim y_1 \ldots y_n$ implies $\exists \sigma' \in S_s$ s.t.: $x_i \sim y_{\sigma'(i)} \quad \forall i = 1, \ldots, s.$ $\square$

Corollary 3.65. *Let R be PID. Then $\forall x \in R, x$ can be uniquely factorized into a finite product of irreducibles ($\Leftrightarrow$ primes).*

(This generalizes the fundamental thm of arithmetic from $R = \mathbb{Z}$ to any PID)

Definition 3.66. (UFD) Let R be ID. We say R is a **unique factorization domain** (UFD) if we have unique factorization into irreducible elements in R.

(Of course: R is PID $\Rightarrow R$ is UFD).

CONCLUSION:

ID $\supsetneq$ UFD $\supsetneq$ PID $\supsetneq$ ED (Euclidean domain) $\supsetneq$ Fields

(eg.: $\mathbb{Z}[\sqrt{-5}]$ is an ID (HW9: $(2 + \sqrt{-5}) \cdot (2 - \sqrt{-5}) = 3 \cdot 3$ is a non-unique factorization into irreducibles) ; $\mathbb{Z}[x]$ is a UFD (HW9: NOT PID ; later: it's UFD (Gauss' Lemma)) ; $\mathbb{Q}[x]$ is an ED.)

As in the case of PID, one has:

Proposition: Let R be UFD. Then p is prime iff p is irreducible.

Proof: p prime $\Rightarrow p$ irreducible $\forall$ ID. Now suppose p is irreducible. Consider $p \mid ab$ i.e.: $pc = ab$ for some $c \in R$. Factorize $a = a_1 \ldots a_x$ and $b = b_1 \ldots b_y$ into irreducibles. Then UF implies that either a_i or b_j associates p. Thus $p \mid a$ or $p \mid b$. $\square$

Definition 3.67. (Euclidean Function) Let R be ID. A **Euclidean function** on R is $N : R \backslash \{0\} \to \mathbb{N} \backslash \{0\}$ s.t.: $\forall a, b \in R$, either (i) $b \mid a$, or (ii) $\exists q \in R, r \in R \backslash \{0\}$, s.t.: $a = bq + r$ with $N(r) < N(b)$.

Example 3.68.   • $R = \mathbb{Z}$, $N(m) := |m|$. Then (ii) is simply the division algorithm for integers.

   • $R = \mathbb{F}[x]$, $N(p(x)) := \deg(p)$. Then (ii) is division algorithm for polynomials.

Definition 3.69. (Euclidean Domain) An ID R is a **Euclidean Domain** if R has (at least) one Euclidean function.

So $\mathbb{Z}, \mathbb{F}[x]$ are EDs.

If the Euclidean function ("norm function") N satisfies $N(ab) = N(a)N(b) \quad \forall a, b \in R$ we say N is **multiplicative**.

(eg.: $R = \mathbb{Z}, N(a) = |a|$ (multiplicative) ; $R = \mathbb{F}[x], N(p) = \deg(p)$ (**NOT** multiplicative))

Proposition 3.70. $\mathbb{Z}[i]$ *is a Euclidean domain with* $N(a+bi) = a^2+b^2, \forall a, b \in \mathbb{Q}$. *($\mathbb{Z}[i] := \{a+bi \mid a, b \in \mathbb{Z}\}$ is called* ***Gaussian integers****)*

Proof: *Let* $\alpha, \beta \in \mathbb{Z}[i]$. *Assume* $\beta \neq 0$. *Then consider* $\alpha/\beta = c_1 + c_2 i, c_i \in \mathbb{Q}$. *Let* $n_1, n_2 \in \mathbb{Z}$ *be the closest integers of* $c_1, c_2 \in \mathbb{Q}$ *(so that* $|n_i - c_i| \leq \frac{1}{2}$*), and let* $q := n_1 + n_2 i$. *Hence,* $\alpha/\beta = q + ((c_1 - n_1) + (c_2 - n_2)i)$.

$\Rightarrow \alpha = \beta q + ((c_1 - n_1) + (c_2 - n_2)i)\beta$. *Then* $N[((c_1-n_1)+(c_2-n_2)i)\beta] = N((c_1-n_1)+(c_2-n_2)i) \cdot N(\beta)$ $\leq [(\frac{1}{2})^2 + (\frac{1}{2})^2]N(\beta) = \frac{1}{2}N(\beta) < N(\beta)$. $\qquad \square$

Theorem 3.71. *If* $R \in ED$, *then* $R \in PID$.

   Therefore, $\mathbb{Z}[i]$ is PID.

Q: What are the (primes $\Leftrightarrow$ irreducibles) in the Gaussian integer $\mathbb{Z}[i]$?

(eg. 5 is prime in $\mathbb{Z}$, but $5 = (2 + i)(2 - i)$ not prime in $\mathbb{Z}[i]$)

Simple Observations:

   • If $n \in \mathbb{Z}$ is **NOT** prime in $\mathbb{Z}$, then n is <u>NOT</u> prime in $\mathbb{Z}[i]$.

   • $U(\mathbb{Z}[i]) = \{\pm 1, \pm i\}$. Therefore, $\alpha \in U(\mathbb{Z}[i]) \Leftrightarrow N(\alpha) = 1$.

   • If $N(\beta) = p$ is prime in $\mathbb{Z}$, then β is prime in $\mathbb{Z}[i]$. (e.g.: $\beta = 2 + i$ is prime in $\mathbb{Z}[i]$, since $N(\beta) = 5$ in $\mathbb{Z}$)
   $\Rightarrow p = N(\beta) = N(\gamma)N(\delta) \Rightarrow N(\gamma) = 1$ or $N(\delta) = 1$. $\Rightarrow \gamma$ or δ is a unit in $\mathbb{Z}[i] \Rightarrow \beta \sim \gamma$ or $\beta \sim \delta$, i.e.: β is irreducible $\Leftrightarrow$ prime.

Lemma 3.72. *Let* p *be a prime integer. Then either (i)* p *is a Gaussian prime in* $\mathbb{Z}[i]$*, or (ii)* $p = (a + bi)(a - bi)$ *for some* $(a + bi), (a - bi)$ *Gaussian primes.*

Proof: Suppose p is **NOT** a Gaussian prime. Then $p = \gamma \cdot \delta$ for nonunits γ, δ. $\Rightarrow p^2 = N(p) = N(\gamma)N(\delta) \Rightarrow N(\gamma) = N(\delta) = p$

$\Rightarrow \gamma, \delta$ Gaussian primes.

$p = (a + bi)(c + di) \Rightarrow \bar{p} = p = (a - bi)(c - di) \Rightarrow p = (a + bi)(a - bi)$. $\qquad \square$

Now let's study Gaussian primes of the form $a + bi$ $(a, b \neq 0)$.

Lemma 3.73. *Let $a + bi \in \mathbb{Z}[i]$ with $a, b \neq 0$. Then ($a + bi$ is Gaussian prime) $\Leftrightarrow$ ($a^2 + b^2 = p$, p prime integer).*

Proof: ($\Leftarrow$): is done in the simple observations.

($\Rightarrow$): Suppose $a + bi$ Gaussian prime. Then $a - bi = \overline{a + bi}$ is also Gaussian prime. Consider $(a + bi)(a - bi) = p_1 \dots p_r$, p_i prime in $\mathbb{Z}$, this is a factorization in $\mathbb{Z}[i]$. By $\mathbb{Z}[i]$ ED, $\mathbb{Z}[i]$ is PID $\Rightarrow$ UFD, $r \leq 2$. Suppose on contrary $r = 2$. Then $(a + bi)(a - bi) = p_1 p_2$.

By UFD again, p_1, p_2 must be (prime as irreducible) in $\mathbb{Z}[i]$.

$\therefore$ By UFD again, $p_1 \sim (a \pm bi)$, $p_2 \sim (a \mp bi) \Rightarrow p_1 = u \cdot (a \pm bi)$ where $u \in U(\mathbb{Z}[i]) = \{\pm 1, \pm i\}$, contradicting $a, b \neq 0$.

$\therefore r = 1 \Rightarrow (a + bi)(a - bi) = p_1 \Rightarrow a^2 + b^2 = p_1$. $\qquad \square$

CONCLUSION:

(i) If $n \in \mathbb{Z}$ is **NOT** a prime number in $\mathbb{Z}$, then n is <u>NOT</u> Gaussian prime.

(ii) Prime $p \in \mathbb{Z}$ is either a Gaussian prime or $p = (a + bi)(a - bi)$, $a \pm bi$ Gaussian primes.

(iii) $a + bi \in \mathbb{Z}[i], a, b \neq 0$ is Gaussian prime $\Leftrightarrow a^2 + b^2 = p$, $p \in \mathbb{Z}$ prime number.

Theorem 3.74. *Let p be a prime integer. Then p is Gaussian prime $\Leftrightarrow p \equiv 3 \pmod{4}$.*

Proof: ($\Leftarrow$) contrapositive. If p is <u>NOT</u> Gaussian prime. Then (ii) says $p = (a + bi)(a - bi) = a^2 + b^2 \equiv 0 \pmod{4}$ or $1 \pmod{4}$ or $2 \pmod{4}$.

$\Rightarrow p \not\equiv 3 \pmod{4}$.

($\Rightarrow$) Contrapositive. Suppose $p \equiv 1 \pmod{4}$. Then by [Lagrange's Lemma], $p \mid m^2 + 1$ for some integer m. $\Rightarrow p \mid (m + i)(m - i)$. But $p \nmid m + i$, otherwise we have $p \cdot \gamma = m + i$ for $\gamma \in \mathbb{Z}[i] \Rightarrow \gamma = \frac{m}{p} + \frac{1}{p} i \notin \mathbb{Z}[i]$. Similarly $p \nmid m - i$.

$\therefore p$ is **NOT** a prime in $\mathbb{Z}[i]$. $\qquad \square$

Corollary 3.75. *The Gaussian primes in $\mathbb{Z}[i]$ must be of the form*
(i) $p \in \mathbb{Z}$, $p \equiv 3 \pmod{4}$ or   (ii) $a + bi \in \mathbb{Z}[i]$, $a, b \neq 0$, $a^2 + b^2 = p$
(or $p \in \mathbb{Z}$, $p \equiv 3 \pmod{4}$)
(This is classification of all primes in $\mathbb{Z}[i]$.)

Corollary 3.76. *(Fermat's Theorem of 2 squares).*
If $p = 4n + 1$ is a prime , then $\exists\, a, b \in \mathbb{Z}$ s.t.: $p = a^2 + b^2$
(e.g.: $5 = 1^2 + 2^2$, $13 = 2^2 + 3^2$, $17 = 1^2 + 4^2$, $1013 =?^2 +?^2$)

Proof: By above, p is **NOT** a Gaussian prime, so $p = (a + bi)(a - bi)$ by (ii). $\therefore p = a^2 + b^2$. $\qquad \square$

One other application: (HW 10)

The integer solutions (x, y, z) of the equation $x^2 + y^2 = z^2$ are all of the form $x = (m^2 - n^2)k$, $y = (2mn)k$, $z = (m^2 + n^2)k$ for integers m, n, k.

Theorem 3.77. *If R is ED , then R is PID .*

Proof: Let N be the "norm function" of R. For any $I \triangleleft R$, take any $a \in I \setminus \{0\}$ such that $N(a) > 0$ is minimal among all $r \in R$. Now for any $i \in R$, one has $a|i$ or $i = aq + r$ with $N(r) < N(a)$. But $r = i - aq \in I$ with smaller norm than a. Contradict.

$\therefore a|i \Rightarrow i = aq \in \langle a \rangle$ Hence , $I = \langle a \rangle$. $\qquad\qquad\square$

3.10   Polynomial

In this section, we'll study ring of polynomials $R[x]$. A main reason why it is important is the following:

Theorem 3.78. *Let $\mathbb{F}$ be a field. Suppose $f(x) \in \mathbb{F}[x]$ is irreducible, then $\mathbb{F}[x]/\langle f(x) \rangle$ is a field.*

Proof: Since $\mathbb{F}[x]$ is ED, then PID. $I = \langle f(x) \rangle$ is prime, since $f(x)$ is irreducible $\Leftrightarrow$ prime. Therefore, $I = \langle f(x) \rangle$ is a maximal ideal, and the result follows from Section on max/prime ideals. As a consequence, one has an injective homomorphism of fields $\phi : \mathbb{F} \to \mathbb{F}[x]/\langle f(x) \rangle =: \mathbb{K}$ with $\phi(a) := a + \langle f(x) \rangle$, i.e.: we can "extend" $\mathbb{F}$ to a large field.

> **Galois theory**: Understand roots of (irreducible) polynomials $f(x) \in \mathbb{F}[x]$
>
> As before, we write $\overline{x} := x + \langle f(x) \rangle \in \mathbb{K}$.
>
> Then in $\mathbb{K}$, $f(\overline{x}) = f(x) + \langle f(x) \rangle = 0_{\mathbb{K}}$.
>
> So $\overline{x}$ is a root of $f(x)$ in $\mathbb{K}$!

Now we know the importance of studying when a polynomial is irreducible. In this section, we'll offer ways to check whether $p \in \mathbb{Q}[x]$ (or $\mathbb{Z}[x]$) is irreducible.

Although $\mathbb{Z}$ is **NOT** a field, it is useful in understanding whether $p(x) \in \mathbb{Q}[x]$ with $\mathbb{Z}$-coefficients in irreducible or not.

Lemma 3.79. *Let $\phi : R \to S$ be unital ring homomorphism (e.g.: $\mathbb{Z} \hookrightarrow \mathbb{Q}$, $\mathbb{Z} \twoheadrightarrow \mathbb{Z}_p$). Then $\tilde{\phi} : R[x] \to S[x]$ with $\tilde{\phi}(a_0 + a_1 x + \cdots + a_n x^n) := \phi(a_0) + \phi(a_1)x + \cdots + \phi(a_n)x^n$ is a unital ring homomorphism.*

Theorem 3.80. *(Reduction Test) Let $f \in \mathbb{Z}[x]$ monic, p prime such that $\tilde{\phi}(f) \in \mathbb{Z}_p[x]$ is irreducible. Then f is irreducible.*

Proof: Let $f = gh$ (WTS: g or $h \in \mathbb{Z}[x]$ are constant)

$\Rightarrow \tilde{f} = \tilde{g}\tilde{h}$ in $\mathbb{Z}_p[x]$

Since f is monic, g and h are monic (up to units ± 1) and hence $\tilde{f}, \tilde{g}, \tilde{h}$ monic with $\deg(\tilde{g}) = \deg(g)$, $\deg(\tilde{h}) = \deg(h)$.

But since $\tilde{f}$ is irreducible, then $\deg(\tilde{g})$ or $\deg(\tilde{h}) = 0$.

$\Rightarrow \deg(g)$ or $\deg(h) = 0$. $\qquad\qquad\square$

Example 3.81. $f(x) = x^3 + 3x + 7 \xrightarrow{p=2} \tilde{f} = \tilde{\phi}(f) = x^3 + x + 1$

$\tilde{f}$ cannot be factorized into $\tilde{g} \cdot \tilde{h}$, since if so either $\deg(\tilde{g})$ or $\deg(\tilde{h}) = 1$, i.e.: $\tilde{g}$ or $\tilde{h} = x$ or $x + 1$.

By factor theorem, this implies $\tilde{f}(0)$ or $\tilde{f}(1) = 0$ in $\mathbb{Z}_2$.

Theorem 3.82. *(Eisenstein's criterion) Suppose* $f(x) = a_n x^n + \cdots + a_1 x + a_0 \in \mathbb{Z}[x]$ *is such that* $\gcd(a_n, \ldots, a_1) = 1$ *(f is primitive) and* p *prime number such that:*
(1) $p \mid a_i \quad \forall 0 \leq i < n$ *; (2)* $p \nmid a_n$ *; (3)* $p^2 \nmid a_0$.
Then $f(x)$ *is irreducible in* $\mathbb{Z}[x]$.

(e.g.: $f(x) = x^4 + 4x + 6 \quad (p = 2) \quad$ is irreducible)

Proof: Use $\sim: \mathbb{Z}[x] \to \mathbb{Z}_p[x]$. Then consider $f = gh \in \mathbb{Z}[x]$
(WTS: g or h is constant) .
$\tilde{f} = [a_n]x^n \in \mathbb{Z}_p[x]$ by (1) where $[a_n] \neq [0]$ in $\mathbb{Z}_p$ by (2)
$\Rightarrow \tilde{g} \cdot \tilde{h} = \tilde{f} \sim x^n$. Then $\tilde{g} \sim x^i$; $\tilde{h} \sim x^{n-i} \quad (0 \leq i \leq n)$
Suppose $i \neq 0, n$. Then $\tilde{g} \sim x^i \Rightarrow$ constant term of g is a multiple of p . Similarly , $\tilde{h} \sim x^{n-i} \Rightarrow$ constant term of h is also multiple of p .
$\Rightarrow$ constant term of $f = gh$ is a multiple of p^2 , which contradict (3) of Eisenstein's criterion .
$\therefore \tilde{g}$ or $\tilde{h}$ is a constant function in $\mathbb{Z}_p[x] \quad$ —— (*)
Now:
- $\deg(g) + \deg(h) = \deg(f) = \deg(\tilde{f}) = \deg(\tilde{g}) + \deg(\tilde{h})$.
- $\deg(\tilde{g}) \leq \deg(g)$; $\deg(\tilde{h}) \leq \deg(h)$
$\therefore \deg(\tilde{g}) = \deg(g)$; $\deg(\tilde{h}) = \deg(h)$
$\therefore \deg(g)$ or $\deg(h) = 0$ by (*) , so either g or h is constant polynomial. $\qquad \square$

Example 3.83. $g(x) = x^4 + 1$. Consider $f(x) = g(x+1) = (x+1)^4 + 1 = x^4 + 4x^3 + 6x^2 + 4x + 1 + 1 = x^4 + 4x^3 + 6x^2 + 4x + 2$. Since $2 \mid 4, 6, 4, 2$ and $2^2 \nmid 2$, then $f(x)$ (and $g(x)$) is irreducible.

Q: How about irreducibility in $\mathbb{Q}[x]$**?**

Theorem 3.84. *(Gauss' Lemma) A nonconstant polynomial* $f \in \mathbb{Z}[x]$ *is irreducible* $\Leftrightarrow f \in \mathbb{Q}[x]$ *is irreducible and* f *is primitive.*

Proof: ($\Leftarrow$) Suppose $f \in \mathbb{Z}[x]$ and $f = g \cdot h$, $g, h \in \mathbb{Z}[x]$.
(WTS: g or h is a unit in $\mathbb{Z}[x]$)
By hypothesis, f is irreducible in $\mathbb{Q}[x]$ and $g, h \in \mathbb{Z}[x] \subseteq \mathbb{Q}[x]$, therefore, $\deg(g) = 0$ or $\deg(h) = 0$.
$\Rightarrow g$ or h must be a constant integer.
By hypothesis again, since f is primitive, this constant integer can only be 1 or -1. $\Rightarrow g$ or h is a unit in $\mathbb{Z}[x]$.
($\Rightarrow$) If f is irreducible in $\mathbb{Z}[x]$, then obviously f is primitive. Otherwise $f = a_n x^n + \cdots + a_0$ and $2 \leq d = \gcd(a_n, \ldots, a_0)$, then $f = d \cdot [\frac{a_n}{d} x^n + \cdots + \frac{a_0}{d}]$ where d and $[\frac{a_n}{d} x^n + \cdots + \frac{a_0}{d}]$ are both NOT units in $\mathbb{Z}[x]$ contradicting f irreducible in $\mathbb{Z}[x]$.
To see why f is irreducible in $\mathbb{Q}[x]$, suppose $f = g \cdot h$, $g, h \in \mathbb{Q}[x]$.
(WTS: g, h are units in $\mathbb{Q}[x]$, i.e.: $g, h \in \mathbb{Q} \setminus \{0\}$)
Let $\lambda, \mu \in \mathbb{N}$ be the smallest positive integers such that $\lambda \cdot g$ and $\mu \cdot h \in \mathbb{Z}[x]$. Let $g' = \lambda \cdot g$, $h' = \mu \cdot h \in \mathbb{Z}[x]$.

<u>Claim:</u> $\lambda = 1$: Suppose on contrary, $\exists p \mid \lambda$. Then "$p \mid \lambda f$" (which means the coefficients of $\lambda f \in \mathbb{Z}[x]$ are multiples of p).

$\Rightarrow p \mid (\lambda \alpha^{-1} g) \cdot (\alpha h) \Rightarrow$ "$p \mid g'h'$" in $\mathbb{Z}[x]$

Therefore, $\tilde{g}' \cdot \tilde{h}' \equiv 0$ in $\mathbb{Z}_p[x] \Rightarrow \tilde{g}' \equiv 0$ or $\tilde{h}' \equiv 0$ in $\mathbb{Z}_p[x]$

$\Rightarrow$ "$p \mid g'$" — (*) or "$p \mid h'$" — (**) in $\mathbb{Z}[x]$

In (*), "$p \mid g'$" $\Leftrightarrow$ "$p \mid \lambda \alpha^{-1} g$" $\Rightarrow (\frac{\lambda}{p}) \alpha^{-1} g$, $\alpha h \in \mathbb{Z}[x]$, contradicting the minimality of λ.

In (**), "$p \mid \alpha h$" $\Rightarrow (\frac{\lambda}{p})(\frac{\alpha}{p})^{-1} g = \lambda \alpha^{-1} g \in \mathbb{Z}[x]$ and $(\frac{\alpha}{p}) h \in \mathbb{Z}[x]$, contradicting the minimality of λ.

$\therefore p \nmid \lambda$ for any prime number, and $\lambda = 1$.

Therefore, we have $\alpha \in \mathbb{Q} \setminus \{0\}$ s.t.: $f = gh$ in $\mathbb{Q}[x]$ and $f = (\alpha^{-1} g)(\alpha h)$ in $\mathbb{Z}[x]$, since we have f irreducible in $\mathbb{Z}[x]$ by hypothesis, $\alpha^{-1} g$ or αh is a unit in $\mathbb{Z}[x] \Rightarrow \alpha^{-1} g = \pm 1$ or $\alpha h = \pm 1$.

$\Rightarrow g = \alpha'$ where $\alpha' \in \mathbb{Q} \setminus \{0\}$ or $h = \alpha''$ where $\alpha'' \in \mathbb{Q} \setminus \{0\}$. $\square$

Theorem 3.85. $\mathbb{Z}[x]$ *is a UFD (but not PID by HW).*

Proof: Let $f \in \mathbb{Z}[x]$ be primitive.

Claim: f can be factorized into $f = g_1 \ldots g_n$, $g_i \in \mathbb{Z}[x]$ primitive irreducible. To see why, use induction on $\deg(f)$. True for $\deg(f) = 0$ or 1. So assume claim holds for $\deg(f) < k$ Then for f with $\deg(f) = k$ and primitive:

• f is irreducible ; • $f = h_1 h_2$, $h_1, h_2 \in \mathbb{Z}[x]$ NOT units.

If $\deg(h_1)$ or $\deg(h_2) = 0$, then h_1 or h_2 equal to constant $\neq \pm 1$ contradicting f is primitive. Hence, $\deg(h_1) \& \deg(h_2) > 0$ by induction. To see why $\mathbb{Z}[x]$ is UFD, note that we can factorize any primitive $f \in \mathbb{Z}[x]$ into irreducibles (primitive).

To see the factorization is unique, suppose $f = g_1 \ldots g_n = h_1 \ldots h_m$, $g_i, h_j \in \mathbb{Z}[x]$ primitive irreducible. Then since $\mathbb{Z}[x] \subseteq (\mathbb{Q}[x]$ UFD) and by Gauss' Lemma $g_i, h_j \in \mathbb{Q}[x]$ irreducible by unique factorization of $\mathbb{Q}[x]$, we have $n = m$, and $\exists \sigma \in S_n$, s.t.: $g_i \sim h_{\sigma(i)}$ (in $\mathbb{Q}[x]) \Rightarrow g_i = h_{\sigma(i)} \cdot u$ where $u \in$ (unit in $\mathbb{Q}[x])$ Since $g_i, h_{\sigma(i)}$ primitive, $u \in \{\pm 1\} \Rightarrow g_i \sim h_{\sigma(i)}$ in $\mathbb{Z}[x]$. $\square$

Remark (more generally):

If R is a UFD, then $R[x]$ is UFD. To prove it, use Field of Fractions (HW ?) $F = \text{Frac}(R)$ and the fact that $F[x]$ is a PID ($\Rightarrow$ UFD).

Chapter 4

Fields

Recall: F is a field if F is an integral domain (unital) and for all $x \in F \setminus \{0\}$, there exists a multiplicative inverse $x^{-1} \in F \setminus \{0\}$.

In general, there are 3 kinds of fields:

1. **Number field** (a subfield of $\mathbb{C}$)

   (e.g., $\mathbb{Q}$, $\mathbb{R}$, $\mathbb{Q}(i)$, ...)

2. **Finite field** (a field F with $|F| < \infty$)

   (e.g., $\mathbb{Z}_p = \mathbb{F}_p$ for p prime. $\mathbb{F}_2[x]/\langle x^3 + x + 1 \rangle$ has $8 = 2^3$ elements.)

3. **Function fields**

   (e.g., $\mathbb{C}(x) = \mathrm{Frac}(\mathbb{C}[x]) = \{f(x)/g(x) \mid f(x), g(x) \in \mathbb{C}[x], g(x) \neq 0\}$)

We'll only focus on (1) or (2).

4.1   Field Extension and Degree

Motivation: Given a field F, construct a "bigger" field K that "contains" F.

For example, let $F = \mathbb{F}_2$ and $K = \mathbb{F}_2[x]/\langle x^3 + x + 1 \rangle$. Then we have an injective unital ring (or field) homomorphism $i : F \to K$ where $i(0) = 0 + \langle x^3 + x + 1 \rangle$ and $i(1) = 1 + \langle x^3 + x + 1 \rangle$. So we say K "contains" F.

Definition 4.1. Let F and K be fields. We say F is a **subfield** of K or K is a **field extension** of F if there is an injective unital field homomorphism $i : F \to K$.

In this case, K is a vector space over F. There is a scalar multiplication map $\cdot : F \times K \to K$ defined by $\alpha \cdot x = i(\alpha)x$ for $\alpha \in F, x \in K$, that satisfies:

- $0 \cdot x = 0$

- $1 \cdot x = x$

- $\alpha \cdot (x + y) = (\alpha \cdot x) + (\alpha \cdot y)$

- $\ldots$

The **degree** of extension is defined as:

$$[K : F] := \dim_F(K) \quad \text{(as a vector space over } F)$$

Example 4.2. 1. $F = \mathbb{R}$, $K = \mathbb{C}$, with the inclusion $i : \mathbb{R} \hookrightarrow \mathbb{C}$.

Then $[\mathbb{C} : \mathbb{R}] = 2$, since $\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}\} = \text{span}_{\mathbb{R}}\{1, i\}$.

Thus, $\dim_{\mathbb{R}}(\mathbb{C}) = 2$ (since $\{1, i\}$ is linearly independent over $\mathbb{R}$).

2. $F = \mathbb{Q}$, $K = \mathbb{R}$, with the inclusion $i : \mathbb{Q} \hookrightarrow \mathbb{R}$.

To find $[\mathbb{R} : \mathbb{Q}]$, consider that $\mathbb{R} = \text{span}_{\mathbb{Q}}\{1, \sqrt{2}, \sqrt{3}, e, \pi, \dots\}$.

The set of transcendental numbers like e, π is infinite and they are linearly independent over $\mathbb{Q}$. Thus, $[\mathbb{R} : \mathbb{Q}] = \infty$.

4.2 Splitting Extension

In field theory, we want to understand roots of polynomials $p(x) \in F[x]$. More precisely, we would like to construct $E : F$ such that E contains some (or all) roots of $F[x]$. The first step is the following:

Theorem 4.3 (Kronecker)**.** *Let F be a field, and $p(x) \in F[x]$ be an irreducible polynomial of degree m. Then $K = F[x]/\langle p(x)\rangle$ is a field extension of F such that $[K : F] = m$. Moreover, there exists an element $\alpha \in K$ such that:*

- $K = F[\alpha] := \{a_n\alpha^n + \cdots + a_1\alpha + a_0 \mid a_i \in F, n \in \mathbb{N}\}$.

- $p(\alpha) = 0$ *in K.*

Proof. Since $p(x) = b_m x^m + \cdots + b_1 x + b_0 \in F[x]$ is irreducible, the ideal $\langle p(x)\rangle$ is a maximal ideal in $F[x]$. This implies that $K = F[x]/\langle p(x)\rangle$ is a field.

Let's study the dimension, $\dim_F(K)$. Consider the element $\alpha := x + \langle p(x)\rangle \in K$. Then in K,

$$
\begin{aligned}
p(\alpha) &= b_m\alpha^m + \cdots + b_1\alpha + b_0 \\
&= b_m(x + \langle p(x)\rangle)^m + \cdots + b_1(x + \langle p(x)\rangle) + b_0(1 + \langle p(x)\rangle) \\
&= (b_m x^m + \cdots + b_1 x + b_0) + \langle p(x)\rangle \\
&= p(x) + \langle p(x)\rangle = 0 + \langle p(x)\rangle = 0_K
\end{aligned}
$$

So, the second part of the theorem is proved. Moreover, this also implies that

$$\alpha^m = -\frac{1}{b_m}(b_{m-1}\alpha^{m-1} + \cdots + b_0)$$

i.e. the set $\{1, \alpha, \dots, \alpha^m\}$ is linearly dependent in K. More precisely, α^m is a linear combination of $\{1, \alpha, \dots, \alpha^{m-1}\}$. Similarly, for any $n \geq m$, α^n is also a linear combination of $\{1, \alpha, \dots, \alpha^{m-1}\}$.

Claim: The set $\{1, \alpha, \dots, \alpha^{m-1}\}$ is a basis of K over F.

Spanning set: Take any element $k \in K$. By definition, $k = (c_l x^l + \cdots + c_1 x + c_0) + \langle p(x) \rangle$ for some $c_i \in F$. This is equal to $c_l \alpha^l + \cdots + c_1 \alpha + c_0$. Since all powers α^j for $j \geq m$ can be reduced to a linear combination of $\{1, \alpha, \ldots, \alpha^{m-1}\}$, any element $k \in K$ can be written as a linear combination of these basis elements. Thus, $K = \mathrm{Span}_F \{1, \alpha, \ldots, \alpha^{m-1}\}$.

Linearly independent: Suppose $d_{m-1} \alpha^{m-1} + \cdots + d_1 \alpha + d_0 = 0_K$ for some $d_i \in F$, not all zero. This is equivalent to $(d_{m-1} x^{m-1} + \cdots + d_1 x + d_0) + \langle p(x) \rangle = 0_K$. Let $g(x) = d_{m-1} x^{m-1} + \cdots + d_0$. The equation means $g(x) \in \langle p(x) \rangle$. This implies that $p(x)$ divides $g(x)$, i.e., $g(x) = p(x) \cdot r(x)$ for some $r(x) \in F[x]$. But this is a contradiction, because $\deg(g(x)) \leq m - 1$ while $\deg(p(x)) = m$. The degree of a non-zero polynomial $g(x)$ cannot be less than the degree of a non-zero polynomial $p(x)$ that divides it. Thus, $g(x)$ must be the zero polynomial, which means all d_i must be zero. The set is linearly independent.

Since $\{1, \alpha, \ldots, \alpha^{m-1}\}$ is a basis for K over F, the dimension is m. Therefore, $[K : F] = m$. $\square$

Example 4.4. Let $F = \mathbb{Q}$, $p(x) = x^3 - 2 \in \mathbb{Q}[x]$ (irreducible by Eisenstein's criterion). Then $[K : \mathbb{Q}] = 3$ for $K = \mathbb{Q}[x]/\langle x^3 - 2 \rangle$. Let $\alpha = x + \langle x^3 - 2 \rangle$. Then $K = \mathrm{Span}_{\mathbb{Q}}\{1, \alpha, \alpha^2\} = \{a_2 \alpha^2 + a_1 \alpha + a_0 \mid a_i \in \mathbb{Q}\}$. We know $\alpha^3 - 2 = 0$, so $\alpha^3 = 2$. K is a field that contains $\mathbb{Q}$ and a cube root of 2. We also know how to do arithmetic in K, for instance:

$$
\begin{aligned}
(2\alpha^2 + \tfrac{3}{4}) \cdot (\alpha - 6) &= 2\alpha^3 - 12\alpha^2 + \tfrac{3}{4}\alpha - \tfrac{18}{4} \\
&= 2(2) - 12\alpha^2 + \tfrac{3}{4}\alpha - \tfrac{9}{2} \\
&= 4 - 12\alpha^2 + \tfrac{3}{4}\alpha - \tfrac{9}{2} \\
&= -12\alpha^2 + \tfrac{3}{4}\alpha - \tfrac{1}{2}
\end{aligned}
$$

Exercise: What is $\alpha^{-1} \in K$?

In the above example, under the extension $K : \mathbb{Q}$, one can treat the polynomial $p(x) = x^3 - 2 \in \mathbb{Q}[x] \subseteq K[x]$ under the extended field. By Kronecker's theorem, $p(x) \in K[x]$ has a root:

$$
p(x) = (x - \alpha)(x^2 + \alpha x + \alpha^2) \in K[x]
$$

However, K does not contain all the roots of $p(x)$.

Definition 4.5. Let F be a field, and $p(x) \in F[x]$. We say a field extension $E : F$ splits $p(x)$ if $p(x) = a(x - a_1) \cdots (x - a_n)$.

Proposition 4.6. *For any field F and any polynomial $p(x) \in F[x]$, there exists a field extension $E : F$ such that E splits $p(x)$.*

Proof. Prove by induction on the degree of $p(x)$ (for any base field F). If $p(x) \in F$ is of degree 1, we are done. By induction hypothesis, assume the theorem holds for all polynomials of degree $< k$ over any field. Consider a polynomial $p(x)$ of degree k. Let $f(x)$ be an irreducible factor of $p(x)$. Then take $K = F[x]/\langle f(x) \rangle$ so that $a := x + \langle f(x) \rangle \in K$ is a root of $f(x)$ and thus of $p(x)$. So

$p(x) = (x - a)g(x)$ in $K[x]$, where $\deg(g(x)) = k - 1$. By the induction hypothesis, there exists an extension $E : K$ such that $g(x)$ splits in $E[x]$, say $g(x) = c(x - a_2)...(x - a_k)$. So we have $p(x) = c(x - a)(x - a_2)...(x - a_k)$ in $E[x]$, and we are done.   □

Now we know for any $p(x) \in F[x]$, there is a field extension $E : F$ such that E contains all the roots of $p(x)$. We want to find 'the smallest' extension that contains all the roots of $p(x)$.

Definition 4.7. Let $E : F$ be a field extension, and $e_1, \ldots, e_n \in E$. Write $F(e_1, \ldots, e_n)$ for the smallest field extension of F containing $e_1, \ldots, e_n$. In other words,

$$F(e_1, \ldots, e_n) := \bigcap_{K \leq E, \; e_1, \ldots, e_n \in K} K$$

Proposition 4.8. $F(e_1, e_2, \ldots, e_n) = (F(e_1))(e_2, \ldots, e_n) = \cdots = ((F(e_1)(e_2)) \ldots (e_{n-1}))(e_n)$.

Proof. Easy.   □

Definition 4.9. Suppose $E : F$ is a field extension such that E splits a polynomial $p(x) \in F[x] \subseteq E[x]$ with roots $a_1, \ldots, a_n \in E$. Then a **splitting field** of $p(x)$ over F is the smallest subfield $F(a_1, \ldots, a_n)$ of E containing all the roots a_i of $p(x)$ in E.

Example 4.10.

1. Let $p(x) = x^2 + 3x + 3 \in \mathbb{Q}[x]$. Then for $E = \mathbb{Q}[x]/\langle p(x) \rangle$, the polynomial $p(x)$ splits automatically into $(x - a)(x - b)$. So $\mathbb{Q}(a, b) = \mathbb{Q}[x]/\langle p(x) \rangle$ is a splitting field.

   On the other hand, $p(x)$ splits in $\mathbb{C}$ and the roots are $-3/2 \pm i\sqrt{3}/2$. So $\mathbb{Q}(i\sqrt{3})$ is also a splitting field of $p(x)$.

2. Let $p(x) = x^3 - 2 \in \mathbb{Q}[x]$. Then $F = \mathbb{Q}[x]/\langle x^3 - 2 \rangle$ has a root $\alpha \in E$. We can form $E = F[x]/\langle x^2 + \alpha x + \alpha^2 \rangle$. The field E is a splitting field, which we can write as $\mathbb{Q}(a, b, c)$ where a, b, c are the roots.

   On the other hand, working in $\mathbb{C}$, the roots are $2^{1/3}, 2^{1/3}\omega, 2^{1/3}\omega^2$ (where $\omega = e^{i2\pi/3}$). Then $\mathbb{Q}(2^{1/3}, 2^{1/3}\omega, 2^{1/3}\omega^2) = \mathbb{Q}(2^{1/3}, \omega)$ is also a splitting field of $p(x)$.

Theorem 4.11. *(will prove in MAT5210) All splitting fields of the same polynomial are isomorphic.*

4.3   Simple Extension

In the previous section, we begin with a irreducible polynomial $p(x) \in F[x]$, and extend to E so that E contains some (or all) roots of $p(x)$. In this section, we change our perspective a little bit – we begin with a field extension $E : F$, and see which elements in E are roots of a polynomial $p(x) \in F[x]$. Such $E : F$ always exists by the virtue of the previous section.

Definition 4.12. Let F be a field and L be a field extension of F (e.g., $F = \mathbb{Q} \subseteq L = \mathbb{C}$). An element $\alpha \in L$ is

- **algebraic** if there exists a non-zero polynomial $p(x) \in F[x]$ such that $p(\alpha) = 0$ (in L).

- Otherwise, α is called **transcendental**.

Example 4.13 (Elements over $\mathbb{Q}$)**.**

- $i = \sqrt{-1}$ is algebraic over $\mathbb{Q}$ (its polynomial is $p(x) = x^2 + 1$).

- $\sqrt[4]{2}, \sqrt{3}$ are algebraic over $\mathbb{Q}$ (their polynomials are $p(x) = x^4 - 2$ and $x^2 - 3$, respectively).

- $\sqrt{2} + \sqrt{3}$ is algebraic over $\mathbb{Q}$ (HW12).

- e, π are transcendental over $\mathbb{Q}$ (there is NO polynomial with $\mathbb{Q}$-coefficients having e or π as a root).

Theorem 4.14. *If α is transcendental over F, then $F[\alpha] \cong F[x]$ as rings (but not as fields).*

As for β algebraic:

Proposition 4.15. *Let $\beta \in L$ be algebraic over F. Consider the set $I = \{f(x) \in F[x] \mid f(\beta) = 0\}$. Then:*

1. *I is a non-zero ideal in $F[x]$.*

2. *$I = \langle p(x) \rangle$ for some unique monic polynomial $p(x) \in F[x]$. (This is because $F[x]$ is a PID).*

3. *$p(x)$ is the unique monic polynomial of smallest degree in I.*

4. *$p(x)$ is irreducible.*

*This polynomial $p(x)$ is called the **minimal polynomial** of β over F.*

Proof. (a) Let's show $p(x)$ has the smallest degree. By the division algorithm, suppose on the contrary there exists $g(x) \in I$ with $\deg(g) < \deg(p)$. Since $I = \langle p(x) \rangle$, $g(x)$ must be a multiple of $p(x)$, which is impossible unless $g(x) = 0$. Contradiction.

(b) Let's show $p(x)$ is irreducible. Suppose on the contrary that $p(x) = h_1(x)h_2(x)$ with $\deg(h_1) > 0$ and $\deg(h_2) > 0$. Then $p(\beta) = h_1(\beta)h_2(\beta) = 0$ in the field L. This implies that either $h_1(\beta) = 0$ or $h_2(\beta) = 0$. This means either $h_1(x) \in I$ or $h_2(x) \in I$. But $\deg(h_1) < \deg(p)$ and $\deg(h_2) < \deg(p)$, which contradicts the fact that $p(x)$ is a polynomial of smallest degree in I. Thus, $p(x)$ must be irreducible. $\square$

Definition 4.16. Let F be a field, $\beta \in L$ (a field extension of F). The polynomial $m_\beta(x) := p(x)$ appearing in Proposition 4.15 above is called the **minimal polynomial** of β over F.

Example 4.17. Minimal polynomials over $\mathbb{Q}$:

- For $\beta = i$, the minimal polynomial is $m_\beta(x) = x^2 + 1$.

- For $\beta = \sqrt[4]{2}$, the minimal polynomial is $m_\beta(x) = x^4 - 2$.

- For $\beta = e^{2\pi i/5}$, the minimal polynomial is $m_\beta(x) = x^4 + x^3 + x^2 + x + 1$.
  (Since $\beta^5 = 1$, and $x^5 - 1 = (x - 1)p(x)$).

Theorem 4.18. *Let $L : F$ be a field extension, and $\beta \in L$ be algebraic. Then the smallest field extension $F(\beta) : F$ containing β is equal to the polynomial ring $F[\beta]$, which is isomorphic to $F[x]/\langle m_\beta(x)\rangle$.*

Proof. Consider the ring homomorphism $\phi : F[x] \to F[\beta]$ defined by evaluation at β:

$$\phi(a_n x^n + \cdots + a_0) = a_n \beta^n + \cdots + a_0$$

The kernel of this map is precisely the ideal $I = \{f(x) \in F[x] \mid f(\beta) = 0\}$, which we know is equal to $\langle m_\beta(x)\rangle$. So the result follows by the First Isomorphism Theorem for rings. Since $\langle m_\beta(x)\rangle$ is a maximal ideal, $F[x]/\langle m_\beta(x)\rangle$ is a field, and therefore $F[\beta]$ is also a field. $\qquad\square$

Definition 4.19. The field extension $F(\beta) := F[\beta]$ of F is called the **simple extension** of F by β.

Corollary 4.20. *If $\alpha, \beta \in L$ are algebraic over F and are roots of the same irreducible polynomial $p(x) \in F[x]$, then $F(\alpha) \cong F(\beta)$.*

Proof. $m_\alpha(x) = m_\beta(x) = p(x)$. So $F(\alpha) \cong F[x]/\langle p(x)\rangle \cong F(\beta)$. $\qquad\square$

Let $E : F$ be a field extension, and $\alpha_1, \ldots, \alpha_k \in E$ be algebraic over F. Then the field extension $F(\alpha_1, \ldots, \alpha_k)$ can be constructed inductively as follows: Define a sequence of fields by $F_0 = F$ and

$$F_{i+1} := F_i(\alpha_{i+1}) \cong \frac{F_i[x]}{\langle m_{\alpha_{i+1}}^{F_i}(x)\rangle} \quad \text{for } 0 \le i \le k - 1.$$

where $m_{\alpha_{i+1}}^{F_i}(x) \in F_i[x]$ is the minimal polynomial of the element α_{i+1} over the field F_i. Then $F_k = F(\alpha_1, \ldots, \alpha_k)$.

4.4   Algebraic Extension

We have seen that a simple extension $F(\beta) : F$ contains an algebraic element β. What about the other elements in $F(\beta) = \mathrm{Span}_F\{1, \beta, \ldots, \beta^{\deg(m_\beta(x))-1}\}$? Are they algebraic? More generally, how about the non-simple extension $F(\beta_1, \ldots, \beta_k)$?

Definition 4.21. An extension $E : F$ is called **algebraic** if every element $\alpha \in E$ is algebraic over F.

Theorem 4.22. *If $[E : F] < \infty$ (i.e., it is a finite extension), then E is an algebraic extension of F.*

Proof. Suppose $[E : F] = n$. Then for any $\alpha \in E$, consider the set $\{1, \alpha, \alpha^2, \ldots, \alpha^n\}$. This is a set of $n + 1$ elements in an n-dimensional vector space, so it must be linearly dependent over F. This implies there exist scalars $c_0, \ldots, c_n \in F$, not all zero, such that $c_0 \cdot 1 + c_1 \alpha + \cdots + c_n \alpha^n = 0$. Thus, α is a root of the non-zero polynomial $p(x) = c_n x^n + \cdots + c_0 \in F[x]$, which means α is algebraic over F. Since α was arbitrary, the extension is algebraic. $\qquad\square$

Corollary 4.23. *Let $E : F$ be a field extension, and $\beta_1, \ldots, \beta_k \in E$ are algebraic over F. Then the extension $F(\beta_1, \ldots, \beta_k) : F$ is algebraic.*

Proof. For the special case when $k = 1$, one has $[F(\beta) : F] = \deg(m_\beta(x))$ which is finite. The proof of the general case is given in Corollary 4.27 below. $\qquad\square$

Theorem 4.24 (Primitive Element Theorem). *Let F be a field with characteristic $\mathrm{char}(F) = 0$. Let $a, b \in K$ be algebraic over F in some extension $K : F$. Then there exists an element $c \in F(a, b)$ such that $F(c) = F(a, b)$. (Such an element c is called a primitive element for the extension).*

Proof. Let $p(x)$ be the minimal polynomial of a over F. Let $q(x)$ be the minimal polynomial of b over F.

Take a field extension $L : K$ such that $p(x)$ and $q(x)$ split completely in $L[x]$. Suppose $a_1, \ldots, a_m \in L$ are the roots of $p(x)$, with $a_1 = a$. Suppose $b_1, \ldots, b_n \in L$ are the roots of $q(x)$, with $b_1 = b$.

Since $\mathrm{char}(F) = 0$, the field F is infinite. This allows us to choose an element $d \in F$ such that

$$d \neq \frac{a_i - a_j}{b_k - b_l}$$

for all i, j and for all $k \neq l$. In particular, we choose $d \in F$ such that for any i and any $j > 1$:

$$d \neq \frac{a_i - a}{b - b_j}$$

(Note that the elements on the right-hand side may not be in F, but there are only a finite number of such values, so we can always find a $d \in F$ that avoids them).

Consider the element $c := a + db$. Then, it is clear that $F(c) \subseteq F(a, b)$ since $a, b \in F(a, b)$ and $d \in F \subseteq F(a, b)$.

To show the other inclusion, $F(a, b) \subseteq F(c)$, it is enough to show that $b \in F(c)$, because if $b \in F(c)$, then $a = c - db \in F(c)$ as well.

Consider the polynomials $r(x) = p(c - dx)$ and $q(x)$, both viewed as polynomials in $F(c)[x]$. We evaluate both polynomials at b:

- $r(b) = p(c - db) = p(a) = 0$.

- $q(b) = 0$.

Let $m(x) \in F(c)[x]$ be the minimal polynomial of b over the field $F(c)$. Since b is a root of both $r(x)$ and $q(x)$ (which are in $F(c)[x]$), it must be that $m(x)$ divides both $r(x)$ and $q(x)$ in $F(c)[x]$.

$$m(x) \mid r(x) \quad \text{and} \quad m(x) \mid q(x)$$

Now let's study the polynomial $m(x) \in F(c)[x] \subseteq L[x]$.

Since $m(x) \mid q(x)$ and $q(x)$ splits in $L[x]$, the roots of $m(x)$ must be a subset of the roots of $q(x)$, which are $\{b = b_1, b_2, \ldots, b_n\}$.

Consider any root b_j of $m(x)$. Since $m(x)$ also divides $r(x)$, this b_j must also be a root of $r(x)$. So, for such a b_j, we must have $r(b_j) = p(c - db_j) = 0$. This implies that $c - db_j$ must be one of the roots of $p(x)$, say a_i.

$$c - db_j = a_i$$

Substituting $c = a + db$:

$$a + db - db_j = a_i$$

$$d(b - b_j) = a_i - a$$

If $j \neq 1$, then $b \neq b_j$. We can write:

$$d = \frac{a_i - a}{b - b_j}$$

But this contradicts our initial choice of d, which was chosen specifically to not be equal to any of these values. Therefore, the only possibility is that $j = 1$, which means $b_j = b_1 = b$, and hence

$$m(x) = x - b$$

in $F(c)[x]$, i.e. $b \in F(c)$ and the theorem is proved. $\square$

4.5 Tower Law

Theorem 4.25 (Tower Law)**.** *Let* $F \subseteq K \subseteq L$ *be field extensions. (e.g.,* $\mathbb{Q} \subseteq \mathbb{Q}(\sqrt{2}) \subseteq \mathbb{Q}(\sqrt{2}, \sqrt{3})$*). Then:*

$$[L : F] = [L : K][K : F]$$

Proof. Let $\{e_1, \ldots, e_n\}$ be a basis of K over F. This means $[K : F] = n$. Let $\{f_1, \ldots, f_m\}$ be a basis of L over K. This means $[L : K] = m$.

Claim: The set $\mathcal{B} = \{e_i f_j \mid 1 \leq i \leq n, 1 \leq j \leq m\}$ is a basis of L over F. If this claim is true, then the dimension of L over F is $n \cdot m$, which proves the theorem. We need to show that $\mathcal{B}$ is a spanning set and is linearly independent over F.

Spanning set: Let x be any element in L. Since $\{f_j\}$ is a basis of L over K, we can write x as a linear combination:

$$x = \sum_{j=1}^{m} \mu_j f_j, \quad \text{where each } \mu_j \in K$$

Now, for each coefficient $\mu_j \in K$, we can express it as a linear combination of the basis elements of K over F. For each j, we have:

$$\mu_j = \sum_{i=1}^{n} \nu_{ij} e_i, \quad \text{where each } \nu_{ij} \in F$$

Substituting this back into the expression for x:

$$x = \sum_{j=1}^{m} \left(\sum_{i=1}^{n} \nu_{ij} e_i \right) f_j = \sum_{j=1}^{m} \sum_{i=1}^{n} \nu_{ij} (e_i f_j)$$

Since every element $x \in L$ can be written as a linear combination of the elements $\{e_i f_j\}$ with coefficients $\nu_{ij} \in F$, this set spans L over F.

Linear Independence: Suppose we have a linear combination of the elements $\{e_i f_j\}$ that equals zero, with coefficients β_{ij} from the field F:

$$\sum_{i=1}^{n} \sum_{j=1}^{m} \beta_{ij} e_i f_j = 0, \quad \text{where } \beta_{ij} \in F$$

We can regroup the terms by factoring out the f_j:

$$\sum_{j=1}^{m} \left(\sum_{i=1}^{n} \beta_{ij} e_i \right) f_j = 0$$

For each j, the inner sum $\sum_{i=1}^{n} \beta_{ij} e_i$ is a linear combination of elements of the basis $\{e_i\}$ with coefficients in F. This means the term in the parenthesis is an element of the field K. So, we have a linear combination of the basis elements $\{f_j\}$ with coefficients from K that equals zero. By the linear independence of the set $\{f_1, \ldots, f_m\}$ over K, all the coefficients must be zero:

$$\sum_{i=1}^{n} \beta_{ij} e_i = 0 \quad \text{for all } j = 1, \ldots, m$$

Now, for each j, we have a linear combination of the basis elements $\{e_i\}$ over F that equals zero. By the linear independence of the set $\{e_1, \ldots, e_n\}$ over F, all the coefficients must be zero:

$$\beta_{ij} = 0 \quad \text{for all } i, j$$

This shows that the set $\{e_i f_j\}$ is linearly independent over F, which completes the proof of the Tower Law. $\qquad \square$

Example 4.26. Let $F = \mathbb{F}_2$ (which is the same as $\mathbb{Z}_2$). Let $L = \mathbb{F}_2[x]/\langle x^3 + x + 1 \rangle$. So, $|L| = 2^3 = 8$ and $[L : F] = 3$.

Are there any proper subfields K with $\mathbb{F}_2 \subset K \subset L$? Suppose such a subfield K exists. Then by the Tower Law, we must have:

$$[L : \mathbb{F}_2] = [L : K][K : \mathbb{F}_2]$$

Substituting the known value:

$$3 = [L : K][K : \mathbb{F}_2]$$

Since 3 is a prime number and the degrees of extensions are integers greater than or equal to 1, the only possibilities are:

1. $[L : K] = 1$ and $[K : \mathbb{F}_2] = 3$. This implies $K = L$.

2. $[L : K] = 3$ and $[K : \mathbb{F}_2] = 1$. This implies $K = \mathbb{F}_2$.

Therefore, there are no proper intermediate subfields between L and $\mathbb{F}_2$.

Corollary 4.27. *Suppose we have a tower of fields $F \subseteq E \subseteq K$. If $K : E$ is an algebraic extension and $E : F$ is an algebraic extension, then $K : F$ is also an algebraic extension.*

Proof. Let $a \in K$. Since $K : E$ is algebraic, a is a root of some polynomial $p(x) = b_n x^n + \cdots + b_1 x + b_0 \in E[x]$.

Since $E : F$ is algebraic, each coefficient $b_i \in E$ is algebraic over F. Consider the tower of fields:

$$F \subseteq F_0 := F(b_0) \subseteq F_1 := F_0(b_1) \subseteq \cdots \subseteq F_n := F_{n-1}(b_n) = F(b_0, \ldots, b_n).$$

Since each b_i is algebraic over F, it is also algebraic over F_{i-1}. Therefore, each extension in this tower, $F_i : F_{i-1}$, is a finite extension. By the Tower Law, $[F_n : F]$ is finite.

$$[F_n : F] = [F_n : F_{n-1}] \ldots [F_1 : F_0][F_0 : F] < \infty.$$

Now, the polynomial $p(x)$ has its coefficients in F_n, so $p(x) \in F_n[x]$. Since a is a root of $p(x)$, the extension $F_n(a) : F_n$ is finite, with $[F_n(a) : F_n] \leq \deg(p) = n$.

Consider the degree of the extension $F_n(a)$ over F:

$$[F_n(a) : F] = [F_n(a) : F_n][F_n : F].$$

This product is finite. Since $F(a)$ is a subfield of $F_n(a)$, we have $[F(a) : F] \leq [F_n(a) : F] < \infty$. Because $F(a) : F$ is a finite extension, it must be an algebraic extension. This implies that the element a is algebraic over F. Since a was an arbitrary element of K, the extension $K : F$ is algebraic. $\square$

**Corollary 4.28.** *The set of all algebraic elements in an extension $E : F$ forms a subfield of E.*

Proof. Let $a, b \in E$ be algebraic over F. We need to show that $a + b$, $a - b$, ab, and a/b (for $b \neq 0$) are also algebraic over F. Consider the field $F(a, b)$. By the Tower Law,

$$[F(a, b) : F] = [F(a, b) : F(a)][F(a) : F].$$

Since a is algebraic over F, $[F(a) : F]$ is finite. Since b is algebraic over F, it is also algebraic over the larger field $F(a)$. Thus, $[F(a, b) : F(a)] = [F(a)(b) : F(a)]$ is also finite. Therefore, $[F(a, b) : F]$ is finite. Since $F(a, b)$ is a finite extension of F, it is an algebraic extension. Any element of $F(a, b)$ is algebraic over F. This includes $a + b, a - b, ab$, and a/b. $\square$

As a final application of the Tower Law: What can we do with a ruler (with no markings) and a compass?

- **Bisect an angle**

Q: Can we **tri-sect** any angle??

- **Double a square**

Q: Can you **double a cube**?