

Grain Boundary Diffusion

Adriaan Riet

September 23, 2015

1 Background

Grain boundary diffusion has been a subject of research since at least the early 20th century [1]. In 1951, Fisher published a method of gathering grain boundary diffusion information, with an approximate derivation of the applicable equations [2]. It was observed that diffusion along a grain boundary was much faster than diffusion in bulk phases. Unfortunately, Fisher’s model incorporated several assumptions to make the derivation more straightforward, and the work has been built upon by Mishin and Yurovitskii [3], as well as Klinger and Rabkin [4], focusing on anisotropy of the grain boundary diffusion and inhomogeneity of the bulk phase, respectively. More work has also been done by Turnbull et al. dealing with orientation effects [5]. It has been noted that Fisher’s derivation was very approximate, and that a better theoretical derivation was introduced by Whipple [6], but that Whipple’s derivation was unable to be applied to experiment [7].

An attempt has been made to relate overall average grain boundary diffusion effects [8], which has been used in obtaining intergranular diffusion characteristics [9]. Further, the problem of the “Kirkendall Effect” during grain boundary diffusion shows that the physical location of the interface can change, which has been addressed [10]. Much of the research in grain boundary diffusion has been using experimental observation and phenomenological models, but atomistic models have the potential to give insight into the modes of diffusion and the underlying causes [11]. Thanks to work done by Green [12, 13] and Kubo [14, 15], there is a solid framework by which an atomistic model can be tied to transport properties.

Unfortunately, numerically modeling the systems involved in grain boundary diffusion is computationally intensive. Also, the ensemble used to in many MD simulations involves a constant energy assumption, which is less helpful than a constant temperature assumption. This problem was overcome by the Nose-Hoover method [16]. MD simulations for diffusion problems run with timescales on the order of fs, while the diffusion characteristics are determined on the order of ns, so that meaningful simulations need to run on the order of 10^6 timesteps [17]. An effort was made by Sørensen et al. to use a hybrid kinetic Monte Carlo-molecular dynamics approach, which used molecular dynamics to determine the modes by which diffusion can occur, and using kinetic monte carlo to determine the rates of diffusion. This allowed for a smaller, less rigorous molecular dynamics simulation, and because the monte carlo method does not deal with these disparate timescales, computational efficiency was improved [18].

Periclase is one of the most important minerals for understanding the properties of the deep mantle, which could be important in gaining understanding about the formation of the earth [19]. Studies have been done by Lacks and Van Orman dealing with the behavior of diffusion and isotope fractionation in silicate melts [20], as well as diffusion of Al in the bulk of periclase [21]. Viscosity, chemical diffusivity and partial molar volumes of silicate liquids near interfaces were found as a function of pressure through MD Simulation as well [22].

We propose a study of grain boundary diffusion in periclase through experimental methods and through simulation techniques, in order to increase the available information and gain further insight on the mechanism and factors in grain boundary diffusion.

References

- [1] R. E. Hoffman and D. Turnbull. Lattice and grain boundary self-diffusion in silver. *Journal of Applied Physics*, 22(5):634–639, 1951. PT: J; NR: 17; TC: 276; J9: J APPL PHYS; PG: 6; GA: UD773; UT: WOS:A1951UD77300020.
- [2] J. C. Fisher. Calculation of diffusion penetration curves for surface and grain boundary diffusion. *Journal of Applied Physics*, 22(1):74–77, 1951. PT: J; NR: 2; TC: 958; J9: J APPL PHYS; PG: 4; GA: UD769; UT: WOS:A1951UD76900014.
- [3] Y.M. Mishin and I. V. Yurovitskii. A generalized-model of grain-boundary diffusion. *Philosophical Magazine A-Physics of Condensed Matter Structure Defects and Mechanical Properties*, 64(6):1239–1249, DEC 1991. PT: J; NR: 18; TC: 12; J9: PHILOS MAG A; PG: 11; GA: GU162; UT: WOS:A1991GU16200004.
- [4] L. Klinger and E. Rabkin. Beyond the fisher model of grain boundary diffusion: Effect of structural inhomogeneity in the bulk. *Acta Materialia*, 47(3):725–734, FEB 5 1999. PT: J; NR: 26; TC: 34; J9: ACTA MATER; PG: 10; GA: 170BQ; UT: WOS:000078785900001.
- [5] D. TURNBULL and RE HOFFMAN. The effect of relative crystal and boundary orientations on grain boundary diffusion rates. *Acta Metallurgica*, 2(3):419–426, 1954. PT: J; NR: 16; TC: 309; J9: ACTA METALL MATER; PG: 8; GA: UL520; UT: WOS:A1954UL52000010.
- [6] R.T.P. Whipple. Concentration contours in grain boundary diffusion. *Philosophical Magazine*, 45(371):1225–1236, 1954. PT: J; NR: 3; TC: 616; J9: PHILOS MAG; PG: 12; GA: UV269; UT: WOS:A1954UV26900001.
- [7] A.D. LeClaire. Analysis of grain boundary diffusion measurements. *British Journal of Applied Physics*, 14(6):351–&, 1963. PT: J; NR: 14; TC: 484; J9: BRIT J APPL PHYS; PG: 0; GA: 7435A; UT: WOS:A19637435A00013.
- [8] Ying Chen and Christopher A. Schuh. Diffusion on grain boundary networks: Percolation theory and effective medium approximations. *Acta Materialia*, 54(18):4709–4720, 10 2006.
- [9] Ying Chen and Christopher A. Schuh. Contribution of triple junctions to the diffusion anomaly in nanocrystalline materials. *Scripta Materialia*, 57(3):253–256, AUG 2007. PT: J; NR: 30; TC: 39; J9: SCRIPTA MATER; PG: 4; GA: 180ZI; UT: WOS:000247406100019.
- [10] L. Klinger and E. Rabkin. Theory of the kirkendall effect during grain boundary interdiffusion. *Acta Materialia*, 59(4):1389–1399, FEB 2011. PT: J; NR: 36; TC: 17; J9: ACTA MATER; PG: 11; GA: 720FE; UT: WOS:000287265100008.
- [11] Y. Mishin and C. Herzig. Grain boundary diffusion: recent progress and future research. *Materials Science and Engineering A-Structural Materials Properties Microstructure and Processing*, 260(1-2):55–71, FEB 1999. PT: J; CT: DOE Panel on Time-Dependent Interfacial Decohesion; CY: JUN 24-27, 1997; CL: GREAT OAKS LANDING, MARYLAND; NR: 59; TC: 119; J9: MAT SCI ENG A-STRUCT; PG: 17; GA: 168HJ; UT: WOS:000078685900007.
- [12] ms green. markoff random processes and the statistical mechanics of time-dependent phenomena .2. irreversible processes in fluids. *journal of chemical physics*, 22(3):398–413, 1954. pt: j; nr: 9; tc: 649; j9: j chem phys; pg: 16; ga: uc157; ut: wos:a1954uc15700009.
- [13] ms green. markoff random processes and the statistical mechanics of time-dependent phenomena. *journal of chemical physics*, 20(8):1281–1295, 1952. pt: j; nr: 19; tc: 437; j9: j chem phys; pg: 15; ga: uc102; ut: wos:a1952uc10200015.

- [14] R. KUBO. Statistical-mechanical theory of irreversible processes .1. general theory and simple applications to magnetic and conduction problems. *Journal of the Physical Society of Japan*, 12(6):570–586, 1957. PT: J; NR: 27; TC: 5413; J9: J PHYS SOC JPN; PG: 17; GA: WJ214; UT: WOS:A1957WJ21400003.
- [15] R. KUBO, M. YOKOTA, and S. NAKAJIMA. Statistical-mechanical theory of irreversible processes .2. response to thermal disturbance. *Journal of the Physical Society of Japan*, 12(11):1203–1211, 1957. PT: J; NR: 18; TC: 397; J9: J PHYS SOC JPN; PG: 9; GA: WJ219; UT: WOS:A1957WJ21900005.
- [16] S. NOSE. A unified formulation of the constant temperature molecular-dynamics methods. *Journal of Chemical Physics*, 81(1):511–519, 1984. PT: J; NR: 27; TC: 5078; J9: J CHEM PHYS; PG: 9; GA: TA661; UT: WOS:A1984TA66100062.
- [17] Gaurav Goel, Liquan Zhang, Daniel J. Lacks, and James A. Van Orman. Isotope fractionation by diffusion in silicate melts: Insights from molecular dynamics simulations. *Geochimica et Cosmochimica Acta*, 93:205–213, SEP 15 2012. PT: J; NR: 38; TC: 3; J9: GEOCHIM COSMOCHIM AC; PG: 9; GA: 000YK; UT: WOS:000308426800012.
- [18] MR Sørensen, Y. Mishin, and AF Voter. Diffusion mechanisms in cu grain boundaries. *Physical Review B*, 62(6):3658–3673, AUG 1 2000. PT: J; NR: 45; TC: 113; J9: PHYS REV B; PG: 16; GA: 343XN; UT: WOS:000088727900017.
- [19] M. W. Ammann, J. P. Brodholt, J. Wookey, and D. P. Dobson. First-principles constraints on diffusion in lower-mantle minerals and a weak d'' layer. *Nature*, 465(7297):462–465, May 2010.
- [20] Daniel J. Lacks, Gaurav Goel, Charles J. Bopp, James A. Van Orman, Charles E. Lesher, and Craig C. Lundstrom. Isotope fractionation by thermal diffusion in silicate melts. *Physical Review Letters*, 108(6):065901, FEB 10 2012. PT: J; NR: 23; TC: 11; J9: PHYS REV LETT; PG: 5; GA: 889VL; UT: WOS:000300102900019.
- [21] James A. Van Orman, Chen Li, and Katherine L. Crispin. Aluminum diffusion and al-vacancy association in periclase. *Physics of the Earth and Planetary Interiors*, 172(1-2):34–42, JAN 2009. PT: J; NR: 29; TC: 13; J9: PHYS EARTH PLANET IN; PG: 9; GA: 397JS; UT: WOS:000262659600007.
- [22] Daniel J. Lacks, David B. Rear, and James A. Van Orman. Molecular dynamics investigation of viscosity, chemical diffusivities and partial molar volumes of liquids along the mgo-sio2 join as functions of pressure. *Geochimica et Cosmochimica Acta*, 71(5):1312–1323, MAR 1 2007. PT: J; NR: 38; TC: 37; J9: GEOCHIM COSMOCHIM AC; PG: 12; GA: 146AE; UT: WOS:000244906100016.