

Controlling Quantum Electrodynamics in Circularly Polarized High Harmonic Generation: Bright, High-Energy Attosecond Waveforms with Tailored Spectro-Temporal Polarization Properties

JILA
CU Boulder and NIST

Kevin M. Dorney¹, Tingting Fan¹, Jennifer L. Ellis¹, Daniel D. Hickstein¹, Christopher A. Mancuso¹, Nathan Brooks¹, Dmitriy Zusin¹, Christian Gentry¹, Ronny Knut¹, Patrik Grychtol¹, Tenio Popmintchev¹, Carlos Hernández-García², Dejan Milošević^{3,4,5}, Henry C. Kapteyn¹, and Margaret M. Murnane¹

¹JILA - Department of Physics, University of Colorado and NIST, Boulder, Colorado, 80309, USA

²Grupo de Investigación en Aplicaciones del Láser y Fótónica, Departamento de Física Aplicada, Universidad de Salamanca, E-37008 Salamanca, Spain

³Academy of Sciences and Arts of Bosnia and Herzegovina, Bistrik 7, 7100 Sarajevo, Bosnia and Herzegovina


⁴Faculty of Science, University of Sarajevo, Zmaja od Bosne 35, 71000 Sarajevo, Bosnia and Herzegovina

⁵Max-Born-Institut, Max-Born-Strasse 2a, 12489 Berlin, Germany



ABSTRACT

- MOTIVATION** - Circularly polarized high-harmonic generation (CPHHG) has recently emerged as a breakthrough light science technique to produce laser-like beams of high-energy, ultrashort, circularly polarized light on a table-top scale system. Typically, CPHHG results in a comb of high-harmonics with alternating circularities, while the attosecond pulse trains (APTs) are *linearly* polarized, thus precluding CPHHG-based studies of sub-fs chiral dynamics.




- EXPERIMENT** - We present experimental and theoretical efforts that demonstrate active control over the quantum electrodynamics in CPHHG, resulting in full control over the spectrotemporal polarization properties of the generated high-harmonics.

RESULTS

1. The spectral helicity distribution in CPHHG can be actively controlled via the intensity ratio of the bicircular driving field, resulting in direct control over the polarization of the underlying APTs.
2. Collective multielectron effects can be exploited in CPHHG to yield a bright harmonic spectrum composed of a single helicity, thus generating circularly polarized APTs..

BICIRCULAR DRIVEN CPHHG

- Single-stage, high-energy Ti:sapphire amplifier (790 nm, 9 mJ, 45 fs, 1 kHz).
- Second harmonic generation (395 nm) and short-wave IR OPA (1300 nm or 2000 nm).
- Fundamental and SHG/OPA beams spatiotemporally overlapped in a collinear geometry in a gas jet (790 nm + 395 nm) or hollow core waveguide (790 nm + OPA).



REFERENCES

¹Dorney, K. M. et al. Helicity-selective enhancement and polarization control of attosecond high harmonic waveforms driven by bichromatic circularly polarized laser fields. *Phys. Rev. Lett.* **119**, 063201 (2017).

²Dorney, K. M. et al. Electronic structure induced spectrotemporal shaping of circularly polarized attosecond waveforms. *In preparation.*

³Higuet, J. et al. High-order harmonic spectroscopy of the Cooper minimum in argon. *Phys. Rev. A* **83**, 053401 (2011).

⁴Schoun, S.B. Attosecond pulse shaping around a Cooper minimum. *Phys. Rev. Lett.* **112**, 153001 (2014).


CONTROLLING QUANTUM ELECTRODYNAMICS IN CPHHG: CUSTOM SPECTROTEMPORAL WAVEFORMS FOR ATTOSECOND CHIRAL SPECTROSCOPY

CONTROLLING THE POLARIZATION STATE OF ATTOSECOND HIGH HARMONIC WAVEFORMS¹


- The polarization of the underlying APTs produced via CPHHG is directly coupled to the spectral intensities of RCP and LCP harmonics.

$$I_{q,RCP} \approx I_{q,LCP} \Rightarrow \text{Linear APTs!} \quad I_{q,RCP} \neq I_{q,LCP} \Rightarrow \text{Elliptical APTs!}$$


- By simply altering the intensity ratio, I_B/I_R , of the bicircular field, we can enhance either RCP or LCP harmonics, while still preserving their circularity!



- Numerical SFA calculations allow for direct access to the subcycle dynamics of the APTs as the intensity ratio is varied.



- Remarkably, a simple perturbative photon model for CPHHG emission accurately recaptures the spectral control afforded by altering the intensity ratio.



CONCLUSIONS AND OUTLOOK


- We demonstrate active control over the spectrotemporal structure of high-harmonic waveforms produced via CPHHG, yielding user-defined harmonic beams for ultrafast chiral spectroscopies.
- The methods presented here are both straightforward and robust, allowing for easy integration into existing setups.
- Future work involves a rigorous theoretical treatment of the CM effect in Ar (and other species) and extension of these techniques to ultraviolet and mid-infrared driven CPHHG, thus enabling transient absorption studies of sub-fs chiral dynamics.

EXPLOITING ELECTRONIC STRUCTURE TO YIELD SINGLE HELICITY ATTOSECOND PULSE TRAINS²


- Formally, CPHHG emission is the result of bound-free dipole transitions from laser-dressed continuum states to the ground state of the parent ion (e.g. Ar⁺).

$$I_{3q \pm 1} \propto |\sqrt{c_+} \langle \Psi_k | \hat{f}_\pm | 3p_+ \rangle + \sqrt{c_-} \langle \Psi_k | \hat{f}_\pm | 3p_- \rangle|^2 \quad \Psi_k^+ \propto R_{kl}(r) Y_{lm}(\hat{r}) Y_{lm}^*(\hat{k})$$


- Thus, electronic structure effects can be exploited to reshape the spectrotemporal structure of the emitted harmonics in a natural way.



- By employing short-wave IR drivers from an OPA, we extend the CPHHG cutoff in Ar beyond the Cooper minimum (CM), resulting in a sharp helicity reversal in the spectral amplitudes of the harmonics.



- Under optimal conditions, the CM effect can be exploited to yield harmonics of a single helicity, which correspond to **completely circular APTs!**



- Preliminary SFA-based calculations accurately recapture the CM-induced helicity reversion in Ar.