

# Limits of foliations in quasi-Fuchsian manifolds

Keaton Quinn

University of Illinois at Chicago

June 10, 2020

# Introduction


Let  $S$  be an oriented closed surface of genus at least 2.

## Introduction

Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.


# Introduction

Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.



## Introduction


Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.



# Introduction

Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.

$$\tilde{M} \simeq \mathbb{H}^3 \simeq \mathbb{B}^3$$




$$\pi_1(M) \cong \pi_1(S) \longrightarrow \Gamma \subset \mathrm{SL}_2 \mathbb{C}$$

# Introduction

Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.

$$\tilde{M} \simeq \mathbb{H}^3 \simeq \mathbb{B}^3 \quad \partial(\mathbb{H}^3) \simeq \mathbb{CP}^1$$




$$\pi_1(M) \simeq \pi_1(S) \longrightarrow \Gamma \subset \text{SL}_2 \mathbb{C}$$

# Introduction

Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.

$$\tilde{M} \simeq \mathbb{H}^3 \simeq \mathbb{B}^3 \quad \partial(\mathbb{H}^3) \simeq \mathbb{CP}^1$$




$$\pi_1(M) \simeq \pi_1(S) \longrightarrow \Gamma \subset SL_2 \mathbb{C}$$

# Introduction

Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.


$$\tilde{M} \simeq \mathbb{H}^3 \simeq \mathbb{B}^3 \quad \partial^*(\mathbb{H}^3) \simeq \mathbb{CP}^1$$



$$\pi_1(M) \simeq \pi_1(S) \longrightarrow \Gamma \subset SL_2 \mathbb{C}$$

# Introduction


Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.



$$\pi_1(M) \cong \pi_1(S) \longrightarrow \Gamma \subset SL_2 \mathbb{C}$$

# Introduction


Let  $S$  be an oriented closed surface of genus at least 2. A quasi-Fuchsian structure  $M$  on  $S \times (0, 1)$  is a complete hyperbolic metric such that there exists a nonempty, compact, geodesically convex subset.



# Introduction


$X = \Omega_+/\Gamma$  and  $Y = \Omega_-/\Gamma$  are called the surfaces at infinity and they inherit both conformal structures and complex projective structures.

These induce  $[X]$  and  $[Y]$  in  $\mathcal{T}(S)$ , the Teichmüller space of  $S$  and we get holomorphic quadratic differentials  $\phi_X$  and  $\phi_Y$  that parametrize the projective structures.




# Introduction

We will consider certain foliations of the ends of  $M$  and investigate their limits as the foliations leave the ends.




We will focus on just one end of  $M$ . From now on let  $X$  denote its surface at infinity and  $\phi$  its holomorphic quadratic differential.


# Visual Metrics




# Visual Metrics




# Visual Metrics




# Visual Metrics




# Visual Metrics




# Epstein Surfaces




# Epstein Surfaces




# Epstein Surfaces




# Epstein Surfaces




# Epstein Surfaces




# Epstein Surfaces



# Epstein Surfaces



# Epstein Surfaces




# Epstein Surfaces

## Theorem

Let  $\Omega$  be a domain in  $\mathbb{CP}^1$  and  $\sigma$  a  $C^k$  conformal metric on  $\Omega$ , then there exists a unique  $C^{k-1}$  map  $\text{Ep}_\sigma : \Omega \rightarrow \mathbb{H}^3$ , called the Epstein map of  $\Omega$  for the metric  $\sigma$ , such that for all  $z \in \Omega$ ,

$$V_{\text{Ep}_\sigma(z)}(z) = \sigma(z).$$

Moreover, the image of a point  $z$  depends only on the 1-jet of  $\sigma$  at  $z$ .




## Properties

This construction is equivariant with respect to the actions of  $\mathrm{SL}_2\mathbb{C}$  on  $\mathbb{CP}^1$  and  $\mathbb{H}^3$ .

# Properties

This construction is equivariant with respect to the actions of  $\mathrm{SL}_2\mathbb{C}$  on  $\mathbb{CP}^1$  and  $\mathbb{H}^3$ .


$$\mathbb{H}^3 \simeq \mathbb{C} \times \mathbb{R}^+$$



## Properties

This construction is equivariant with respect to the actions of  $\mathrm{SL}_2\mathbb{C}$  on  $\mathbb{CP}^1$  and  $\mathbb{H}^3$ .


$$\mathbb{H}^3 \simeq \mathbb{C} \times \mathbb{R}^+$$



# Properties

This construction is equivariant with respect to the actions of  $\mathrm{SL}_2\mathbb{C}$  on  $\mathbb{CP}^1$  and  $\mathbb{H}^3$ .


$$\mathbb{H}^3 \simeq \mathbb{C} \times \mathbb{R}^+$$



# Properties

This construction is equivariant with respect to the actions of  $\mathrm{SL}_2\mathbb{C}$  on  $\mathbb{CP}^1$  and  $\mathbb{H}^3$ .


$$\mathbb{H}^3 \simeq \mathbb{C} \times \mathbb{R}^+$$



# Properties

This construction is equivariant with respect to the actions of  $\mathrm{SL}_2\mathbb{C}$  on  $\mathbb{CP}^1$  and  $\mathbb{H}^3$ .


$$\mathbb{H}^3 \simeq \mathbb{C} \times \mathbb{R}^+$$



## Properties

From  $\text{Ep}_{N_*\sigma}(N \cdot z) = N \cdot \text{Ep}_\sigma(z)$  we get a variant of this Epstein construction for quotients.


In the quasi-Fuchsian case this means if we take a conformal metric  $\sigma$  on the surface at infinity  $X$ , we get an Epstein surface  $\text{Ep}_\sigma : X \rightarrow M$ .  
 (Caution: we say surface even though it may fail to be immersed)



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



# Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



# Properties

The Epstein construction behaves well under a normal flow.


Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



# Properties

The Epstein construction behaves well under a normal flow.

Flowing  $\text{Ep}_\sigma$  for time  $t$  in the normal direction gives a parallel copy of the Epstein surface at distance  $t$  from the original copy. This parallel surface is also obtained as the Epstein surface  $\text{Ep}_{e^{2t}\sigma}$ .



## The Poincaré Family

We call the unique hyperbolic conformal metric  $h$  on  $X$  the Poincaré metric and its Epstein surface  $\text{Ep}_h : X \rightarrow M$  the Poincaré surface of  $X$ .

## The Poincaré Family

We call the unique hyperbolic conformal metric  $h$  on  $X$  the Poincaré metric and its Epstein surface  $\text{Ep}_h : X \rightarrow M$  the Poincaré surface of  $X$ .

The family obtained by parallel flowing  $\text{Ep}_h$  we call the Poincaré Family, and this family is given as  $(\text{Ep}_{e^{2t}h})_{t \geq 0}$ .

## The Poincaré Family

We call the unique hyperbolic conformal metric  $h$  on  $X$  the Poincaré metric and its Epstein surface  $\text{Ep}_h : X \rightarrow M$  the Poincaré surface of  $X$ .

The family obtained by parallel flowing  $\text{Ep}_h$  we call the Poincaré Family, and this family is given as  $(\text{Ep}_{e^{2t}h})_{t \geq 0}$ .

For  $t > c = c(\phi, h)$ , the surfaces  $\text{Ep}_{e^{2t}h}$  are embedded, and so for  $t$  sufficiently large, this family forms a foliation of the end of  $M$ .

## The Poincaré Family

Notice that the family of conformal metrics  $e^{2t} h$  for the Poincaré family has the property that it can be rescaled to be equal to  $h$  for all  $t$ .

## The Poincaré Family

Notice that the family of conformal metrics  $e^{2t}h$  for the Poincaré family has the property that it can be rescaled to be equal to  $h$  for all  $t$ .

Specifically, define  $\epsilon = e^{-2t}$  then the conformal metrics  $\rho_\epsilon$  of the Poincaré family satisfy  $\epsilon\rho_\epsilon = h$  for all  $\epsilon$ .

## The Poincaré Family

Notice that the family of conformal metrics  $e^{2t}h$  for the Poincaré family has the property that it can be rescaled to be equal to  $h$  for all  $t$ .

Specifically, define  $\epsilon = e^{-2t}$  then the conformal metrics  $\rho_\epsilon$  of the Poincaré family satisfy  $\epsilon\rho_\epsilon = h$  for all  $\epsilon$ .

We can consider weakening this condition to only ask that  $\epsilon\rho_\epsilon \rightarrow h$  as  $\epsilon \rightarrow 0$ .

## The Poincaré Family

Notice that the family of conformal metrics  $e^{2t}h$  for the Poincaré family has the property that it can be rescaled to be equal to  $h$  for all  $t$ .

Specifically, define  $\epsilon = e^{-2t}$  then the conformal metrics  $\rho_\epsilon$  of the Poincaré family satisfy  $\epsilon\rho_\epsilon = h$  for all  $\epsilon$ .

We can consider weakening this condition to only ask that  $\epsilon\rho_\epsilon \rightarrow h$  as  $\epsilon \rightarrow 0$ .

The families of surfaces we consider are in this sense asymptotic to the Poincaré family.

# Asymptotically Poincaré Families

## Definition

Let  $S_\epsilon$  for  $\epsilon \in (0, 1)$  be a family of embedded Epstein surfaces for the conformal metrics  $\sigma(\epsilon)$ . We call this family asymptotically Poincaré if

# Asymptotically Poincaré Families

## Definition

Let  $S_\epsilon$  for  $\epsilon \in (0, 1)$  be a family of embedded Epstein surfaces for the conformal metrics  $\sigma(\epsilon)$ . We call this family asymptotically Poincaré if

1. there exists a scaling function  $f : [0, 1) \rightarrow [0, \infty)$  so that the path

$$f\sigma : (0, 1) \rightarrow \text{Met}^\infty(X)$$

is differentiable and converges to  $h$ , as  $\epsilon \rightarrow 0$ , in the space  $\text{Met}^\infty(X)$  of smooth metrics on  $X$ ,

# Asymptotically Poincaré Families

## Definition

Let  $S_\epsilon$  for  $\epsilon \in (0, 1)$  be a family of embedded Epstein surfaces for the conformal metrics  $\sigma(\epsilon)$ . We call this family asymptotically Poincaré if

1. there exists a scaling function  $f : [0, 1) \rightarrow [0, \infty)$  so that the path

$$f\sigma : (0, 1) \rightarrow \text{Met}^\infty(X)$$

is differentiable and converges to  $h$ , as  $\epsilon \rightarrow 0$ , in the space  $\text{Met}^\infty(X)$  of smooth metrics on  $X$ ,

2. the function  $f$  is smooth,  $f(0) = 0$ , and  $f'(0) \neq 0$ ,

# Asymptotically Poincaré Families

## Definition

Let  $S_\epsilon$  for  $\epsilon \in (0, 1)$  be a family of embedded Epstein surfaces for the conformal metrics  $\sigma(\epsilon)$ . We call this family asymptotically Poincaré if

1. there exists a scaling function  $f : [0, 1) \rightarrow [0, \infty)$  so that the path

$$f\sigma : (0, 1) \rightarrow \text{Met}^\infty(X)$$

is differentiable and converges to  $h$ , as  $\epsilon \rightarrow 0$ , in the space  $\text{Met}^\infty(X)$  of smooth metrics on  $X$ ,

2. the function  $f$  is smooth,  $f(0) = 0$ , and  $f'(0) \neq 0$ ,
3. the continuous extension  $\gamma : [0, 1) \rightarrow \text{Met}^\infty(X)$  of  $f\sigma$  is differentiable.

# Asymptotically Poincaré Families Foliate

## Proposition (Q.)

*Let  $S_\epsilon$  be an asymptotically Poincaré family for the conformal metrics  $\sigma(\epsilon)$ . There exists a  $\delta > 0$  (depending on the family) such that for  $\epsilon \in (0, \delta)$  the surfaces  $S_\epsilon$  foliate the end of  $M$ .*

# Asymptotically Poincaré Families Foliate

## Proposition (Q.)

Let  $S_\epsilon$  be an asymptotically Poincaré family for the conformal metrics  $\sigma(\epsilon)$ . There exists a  $\delta > 0$  (depending on the family) such that for  $\epsilon \in (0, \delta)$  the surfaces  $S_\epsilon$  foliate the end of  $M$ .

### Proof.

- ▶ The 1-parameter family of associated Epstein maps gives a map  $\text{Ep}_\sigma : [0, 1) \times X \rightarrow M \cup X$  that restricts to the identity on the boundary  $\{0\} \times X \rightarrow X$ .

# Asymptotically Poincaré Families Foliate

## Proposition (Q.)

Let  $S_\epsilon$  be an asymptotically Poincaré family for the conformal metrics  $\sigma(\epsilon)$ . There exists a  $\delta > 0$  (depending on the family) such that for  $\epsilon \in (0, \delta)$  the surfaces  $S_\epsilon$  foliate the end of  $M$ .

### Proof.

- ▶ The 1-parameter family of associated Epstein maps gives a map  $\text{Ep}_\sigma : [0, 1) \times X \rightarrow M \cup X$  that restricts to the identity on the boundary  $\{0\} \times X \rightarrow X$ .
- ▶ The function given by  $E(\epsilon, x) = \text{Ep}_\sigma(\epsilon^2, x)$  has invertible derivative on the boundary and so it is a local  $C^1$ -diffeomorphism from a neighborhood of the boundary into  $M \cup X$ .

# Asymptotically Poincaré Families Foliate

## Proposition (Q.)

Let  $S_\epsilon$  be an asymptotically Poincaré family for the conformal metrics  $\sigma(\epsilon)$ . There exists a  $\delta > 0$  (depending on the family) such that for  $\epsilon \in (0, \delta)$  the surfaces  $S_\epsilon$  foliate the end of  $M$ .

### Proof.

- ▶ The 1-parameter family of associated Epstein maps gives a map  $\text{Ep}_\sigma : [0, 1) \times X \rightarrow M \cup X$  that restricts to the identity on the boundary  $\{0\} \times X \rightarrow X$ .
- ▶ The function given by  $E(\epsilon, x) = \text{Ep}_\sigma(\epsilon^2, x)$  has invertible derivative on the boundary and so it is a local  $C^1$ -diffeomorphism from a neighborhood of the boundary into  $M \cup X$ .
- ▶ Therefore, the restriction of  $E$  to  $[0, \delta^2) \times X$ , for some small enough  $\delta$ , is a diffeomorphism onto a collar neighborhood of  $X$  in  $M \cup X$ .

# Main Results

## Theorem (Q.)

*Let  $S_\epsilon$  for  $\epsilon \in (0, 1)$  be an asymptotically Poincaré family of surfaces for the conformal metrics  $\sigma(\epsilon)$ . If  $h$  is the Poincaré metric of  $X$  and  $\phi$  the holomorphic quadratic differential at infinity, then in Teichmüller space we have*

$$[I(\sigma(\epsilon))] \rightarrow [h] \quad \text{and} \quad [II(\sigma(\epsilon))] \rightarrow [h] \quad \text{as } \epsilon \rightarrow 0,$$

*where  $[g]$  denotes the point in  $\mathcal{T}(X)$  represented by the Riemannian metric  $g$ .*

*Moreover, the tangent vectors at  $\epsilon = 0$  in  $T_{[h]}\mathcal{T}(X)$  are given by*

$$[\dot{I}(\sigma(\epsilon))] = 4f'(0)\operatorname{Re}(\phi) \quad \text{and} \quad [\dot{II}(\sigma(\epsilon))] = 0.$$

# Riemannian Model of $\mathcal{T}(X)$

We model  $\mathcal{T}(X)$  as the quotient

$$\text{Met}^\infty(X)/(\text{Diff}_0^\infty(X) \ltimes P^\infty(X))$$

of smooth metrics on  $X$  by the semi-direct product of  $\text{Diff}_0^\infty(X)$ , the group of smooth diffeomorphisms isotopic to the identity, and  $P^\infty(X)$ , the group of smooth positive functions on  $X$ .

# Riemannian Model of $\mathcal{T}(X)$

We model  $\mathcal{T}(X)$  as the quotient

$$\text{Met}^\infty(X)/(\text{Diff}_0^\infty(X) \ltimes P^\infty(X))$$

of smooth metrics on  $X$  by the semi-direct product of  $\text{Diff}_0^\infty(X)$ , the group of smooth diffeomorphisms isotopic to the identity, and  $P^\infty(X)$ , the group of smooth positive functions on  $X$ .

However, we will first work with Sobolev tensors.

## Riemannian Model of $\mathcal{T}(X)$

We model  $\mathcal{T}(X)$  as the quotient

$$\text{Met}^\infty(X)/(\text{Diff}_0^\infty(X) \ltimes P^\infty(X))$$

of smooth metrics on  $X$  by the semi-direct product of  $\text{Diff}_0^\infty(X)$ , the group of smooth diffeomorphisms isotopic to the identity, and  $P^\infty(X)$ , the group of smooth positive functions on  $X$ .

However, we will first work with Sobolev tensors.

Denote by  $\text{Met}^s(X)$  the set of Riemannian metrics of Sobolev class  $H^s$  for a fixed  $s > 3$  and note that  $\text{Met}^\infty(X) = \cap_{s>3} \text{Met}^s(X)$

With these regularity assumptions  $T_h \text{Met}^s(X)$  is a Hilbert space.

## Riemannian Model of $\mathcal{T}(X)$


$T_h \text{Met}^s(X)$  decomposes as the orthogonal direct sum

$$\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \oplus \{\mathcal{L}_V h + uh \mid u \in H^s, V \in \Gamma(TX)\}.$$

## Riemannian Model of $\mathcal{T}(X)$

$T_h \text{Met}^s(X)$  decomposes as the orthogonal direct sum

$$\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \oplus \{\mathcal{L}_V h + uh \mid u \in H^s, V \in \Gamma(TX)\}.$$




$$\Gamma^s(\Sigma^2)$$

## Riemannian Model of $\mathcal{T}(X)$

$T_h \text{Met}^s(X)$  decomposes as the orthogonal direct sum

$$\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \oplus \{\mathcal{L}_V h + uh \mid u \in H^s, V \in \Gamma(TX)\}.$$




$$\Gamma^s(\Sigma^2)$$

# Riemannian Model of $\mathcal{T}(X)$

$T_h \text{Met}^s(X)$  decomposes as the orthogonal direct sum

$$\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \oplus \{\mathcal{L}_V h + uh \mid u \in H^s, V \in \Gamma(TX)\}.$$




$$\Gamma^s(\Sigma^2)$$

## Riemannian Model of $\mathcal{T}(X)$

$T_h \text{Met}^s(X)$  decomposes as the orthogonal direct sum

$$\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \oplus \{\mathcal{L}_V h + uh \mid u \in H^s, V \in \Gamma(TX)\}.$$




$$\Gamma^s(\Sigma^2)$$

# Riemannian Model of $\mathcal{T}(X)$

$T_h \text{Met}^s(X)$  decomposes as the orthogonal direct sum

$$\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \oplus \{\mathcal{L}_V h + uh \mid u \in H^s, V \in \Gamma(TX)\}.$$



$$\Gamma^s(\Sigma)$$

## Riemannian Model of $\mathcal{T}(X)$

The set  $\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\}$  is a model for the tangent space to the quotient  $\text{Met}^s(X)/(\text{Diff}_0^s(X) \ltimes P^s(X))$  at  $[h]$ .

This is true for any  $s$ , and each such  $g$  is a smooth tensor. So this set is identified with the tangent space to Teichmüller space.

## Riemannian Model of $\mathcal{T}(X)$

The set  $\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\}$  is a model for the tangent space to the quotient  $\text{Met}^s(X)/(\text{Diff}_0^s(X) \ltimes P^s(X))$  at  $[h]$ .

This is true for any  $s$ , and each such  $g$  is a smooth tensor. So this set is identified with the tangent space to Teichmüller space.

$$\begin{aligned} T_{[h]}\mathcal{T}(X) &= \{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \\ &= \{\text{Re}(\psi) \mid \psi \text{ a holomorphic quadratic differential on } X\}. \end{aligned}$$

## Riemannian Model of $\mathcal{T}(X)$

The set  $\{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\}$  is a model for the tangent space to the quotient  $\text{Met}^s(X)/(\text{Diff}_0^s(X) \ltimes P^s(X))$  at  $[h]$ .

This is true for any  $s$ , and each such  $g$  is a smooth tensor. So this set is identified with the tangent space to Teichmüller space.

$$\begin{aligned} T_{[h]}\mathcal{T}(X) &= \{g \mid \text{tr}_h(g) = 0 = \text{div}_h(g)\} \\ &= \{\text{Re}(\psi) \mid \psi \text{ a holomorphic quadratic differential on } X\}. \end{aligned}$$

The projection  $\text{Met}^\infty(X) \rightarrow \mathcal{T}(X)$  is continuous and its derivative at  $h$  is given by orthogonal projection onto  $\{\text{Re}(\psi)\}$

## Proof

Returning to our theorem, we are given that  $\gamma : [0, 1) \rightarrow \text{Met}^\infty(X)$  is continuous and differentiable at  $\epsilon = 0$ . Therefore, we also have that  $\gamma$  is continuous to  $\text{Met}^s(X)$ , for each  $s$ , and is differentiable at  $\epsilon = 0$ .

## Proof

Returning to our theorem, we are given that  $\gamma : [0, 1) \rightarrow \text{Met}^\infty(X)$  is continuous and differentiable at  $\epsilon = 0$ . Therefore, we also have that  $\gamma$  is continuous to  $\text{Met}^s(X)$ , for each  $s$ , and is differentiable at  $\epsilon = 0$ .

In  $\text{Met}^s(X)$  we can write

$$I(\sigma(\epsilon)) = \frac{1}{4\epsilon f'(0)}h + \frac{1}{4f'(0)}\dot{y} + \frac{1}{2}h + \text{Re}(\phi) + O(\epsilon),$$

## Proof

Returning to our theorem, we are given that  $\gamma : [0, 1) \rightarrow \text{Met}^\infty(X)$  is continuous and differentiable at  $\epsilon = 0$ . Therefore, we also have that  $\gamma$  is continuous to  $\text{Met}^s(X)$ , for each  $s$ , and is differentiable at  $\epsilon = 0$ .

In  $\text{Met}^s(X)$  we can write

$$I(\sigma(\epsilon)) = \frac{1}{4\epsilon f'(0)}h + \frac{1}{4f'(0)}\dot{\gamma} + \frac{1}{2}h + \text{Re}(\phi) + O(\epsilon),$$

so that

$$I_\epsilon := 4\epsilon f'(0)I(\sigma(\epsilon)) = h + \epsilon(\dot{\gamma} + 2f'(0)h + 4f'(0)\text{Re}(\phi)) + O(\epsilon^2).$$

## Proof

From the expression


$$I_\epsilon = h + \epsilon(\dot{\gamma} + 2f'(0)h + 4f'(0)\operatorname{Re}(\phi)) + O(\epsilon^2)$$

we can see that  $I_\epsilon \rightarrow h$  in  $\operatorname{Met}^s(X)$  for all  $s$ , implying  $I_\epsilon \rightarrow h$  in  $\operatorname{Met}^\infty(X)$ .


Moreover, the derivative at  $\epsilon = 0$  is

$$\dot{I}_\epsilon = \dot{\gamma} + 2f'(0)h + 4f'(0)\operatorname{Re}(\phi).$$


# Proof




## Proof




## Proof



## Proof



## Proof



Therefore, as  $I_\epsilon \rightarrow h$  in  $\text{Met}^\infty(X)$  we have that  $[I(\sigma(\epsilon))] = [I_\epsilon] \rightarrow [h]$  in  $\mathcal{T}(X)$ .

Therefore, as  $I_\epsilon \rightarrow h$  in  $\text{Met}^\infty(X)$  we have that  $[I(\sigma(\epsilon))] = [I_\epsilon] \rightarrow [h]$  in  $\mathcal{T}(X)$ .

And since

$$\dot{I}_\epsilon = \dot{\gamma} + 2f'(0)h + 4f'(0)\text{Re}(\phi),$$

we see

$$[\dot{I}(\sigma(\epsilon))] = 4f'(0)\text{Re}(\phi).$$

Therefore, as  $I_\epsilon \rightarrow h$  in  $\text{Met}^\infty(X)$  we have that  $[I(\sigma(\epsilon))] = [I_\epsilon] \rightarrow [h]$  in  $\mathcal{T}(X)$ .

And since

$$\dot{I_\epsilon} = \dot{\gamma} + 2f'(0)h + 4f'(0)\text{Re}(\phi),$$

we see

$$[\dot{I(\sigma(\epsilon))}] = 4f'(0)\text{Re}(\phi).$$

If we apply the same arguments to the Taylor series for  $\dot{II}(\sigma(\epsilon))$  then we see  $[\dot{II}(\sigma(\epsilon))] \rightarrow [h]$  and  $[\dot{II}(\sigma(\epsilon))] = 0$ .



# Applications

We apply this result to two foliations by constant curvature surfaces.

# Applications

We apply this result to two foliations by constant curvature surfaces.

## Theorem (Labourie, 1992)

*Let  $E$  be an end of a quasi-Fuchsian manifold  $M$ , then for each  $k \in (-1, 0)$ , there exists a unique (incompressible) surface embedded in  $E$  with constant Gaussian curvature  $k$ . Moreover, this family of surfaces foliates the end  $E$ .*

# Applications

We apply this result to two foliations by constant curvature surfaces.

## Theorem (Labourie, 1992)


*Let  $E$  be an end of a quasi-Fuchsian manifold  $M$ , then for each  $k \in (-1, 0)$ , there exists a unique (incompressible) surface embedded in  $E$  with constant Gaussian curvature  $k$ . Moreover, this family of surfaces foliates the end  $E$ .*

## Theorem (Mazzeo-Pacard, 2011)


*Each end of a quasi-Fuchsian manifold admits a unique foliation by constant mean curvature surfaces.*

# A Conjecture of Labourie

Labourie called the constant Gaussian curvature surfaces  $k$ -surfaces and he discusses how their first and second fundamental forms may be considered as paths in Teichmüller space.



# A Conjecture of Labourie



He shows that as  $k \rightarrow 0$ , the paths  $[I_k]$  and  $[II_k]$  converge to  $[X] = [h]$  and he asks after the tangent vectors to these paths at  $k = 0$ .

He conjectures  $[I_k]$  is related to the holomorphic quadratic differential  $\phi$ .

# A Conjecture of Labourie

## Theorem (Q.)

*Let  $I_k$  and  $\Pi_k$  be the first and second fundamental forms of the family of  $k$ -surfaces in an end of a quasi-Fuchsian manifold. Let  $\phi$  be the holomorphic quadratic differential at infinity of  $M$ . Then, as  $k \rightarrow 0$ , the tangent vectors to  $[I_k]$  and  $[\Pi_k]$  in Teichmüller space are given by*

$$[\dot{I}_k] = -\operatorname{Re}(\phi) \quad \text{and} \quad [\dot{\Pi}_k] = 0.$$

# A Conjecture of Labourie

## Theorem (Q.)

*Let  $I_k$  and  $II_k$  be the first and second fundamental forms of the family of  $k$ -surfaces in an end of a quasi-Fuchsian manifold. Let  $\phi$  be the holomorphic quadratic differential at infinity of  $M$ . Then, as  $k \rightarrow 0$ , the tangent vectors to  $[I_k]$  and  $[II_k]$  in Teichmüller space are given by*

$$[\dot{I}_k] = -\operatorname{Re}(\phi) \quad \text{and} \quad [\dot{II}_k] = 0.$$

This will follow from our main theorem if we can show the  $k$ -surfaces form an asymptotically Poincaré family. To do this we must present the  $k$ -surfaces as Epstein surfaces.

## Proof

The Gaussian curvature of an Epstein surface for a conformal metric  $\sigma$  can be calculated as

$$K(I(\sigma)) = \frac{4K(\sigma)}{(1 - K(\sigma))^2 - 16|B(\sigma)|^2\sigma^{-2}}.$$

## Proof

The Gaussian curvature of an Epstein surface for a conformal metric  $\sigma$  can be calculated as

$$K(I(\sigma)) = \frac{4K(\sigma)}{(1 - K(\sigma))^2 - 16|B(\sigma)|^2\sigma^{-2}}.$$

Here

- ▶  $K(\sigma)$  is the Gaussian curvature of  $\sigma$  itself.

## Proof

The Gaussian curvature of an Epstein surface for a conformal metric  $\sigma$  can be calculated as

$$K(I(\sigma)) = \frac{4K(\sigma)}{(1 - K(\sigma))^2 - 16|B(\sigma)|^2\sigma^{-2}}.$$

Here

- ▶  $K(\sigma)$  is the Gaussian curvature of  $\sigma$  itself.
- ▶  $B(\sigma)$  is the Schwarzian tensor of  $\sigma$ . This is a quadratic differential (not necessarily holomorphic) depending on  $\sigma$  and its first two derivatives.

## Proof

The Gaussian curvature of an Epstein surface for a conformal metric  $\sigma$  can be calculated as

$$K(I(\sigma)) = \frac{4K(\sigma)}{(1 - K(\sigma))^2 - 16|B(\sigma)|^2\sigma^{-2}}.$$

Here

- ▶  $K(\sigma)$  is the Gaussian curvature of  $\sigma$  itself.
- ▶  $B(\sigma)$  is the Schwarzian tensor of  $\sigma$ . This is a quadratic differential (not necessarily holomorphic) depending on  $\sigma$  and its first two derivatives.

To find a  $k$ -surface, we therefore need to solve  $K(I(\sigma)) = k$  for  $\sigma$ .

## Proof

We focus on solving the equivalent equation,

$$4K(\sigma) = k \left( (1 - K(\sigma))^2 - 16|B(\sigma)|^2\sigma^{-2} \right) \quad (*)$$

# Proof

We focus on solving the equivalent equation,

$$4K(\sigma) = k \left( (1 - K(\sigma))^2 - 16|B(\sigma)|^2\sigma^{-2} \right) \quad (*)$$

Our technique is to use the Implicit Function Theorem to obtain solutions for each  $k$  near 0.

Notice though, that when  $k = 0$  this reads  $K(\sigma) = 0$ , which has no solutions on  $X$ .

## Proof

Since there are no solutions for  $k = 0$  we rescale the equation by considering the function  $f(k) = \frac{1-\sqrt{1+k}}{1+\sqrt{1+k}}$ .

## Proof

Since there are no solutions for  $k = 0$  we rescale the equation by considering the function  $f(k) = \frac{1-\sqrt{1+k}}{1+\sqrt{1+k}}$ .

If  $\sigma$  solves  $(*)$  then  $\tau = f(k)\sigma$  solves

$$(2 + k)(1 + K(\tau))^2$$

$$+ 2\sqrt{1+k} \left(1 - K(\tau)^2\right) + 16 \left(2\sqrt{1+k} - 2 - k\right) \frac{|B(\tau)|^2}{\tau^2} = 0$$

or

$$F(k, \tau) = 0.$$

## Proof

Since there are no solutions for  $k = 0$  we rescale the equation by considering the function  $f(k) = \frac{1-\sqrt{1+k}}{1+\sqrt{1+k}}$ .

If  $\sigma$  solves  $(*)$  then  $\tau = f(k)\sigma$  solves

$$(2 + k)(1 + K(\tau))^2$$

$$+ 2\sqrt{1+k} \left(1 - K(\tau)^2\right) + 16 \left(2\sqrt{1+k} - 2 - k\right) \frac{|B(\tau)|^2}{\tau^2} = 0$$

or

$$F(k, \tau) = 0.$$

Note that  $F(0, \tau) = 0$  is equivalent to  $K(\tau) = -1$ , which has the solution  $\tau = h$ .

## Proof

Since we have a solution for  $k = 0$ , we can then use the Implicit Function Theorem to get solution for  $k$  near zero as well.

The function  $F$ , while defined on smooth conformal metrics  $\text{Conf}^\infty(X)$ , extends to one of Sobolev conformal metrics  $\text{Conf}^s(X)$ . We have

$$F : (-1, 0) \times \text{Conf}^s(X) \rightarrow H^{s-2}(X).$$

# Proof

Since we have a solution for  $k = 0$ , we can then use the Implicit Function Theorem to get solution for  $k$  near zero as well.

The function  $F$ , while defined on smooth conformal metrics  $\text{Conf}^\infty(X)$ , extends to one of Sobolev conformal metrics  $\text{Conf}^s(X)$ . We have

$$F : (-1, 0) \times \text{Conf}^s(X) \rightarrow H^{s-2}(X).$$

We compute

$$DF_{(0,h)}(\dot{k}, \dot{\tau}) = 4DK_h(\dot{\tau}) = -2(\Delta_h - Id)\frac{\dot{\tau}}{h}$$

and see that  $D_2F_{(0,h)} = 4DK_h : H^s(X) \rightarrow H^{s-2}(X)$  is an isomorphism.

## Proof

Consequently, there exists a neighborhood  $V$  of 0 and a curve  $\gamma : V \rightarrow \text{Conf}^s(X)$  with  $\gamma(0) = h$  and  $F(k, \gamma(k)) = 0$ .

## Proof

Consequently, there exists a neighborhood  $V$  of 0 and a curve  $\gamma : V \rightarrow \text{Conf}^s(X)$  with  $\gamma(0) = h$  and  $F(k, \gamma(k)) = 0$ .

Elliptic regularity theory applied to  $DK_h$  tells us that there is a  $\delta > 0$  such that

- ▶  $\gamma(k)$  is a smooth conformal metric for each  $k \in (-\delta, \delta)$ ,

## Proof

Consequently, there exists a neighborhood  $V$  of 0 and a curve  $\gamma : V \rightarrow \text{Conf}^s(X)$  with  $\gamma(0) = h$  and  $F(k, \gamma(k)) = 0$ .

Elliptic regularity theory applied to  $DK_h$  tells us that there is a  $\delta > 0$  such that

- ▶  $\gamma(k)$  is a smooth conformal metric for each  $k \in (-\delta, \delta)$ ,
- ▶ the path  $\gamma : (-\delta, \delta) \rightarrow \text{Conf}^\infty(X)$  is a smooth path.

## Proof

Consequently, there exists a neighborhood  $V$  of 0 and a curve  $\gamma : V \rightarrow \text{Conf}^s(X)$  with  $\gamma(0) = h$  and  $F(k, \gamma(k)) = 0$ .

Elliptic regularity theory applied to  $DK_h$  tells us that there is a  $\delta > 0$  such that

- ▶  $\gamma(k)$  is a smooth conformal metric for each  $k \in (-\delta, \delta)$ ,
- ▶ the path  $\gamma : (-\delta, \delta) \rightarrow \text{Conf}^\infty(X)$  is a smooth path.

To complete the proof we take  $\sigma(k) := f(k)^{-1}\gamma(k)$  for  $k \in (-\delta, 0)$ .

The family of Epstein surfaces for  $\sigma(k)$  then satisfy the definition of an asymptotically Poincaré family and our main theorem applies. □

# Constant Mean Curvature Surfaces

## Theorem (Mazzeo-Pacard, 2011)

*The ends of a quasi-Fuchsian manifold admit unique foliations by constant mean curvature surfaces.*

## Theorem (Q.)

*Let  $I_k$  and  $II_k$  be the first and second fundamental forms of the Epstein surface with constant mean curvature  $-\sqrt{1+k}$ . Let  $\phi$  be the holomorphic quadratic differential at infinity. Then, as  $k \rightarrow 0$ , the tangent vectors to  $[I_k]$  and  $[II_k]$  in Teichmüller space are given by*

$$[I_k] = -\operatorname{Re}(\phi) \quad \text{and} \quad [II_k] = 0.$$

## Proof

The proof proceeds the same as in the constant Gaussian curvature case. However, here we wish to solve the equation

$$H(\text{Ep}_\sigma) = \frac{K(\sigma)^2 - 1 - 16|B(\sigma)|^2\sigma^{-2}}{(K(\sigma) - 1)^2 - 16|B(\sigma)|^2\sigma^{-2}} = -\sqrt{1 + k}.$$

## Proof

The proof proceeds the same as in the constant Gaussian curvature case. However, here we wish to solve the equation

$$H(\text{Ep}_\sigma) = \frac{K(\sigma)^2 - 1 - 16|B(\sigma)|^2\sigma^{-2}}{(K(\sigma) - 1)^2 - 16|B(\sigma)|^2\sigma^{-2}} = -\sqrt{1 + k}.$$

Using the same scaling function  $f(k)$  we are led to consider solutions to a function  $G(k, \tau) = 0$ , which has partial derivative

$$D_2G_{(0,h)} = 2DK_h(\dot{\tau}).$$

## Proof

The proof proceeds the same as in the constant Gaussian curvature case. However, here we wish to solve the equation

$$H(\text{Ep}_\sigma) = \frac{K(\sigma)^2 - 1 - 16|B(\sigma)|^2\sigma^{-2}}{(K(\sigma) - 1)^2 - 16|B(\sigma)|^2\sigma^{-2}} = -\sqrt{1 + k}.$$

Using the same scaling function  $f(k)$  we are led to consider solutions to a function  $G(k, \tau) = 0$ , which has partial derivative

$$D_2G_{(0,h)} = 2DK_h(\dot{\tau}).$$

The Implicit Function Theorem then gives solutions for  $k$  near zero. □

Thank you!

