References

- Alvarez-Jimenez, L., Moreno-Cabañas, A., Ramirez-Jimenez, M., Morales-Palomo, F., Ortega, J. F., & Mora-Rodriguez, R. (2021). Effectiveness of statins vs. exercise on reducing postprandial hypertriglyceridemia in dyslipidemic population: A systematic review and network meta-analysis. *Journal of Sport and Health Science*, 1–11.

 https://doi.org/10.1016/j.jshs.2021.07.006
- Cavero-Redondo, I., Deeks, J. J., Alvarez-Bueno C., Jolly K., Saz-Lara A., Price M., et al. (2021) Comparative effect of physical exercise versus statins on improving arterial stiffness in patients with high cardiometabolic risk: A network meta-analysis. *PLoS Med* 18(2): e1003543. https://doi.org/10.1371/journal.pmed.1003543
- Deichmann R. E, Lavie C. J., Asher T., DiNicolantonio J. J., O'Keefe J. H., Thompson P.D. (2015). The Interaction Between Statins and Exercise: Mechanisms and Strategies to Counter the Musculoskeletal Side Effects of This Combination Therapy. *The Ochsner Journal*, 15:429–437. PMID: 26730228; PMCID: PMC4679305.
- Gui, Y.-jun, Liao, C.-xiu, Liu, Q., Guo, Y., Yang, T., Chen, J.-yuan, Wang, Y.-ting, Hu, J.-hui, & Xu, D.-yan. (2017). Efficacy and safety of statins and exercise combination therapy compared to statin monotherapy in patients with dyslipidaemia: A systematic review and meta-analysis. *European Journal of Preventive Cardiology*, 24(9), 907–916. https://doi.org/10.1177/2047487317691874