

Nonstandard Analysis, The 1972-SciAm Article

Ideas That Common People Brand Nonstandard

Martin Davis¹ Reuben Hersh²

¹Original Author

²Original Author

Aug 2025 / Free Learner's School Conversations

Outline

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

1 Introduction

2 Summary

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

Nonstandard Analysis

Common Words, Uncommon Meanings, Yet Again!

- This Is A Review of A Davis-Hersh¹ Article in Scientific American, June 1972.
 - You Perhaps Haven't Even Heard “Nonstandard Analysis” in The Context of Calculus-1.
 - Analysis Refers to Foundations of Calculus.
 - Nonstandard Only Because The Word ‘Standard’ Was Taken to Mean Formalization of Calculus Using Some Other (Beautiful) Way!
- These Slides Are A Gentle Introduction to Nonstandard Analysis, Are for Personal Benefit, Only Indirectly for Actual Presentation.
- There's No Need to Get Intimidated by Big Words.
- Let's Be Fearless in Guided Imagination, and Make Inquiry A Habit.

¹Both Were Mathematicians with a Gift of Writing Extraordinary Expository Articles.

A Bit of History

Everything Has A History!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, **Avoided The *Unthinkable* Infinity (Infinitely Large) & The *Strange* Infinitesimals (Infinitely Small).**
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The **Banned Infinitesimals**:
 - Small, **Infinitely Small** Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These **Infinitesimal Numbers That Exist in A Sort of Neverland!**
- Infinitesimals Defied **Logic Prevalent at The Time!**
- And So, Karl Weierstrass et. al. **Banished Infinitesimals from Calculus by The 1890s!**

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California, is Named

A Bit of History

Everything Has A History!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, **Avoided The *Unthinkable* Infinity (Infinitely Large) & The *Strange* Infinitesimals (Infinitely Small).**
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The *Banned Infinitesimals*:
 - Small, *Infinitely Small* Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
- Infinitesimals Defied *Logic Prevalent at The Time!*
- And So, Karl Weierstrass et. al. **Banished Infinitesimals from Calculus by The 1890s!**

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California, is Named

A Bit of History

Everything Has A History!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, *Avoided The Unthinkable Infinity (Infinitely Large) & The Strange Infinitesimals (Infinitely Small)*.
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The **Banned Infinitesimals**:
 - Small, *Infinitely Small* Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
- Infinitesimals Defied *Logic Prevalent at The Time!*
- And So, Karl Weierstrass et. al. **Banished Infinitesimals from Calculus by The 1890s!**

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California, is Named

A Bit of History

Everything Has A History!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, *Avoided The Unthinkable Infinity (Infinitely Large) & The Strange Infinitesimals (Infinitely Small)*.
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The *Banned Infinitesimals*:
 - Small, **Infinitely Small** Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
 - Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
 - Infinitesimals Defied *Logic Prevalent at The Time!*
 - And So, Karl Weierstrass et. al. *Banished Infinitesimals from Calculus by The 1890s!*

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California, Is Named

A Bit of History

Everything Has A History!

Nonstandard
Analysis

MD, RH, KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, *Avoided The Unthinkable Infinity (Infinitely Large) & The Strange Infinitesimals (Infinitely Small)*.
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The *Banned Infinitesimals*:
 - Small, **Infinitely Small** Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
- Infinitesimals Defied Logic Prevalent at The Time!
- And So, Karl Weierstrass et. al. *Banished Infinitesimals from Calculus by The 1890s!*

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California, Is Named

A Bit of History

Everything Has A History!


Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, *Avoided The Unthinkable Infinity (Infinitely Large) & The Strange Infinitesimals (Infinitely Small)*.
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The *Banned Infinitesimals*:
 - Small, **Infinitely Small** Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
 - Infinitesimals Defied Logic Prevalent at The Time!
 - And So, Karl Weierstrass et. al. *Banished Infinitesimals from Calculus by The 1890s!*

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California is Named! 

A Bit of History

Everything Has A History!


Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, *Avoided The Unthinkable Infinity (Infinitely Large) & The Strange Infinitesimals (Infinitely Small)*.
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The *Banned Infinitesimals*:
 - Small, **Infinitely Small** Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
- Infinitesimals Defied **Logic Prevalent at The Time!**
 - And So, Karl Weierstrass et. al. *Banished Infinitesimals from Calculus by The 1890s!*

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California is Named! 

A Bit of History

Everything Has A History!


Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Influential Ancient Greeks, Euclid, Aristotle, & Archimedes, *Avoided The Unthinkable Infinity (Infinitely Large) & The Strange Infinitesimals (Infinitely Small)*.
- Ideas of Calculus Over Time Culminated into **Leibniz's and Newton's Formulation of Calculus** in The 1680s.
- Both Used The *Banned Infinitesimals*:
 - Small, **Infinitely Small** Non-Archimedean Numbers!
 - Smaller Than Any Finite Number, Yet Bigger Than 0!
- Several Philosophers/Mathematicians Criticized² Leibniz and Newton for These *Infinitesimal Numbers That Exist in A Sort of Neverland!*
- Infinitesimals Defied **Logic Prevalent at The Time!**
- And So, Karl Weierstrass et. al. **Banished Infinitesimals from Calculus by The 1890s!**

²Prominent Among Them Was Bishop George Berkeley, The Famous Clergyman for Whom the University of Berkeley, California is Named! 

A Bit More of History

Everything Has A History!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between **The Continuous And The Discrete** Is Reignited.

A Bit More of History

Everything Has A History!

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between **The Continuous And The Discrete** Is Reignited.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

A Bit More of History

Everything Has A History!

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between **The Continuous And The Discrete** Is Reignited.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

A Bit More of History

Everything Has A History!

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between **The Continuous And The Discrete** Is Reignited.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

A Bit More of History

Everything Has A History!

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between **The Continuous And The Discrete** Is Reignited.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

A Bit More of History

Everything Has A History!

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between The Continuous And The Discrete Is Reignited.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

A Bit More of History

Everything Has A History!

- Weierstrass Formalized Calculus Using The Allegedly Non-intuitive Idea of **Limits**.
- However, In 1966, The Logician Abraham Robinson Reintroduced Infinitesimals And Provided The **Necessary Rigor** to Them!
- Several Mathematicians Criticized Robinson's Work!
- But in 2020s, It Is Considered Original Body of Work That Is **Equivalent To Standard Analysis**; Strangely, Logic Prevalent Now Necessitates Infinitesimals!
- In The End, Your Approach to Calculus Reduces to **Your Philosophy of Mathematics**: Choice Is Yours!
- The War between **The Continuous And The Discrete** Is Reignited.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Does That Make You Uncomfortable?

Be My Guest!

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- On The First Encounter, *Nonstandard Analysis Feels, Well, *Nonstandard*!*
- It Makes Us *Ponder The Nature of Mathematics.*
- Although It Stretches Our Imagination, Its Logic And Equivalence with Standard Analysis Is *Fun!*
 - After All, When *Applied to Practical Problems*, Both Approaches *Produce Exactly The Same Results!*

Does That Make You Uncomfortable?

Be My Guest!

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- On The First Encounter, **Nonstandard Analysis Feels, Well, *Nonstandard*!**
- It Makes Us **Ponder The Nature of Mathematics.**
- Although It Stretches Our Imagination, Its Logic And Equivalence with Standard Analysis Is **Fun!**
 - After All, When *Applied to Practical Problems*, Both Approaches **Produce Exactly The Same Results!**

Does That Make You Uncomfortable?

Be My Guest!

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- On The First Encounter, **Nonstandard Analysis Feels, Well, *Nonstandard*!**
- It Makes Us **Ponder The Nature of Mathematics.**
- Although It Stretches Our Imagination, Its Logic And Equivalence with Standard Analysis Is **Fun!**
 - After All, When *Applied to Practical Problems*, Both Approaches **Produce Exactly The Same Results!**

Does That Make You Uncomfortable?

Be My Guest!

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- On The First Encounter, **Nonstandard Analysis Feels, Well, *Nonstandard*!**
- It Makes Us **Ponder The Nature of Mathematics.**
- Although It Stretches Our Imagination, Its Logic And Equivalence with Standard Analysis Is **Fun!**
 - After All, When *Applied to Practical Problems*, Both Approaches **Produce Exactly The Same Results!**

Does That Make You Uncomfortable?

Be My Guest!

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- On The First Encounter, **Nonstandard Analysis Feels, Well, *Nonstandard*!**
- It Makes Us **Ponder The Nature of Mathematics.**
- Although It Stretches Our Imagination, Its Logic And Equivalence with Standard Analysis Is **Fun!**
 - After All, When *Applied to Practical Problems*, Both Approaches **Produce Exactly The Same Results!**

Geometry

Let's Introduce Infinitesimals in Geometry.




The Area And Circumference of The Unit Circle

Argument 1: A *Logically Unacceptable* Argument Based on Infinitesimals

Area of Unit Circle = $\frac{1}{2}$ Its Circumference
(Would Euclid Accept It?)

Any circle can be thought of as composed of **infinitely many straight-line segments**, all equal to each other and **infinitely short**. It is then the sum of **infinitesimal triangles**, all of which have altitude 1.

Area of a triangle = half the base times the altitude. Therefore, the sum of the areas of the triangles is half the sum of the bases. But the sum of the areas of the triangles is the area of the circle, and the sum of the bases of the triangles is its circumference. **Therefore, the area of the unit circle = half its circumference.** \square



Seriously?

Is That (**Argument 1** [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by **Nicholas of Cusa** in The 15th Century!
- We Wonder If It Is Even Logical! Clearly, **Objections Galore!**
 - The **Very Notion of A Triangle with An Infinitely Small Base Is Illusive!**
 - The Base of The Triangle, b , **However Small**, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, **No Number of Triangles Can Make A Circle with Positive Circumference!**
 - ② If $b > 0$, **Infinitely Many Triangles Make An Infinitely Large Circumference!**
 - In Either Case, How Can We Ever Get **A Finite Circle from Infinite Infinitely Small Pieces?**
 - ① Every Real Number Must Be Archimedean!
 - ② Since **No Infinitesimal Is Archimedean**, They Were Rejected by Euclid, Aristotle, Archimedes.

Seriously?

Is That (Argument 1 [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by Nicholas of Cusa in The 15th Century!
- We Wonder If It Is Even Logical! Clearly, Objections Galore!
 - The Very Notion of A Triangle with An Infinitely Small Base Is Illusive!
 - The Base of The Triangle, b , However Small, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, No Number of Triangles Can Make A Circle with Positive Circumference!
 - ② If $b > 0$, Infinitely Many Triangles Make An Infinitely Large Circumference!
 - In Either Case, How Can We Ever Get A Finite Circle from Infinite Infinitely Small Pieces?
 - ① Every Real Number Must Be Archimedean!
 - ② Since No Infinitesimal Is Archimedean, They Were Rejected by Euclid, Aristotle, Archimedes.

Seriously?

Is That (**Argument 1** [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by **Nicholas of Cusa in The 15th Century!**
- We Wonder If It Is Even Logical! Clearly, **Objections Galore!**
 - The Very Notion of A Triangle with An Infinitely Small Base Is Illusive!
 - The Base of The Triangle, b , However Small, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, No Number of Triangles Can Make A Circle with Positive Circumference!
 - ② If $b > 0$, Infinitely Many Triangles Make An Infinitely Large Circumference!
 - In Either Case, How Can We Ever Get A Finite Circle from Infinite Infinitely Small Pieces?
 - ① Every Real Number Must Be Archimedean!
 - ② Since No Infinitesimal Is Archimedean, They Were Rejected by Euclid, Aristotle, Archimedes.

Seriously?

Is That (**Argument 1** [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by **Nicholas of Cusa in The 15th Century!**
- We Wonder If It Is Even Logical! Clearly, **Objections Galore!**
 - The **Very Notion of A Triangle with An Infinitely Small Base Is Illusive!**
 - The Base of The Triangle, *b*, **However Small**, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, **No Number of Triangles Can Make A Circle with Positive Circumference!**
 - ② If $b > 0$, **Infinitely Many Triangles Make An Infinitely Large Circumference!**
 - In Either Case, How Can We Ever Get A Finite Circle from Infinite Infinitely Small Pieces?
 - ① Every Real Number Must Be Archimedean!
 - ② Since **No Infinitesimal Is Archimedean**, They Were Rejected by Euclid, Aristotle, Archimedes.

Seriously?

Is That (**Argument 1** [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by **Nicholas of Cusa in The 15th Century!**
- We Wonder If It Is Even Logical! Clearly, **Objections Galore!**
 - The **Very Notion of A Triangle with An Infinitely Small Base Is Illusive!**
 - The Base of The Triangle, b , **However Small**, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, No Number of Triangles Can Make A Circle with Positive Circumference!
 - ② If $b > 0$, Infinitely Many Triangles Make An Infinitely Large Circumference!
 - In Either Case, How Can We Ever Get A Finite Circle from Infinite Infinitely Small Pieces?
 - ① Every Real Number Must Be Archimedean!
 - ② Since No Infinitesimal Is Archimedean, They Were Rejected by Euclid, Aristotle, Archimedes.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Seriously?

Is That (Argument 1 [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by Nicholas of Cusa in The 15th Century!
- We Wonder If It Is Even Logical! Clearly, Objections Galore!
 - The Very Notion of A Triangle with An Infinitely Small Base Is Illusive!
 - The Base of The Triangle, b , However Small, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, No Number of Triangles Can Make A Circle with Positive Circumference!
 - ② If $b > 0$, Infinitely Many Triangles Make An Infinitely Large Circumference!
 - In Either Case, How Can We Ever Get A Finite Circle from Infinite Infinitely Small Pieces?
 - ① Every Real Number Must Be Archimedean!
 - ② Since No Infinitesimal Is Archimedean, They Were Rejected by Euclid, Aristotle, Archimedes.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Seriously?

Is That (**Argument 1** [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by **Nicholas of Cusa in The 15th Century!**
- We Wonder If It Is Even Logical! Clearly, **Objections Galore!**
 - The **Very Notion of A Triangle with An Infinitely Small Base Is Illusive!**
 - The Base of The Triangle, b , **However Small**, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, **No Number of Triangles Can Make A Circle with Positive Circumference!**
 - ② If $b > 0$, **Infinitely Many Triangles Make An Infinitely Large Circumference!**
 - In Either Case, How Can We Ever Get **A Finite Circle from Infinite Infinitely Small Pieces?**
 - ① Every Real Number Must Be Archimedean!
 - ② Since **No Infinitesimal Is Archimedean**, They Were Rejected by Euclid, Aristotle, Archimedes.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Seriously?

Is That (Argument 1 [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by Nicholas of Cusa in The 15th Century!
- We Wonder If It Is Even Logical! Clearly, Objections Galore!
 - The Very Notion of A Triangle with An Infinitely Small Base Is Illusive!
 - The Base of The Triangle, b , However Small, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, No Number of Triangles Can Make A Circle with Positive Circumference!
 - ② If $b > 0$, Infinitely Many Triangles Make An Infinitely Large Circumference!
 - In Either Case, How Can We Ever Get A Finite Circle from Infinite Infinitely Small Pieces?
 - ① Every Real Number Must Be Archimedean!
 - ② Since No Infinitesimal Is Archimedean, They Were Rejected by Euclid, Aristotle, Archimedes.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Seriously?

Is That (**Argument 1** [8]) A Valid Mathematical Proof?

- That *Argument* [8] Was Published by **Nicholas of Cusa in The 15th Century!**
- We Wonder If It Is Even Logical! Clearly, **Objections Galore!**
 - The **Very Notion of A Triangle with An Infinitely Small Base Is Illusive!**
 - The Base of The Triangle, b , **However Small**, Must Clearly Be ≥ 0 , Right?
 - ① If $b = 0$, **No Number of Triangles Can Make A Circle with Positive Circumference!**
 - ② If $b > 0$, **Infinitely Many Triangles Make An Infinitely Large Circumference!**
 - In Either Case, How Can We Ever Get **A Finite Circle from Infinite Infinitely Small Pieces?**
 - ① Every Real Number Must Be Archimedean!
 - ② Since **No Infinitesimal Is Archimedean**, They Were Rejected by Euclid, Aristotle, Archimedes.

Nonstandard
Analysis

MD,RH,KM

Introduc-
tion

Summary

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- Archimedes's Work Came in Two Streams of Tradition:
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His Method of Exhaustion.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since Infinitesimals Don't Exist³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to Infinitesimals in It.
 - ② Ironically, You May Find Its Logic Akin to Weierstrass's Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- **Archimedes's Work Came in Two Streams of Tradition:**
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His **Method of Exhaustion**.
 - Discovered in Constantinople in 1906!
 - Relies on An “**Indirect Argument**” And Purely Finite **Constructions**.
 - Finds The Volumes of **Surfaces of Revolution** (e.g., A Paraboloid).
 - Since **Infinitesimals Don't Exist**³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to **Infinitesimals** in It.
 - ② Ironically, You May Find Its Logic Akin to **Weierstrass's** Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- **Archimedes's Work Came in Two Streams of Tradition:**
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His **Method of Exhaustion**.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since **Infinitesimals Don't Exist**³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to **Infinitesimals** in It.
 - ② Ironically, You May Find Its Logic Akin to **Weierstrass's** Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- **Archimedes's Work Came in Two Streams of Tradition:**
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His **Method of Exhaustion**.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since **Infinitesimals Don't Exist**³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to **Infinitesimals** in It.
 - ② Ironically, You May Find Its Logic Akin to **Weierstrass's** Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Archimedes's Work Came in Two Streams of Tradition:
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His Method of Exhaustion.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since Infinitesimals Don't Exist³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to Infinitesimals in It.
 - ② Ironically, You May Find Its Logic Akin to Weierstrass's Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Archimedes's Work Came in Two Streams of Tradition:
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His Method of Exhaustion.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since Infinitesimals Don't Exist³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to Infinitesimals in It.
 - ② Ironically, You May Find Its Logic Akin to Weierstrass's Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Archimedes's Work Came in Two Streams of Tradition:
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His Method of Exhaustion.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since Infinitesimals Don't Exist³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to Infinitesimals in It.
 - ② Ironically, You May Find Its Logic Akin to Weierstrass's Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Archimedes's Work Came in Two Streams of Tradition:
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His Method of Exhaustion.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since Infinitesimals Don't Exist³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to Infinitesimals in It.
 - ② Ironically, You May Find Its Logic Akin to Weierstrass's Formulation.

³Or Do They?

Archimedes's Work: Lost And Found

Riddle of The Quadrature of The Parabola

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Archimedes's Work Came in Two Streams of Tradition:
1) Continuous, 2) After A Gap of 1000 Years.
- Of Present Interest Is His Method of Exhaustion.
 - Discovered in Constantinople in 1906!
 - Relies on An “Indirect Argument” And Purely Finite Constructions.
 - Finds The Volumes of Surfaces of Revolution (e.g., A Paraboloid).
 - Since Infinitesimals Don't Exist³, It Gives A Logically Acceptable, Rigorous Proof of His Results!
 - ① No References Are Made to Infinitesimals in It.
 - ② Ironically, You May Find Its Logic Akin to Weierstrass's Formulation.

³Or Do They?

The Area And Circumference of The Unit Circle

Argument 2: A *Logical, But Pedantic*, Argument Avoiding Infinitesimals?

Area of Unit Circle = $\frac{1}{2}$ Its Circumference (Flawless Logic?)

Let S Be The Proposition^a That The Area of A Unit Circle (\mathbb{A}_C) = Its Half-circumference (\mathbb{H}_C).

If S Is false, Then

$$\text{Either } \mathbb{A}_C > \mathbb{H}_C \quad \text{Or} \quad \mathbb{H}_C > \mathbb{A}_C \quad (1)$$

Let \mathbb{D} Be The *Positive Difference* between \mathbb{A}_C And \mathbb{H}_C .
Therefore, If $\mathbb{H}_C > \mathbb{A}_C$, Then

$$\mathbb{D} = \mathbb{H}_C - \mathbb{A}_C \quad (2)$$

Otherwise,

$$\mathbb{D} = \mathbb{A}_C - \mathbb{H}_C \quad (3)$$

^aTherefore, S Must Be Either true Or false.

The Area And Circumference of The Unit Circle

Argument 2: Continued ...

Area of Unit Circle = $\frac{1}{2}$ Its Circumference (Flawless Logic?)

...

Now We Use **Proof by Contradiction** to Show that Both Eq. [2] and Eq. [3] Lead to Contradiction with Eq. [1].

Let's start with Eq. [2] first.

We Can **Circumscribe @ The Circle A Regular Polygon with as Many Sides as We Wish.**

Since The Polygon Is Composed of Finite Number of Finite Triangles with Altitude = 1, **Area of The Polygon equals Its Half-perimeter.**

$$\mathbb{A}_P = \mathbb{H}_P \quad (4)$$

As The Number of Sides of The Polygon Circumscribing The Circle Increases, The Difference in Their Areas, $\mathbb{A}_P - \mathbb{A}_C$, Reduces.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

The Area And Circumference of The Unit Circle

Argument 2: Continued ...

Area of Unit Circle = $\frac{1}{2}$ Its Circumference (Flawless Logic?)

...

We Can Increase The Number of Sides of The Polygon Such That $\mathbb{A}_P - \mathbb{A}_C < \frac{\mathbb{D}}{2}$

Then, from Equations [2] and [4],

$$\begin{aligned}\mathbb{H}_P - (\mathbb{H}_C - \mathbb{D}) &< \frac{\mathbb{D}}{2} \\ \therefore \mathbb{H}_P - \mathbb{H}_C + \mathbb{D} &< \frac{\mathbb{D}}{2} \\ \therefore \mathbb{H}_P &< \mathbb{H}_C - \frac{\mathbb{D}}{2}\end{aligned}\tag{5}$$

This Is A Contradiction; The Polygon Circumscribes The Circle, How Can Its Perimeter Be Smaller Than That of The Circle?

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

The Area And Circumference of The Unit Circle

Argument 2: Continued ...

Area of Unit Circle = $\frac{1}{2}$ Its Circumference (Flawless Logic?)

...

We Reason Similarly for Eq. [3]: Increase The Number of Sides of The Polygon Such That $\mathbb{H}_P - \mathbb{H}_C < \frac{\mathbb{D}}{2}$ (Note: Half-Perimeters). Then, from Equations [3] and [4],

$$\begin{aligned}\mathbb{A}_P - (\mathbb{A}_C - \mathbb{D}) &< \frac{\mathbb{D}}{2} \\ \therefore \mathbb{A}_P - \mathbb{A}_C + \mathbb{D} &< \frac{\mathbb{D}}{2} \\ \therefore \mathbb{A}_P &< \mathbb{A}_C - \frac{\mathbb{D}}{2}\end{aligned}\tag{6}$$

A Contradiction Again; How Can The Area of A Circle Be Greater Than That of The Circumscribing Polygon?

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

The Area And Circumference of The Unit Circle

Argument 2 Ends

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

Area of Unit Circle = $\frac{1}{2}$ Its Circumference (Flawless Logic?)

...

Thus, When $\mathbb{A}_C \neq \mathbb{H}_C$, We Reach A Contradiction.
Therefore, $\mathbb{A}_C = \mathbb{H}_C$. □

Comparing The Two ‘Proofs’

One Direct And One Indirect

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

We Have Two Proofs That $\mathbb{A}_C = \mathbb{H}_C$: The **Direct Proof of Nicholas of Cusa [8] Using Infinitesimals** And The **Indirect Argument of Archimedes [15] Avoiding Infinitesimals!**

- The **Direct Proof Is Strange, The Indirect Proof Pedantic!**
- They Agree And Reflect the ‘Beliefs’ of Their Creators: Infinitesimals Made Archimedes Uncomfortable, But Inspired Nicholas of Cusa⁴.
- Nicholas of Cusa, Johannes Kepler, Blaise Pascal Reveled Mysticism of ∞ . Newton, Leibniz, Bernoulli Brothers, L’Hôpital, Euler Vowed to Unravel It.
 - Not Everyone Who Contributed to Formalizing Calculus Using Infinitesimals Believed Their Existence!
 - Newton Accepted **But Avoided Them in *Principia***, Leibniz Accepted **But Didn’t Claim Their *Existence*!**

⁴Who Was A Cardinal of The Church

Comparing The Two ‘Proofs’

One Direct And One Indirect

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

We Have Two Proofs That $\mathbb{A}_C = \mathbb{H}_C$: The **Direct Proof of Nicholas of Cusa [8] Using Infinitesimals** And The **Indirect Argument of Archimedes [15] Avoiding Infinitesimals!**

- The **Direct Proof Is Strange, The Indirect Proof Pedantic!**
- They Agree And Reflect the ‘Beliefs’ of Their Creators: Infinitesimals Made Archimedes Uncomfortable, But Inspired Nicholas of Cusa⁴.
- Nicholas of Cusa, Johannes Kepler, Blaise Pascal Reveled Mysticism of ∞ . Newton, Leibniz, Bernoulli Brothers, L’Hôpital, Euler Vowed to Unravel It.
 - Not Everyone Who Contributed to Formalizing Calculus Using Infinitesimals Believed Their Existence!
 - Newton Accepted **But Avoided Them in *Principia***, Leibniz Accepted **But Didn’t Claim Their *Existence*!**

⁴Who Was A Cardinal of The Church

Comparing The Two ‘Proofs’

One Direct And One Indirect

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

We Have Two Proofs That $\mathbb{A}_C = \mathbb{H}_C$: The **Direct Proof of Nicholas of Cusa [8] Using Infinitesimals** And The **Indirect Argument of Archimedes [15] Avoiding Infinitesimals!**

- The **Direct Proof Is Strange, The Indirect Proof Pedantic!**
- They Agree And Reflect the ‘Beliefs’ of Their Creators: Infinitesimals Made Archimedes Uncomfortable, But Inspired Nicholas of Cusa⁴.
- Nicholas of Cusa, Johannes Kepler, Blaise Pascal Reveled Mysticism of ∞ . Newton, Leibniz, Bernoulli Brothers, L'Hôpital, Euler Vowed to Unravel It.
 - Not Everyone Who Contributed to Formalizing Calculus Using Infinitesimals Believed Their Existence!
 - Newton Accepted But Avoided Them in *Principia*, Leibniz Accepted But Didn't Claim Their *Existence!*

⁴Who Was A Cardinal of The Church

Comparing The Two ‘Proofs’

One Direct And One Indirect

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

We Have Two Proofs That $\mathbb{A}_C = \mathbb{H}_C$: The **Direct Proof of Nicholas of Cusa [8] Using Infinitesimals** And The **Indirect Argument of Archimedes [15] Avoiding Infinitesimals!**

- The **Direct Proof Is Strange, The Indirect Proof Pedantic!**
- They Agree And Reflect the ‘Beliefs’ of Their Creators: Infinitesimals Made Archimedes Uncomfortable, But Inspired Nicholas of Cusa⁴.
- Nicholas of Cusa, Johannes Kepler, Blaise Pascal Reveled Mysticism of ∞ . Newton, Leibniz, Bernoulli Brothers, L’Hôpital, Euler Vowed to Unravel It.
 - Not Everyone Who Contributed to Formalizing Calculus Using Infinitesimals Believed Their Existence!
 - Newton Accepted But Avoided Them in *Principia*, Leibniz Accepted But Didn’t Claim Their *Existence!*

⁴Who Was A Cardinal of The Church

Comparing The Two ‘Proofs’

One Direct And One Indirect

We Have Two Proofs That $\mathbb{A}_C = \mathbb{H}_C$: The **Direct Proof of Nicholas of Cusa [8] Using Infinitesimals** And The **Indirect Argument of Archimedes [15] Avoiding Infinitesimals!**

- The **Direct Proof Is Strange, The Indirect Proof Pedantic!**
- They Agree And Reflect the ‘Beliefs’ of Their Creators: Infinitesimals Made Archimedes Uncomfortable, But Inspired Nicholas of Cusa⁴.
- Nicholas of Cusa, Johannes Kepler, Blaise Pascal Reveled Mysticism of ∞ . Newton, Leibniz, Bernoulli Brothers, L’Hôpital, Euler Vowed to Unravel It.
 - Not Everyone Who Contributed to Formalizing Calculus Using Infinitesimals Believed Their Existence!
 - Newton Accepted But Avoided Them in *Principia*, Leibniz Accepted But Didn’t Claim Their *Existence!*

⁴Who Was A Cardinal of The Church

Comparing The Two ‘Proofs’

One Direct And One Indirect

We Have Two Proofs That $\mathbb{A}_C = \mathbb{H}_C$: The **Direct Proof of Nicholas of Cusa [8] Using Infinitesimals** And The **Indirect Argument of Archimedes [15] Avoiding Infinitesimals!**

- The **Direct Proof Is Strange, The Indirect Proof Pedantic!**
- They Agree And Reflect the ‘Beliefs’ of Their Creators: Infinitesimals Made Archimedes Uncomfortable, But Inspired Nicholas of Cusa⁴.
- Nicholas of Cusa, Johannes Kepler, Blaise Pascal Reveled Mysticism of ∞ . Newton, Leibniz, Bernoulli Brothers, L’Hôpital, Euler Vowed to Unravel It.
 - Not Everyone Who Contributed to Formalizing Calculus Using Infinitesimals Believed Their Existence!
 - Newton Accepted **But Avoided Them in *Principia***, Leibniz Accepted **But Didn’t Claim Their *Existence*!**

⁴Who Was A Cardinal of The Church

Infinitesimals in Dynamics

From Geometry to Dynamics

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Since Antiquity, Geometry Played A Crucial Role in Providing Analysis Problems.

Dynamics Was a Relatively Recent Experience. All Kinds of **Moving Bodies** Started to Appear After Newton. Tools of Calculus Help Tremendously to Analyze **Motion**.

Motion Was, After All, Difficult to Ignore. Since The Time of Zeno⁵ of Elea And, His Teacher, Parmenides, We Were Confused about Motion. Zeno Asserted That The **Universe Was Static And All Motion Was Illusion**. He Postulated Several Paradoxes to Confound (Illuminate?) People. But After Newton, We Had to Study Motion Systematically, Which Would **Remove Zeno's Paradoxes about The Ubiquitous Motion** (If Not Erase His Philosophy) ...

Let's Analyze An Everyday Motion by The Two Approaches (**Standard: Weierstrass, Nonstandard: Robinson**).

⁵Around 450 BC in Ancient Greece.


Analysis of The Motion of A Falling Body

1: The Basic Setup

Consider a Falling Stone.

Its Motion Is Described by Giving Its Position (Displacement from A Reference) as A *Function* of Time. As It Falls, Its Velocity Increases, So That The Velocity at Each Instant Is Also A Variable *Function* of Time. Newton Called The *Instantaneous Position* Function The 'Fluent' And The *Instantaneous Velocity* Function the 'Fluxion'.

If One Is Given, The Other Can Be Determined; This Connection Is The Heart of The Infinitesimal Calculus Fashioned by Newton and Leibniz.



Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its **Instantaneous Position as A Function of Time**.
 - We Understand The Idea of The **Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel** as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend **How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?**
 - Determining This Velocity Is Challenging When We Realize that **The Body Constantly Moves Faster and Faster ...**

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its **Instantaneous Position as A Function of Time**.
 - We Understand The Idea of The **Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel** as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend **How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?**
 - Determining This Velocity Is Challenging When We Realize that **The Body Constantly Moves Faster and Faster ...**

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its *Instantaneous Position as A Function of Time*.
 - We Understand The Idea of The *Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel* as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend *How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?*
 - Determining This Velocity Is Challenging When We Realize that *The Body Constantly Moves Faster and Faster ...*

⁶Exact 'Time-keeping' Is As Fascinating As It Is Challenging.

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its *Instantaneous Position as A Function of Time*.
 - We Understand The Idea of The *Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel* as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend *How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?*
 - Determining This Velocity Is Challenging When We Realize that *The Body Constantly Moves Faster and Faster ...*

⁶Exact 'Time-keeping' Is As Fascinating As It Is Challenging.

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its **Instantaneous Position as A *Function* of Time**.
 - We Understand The Idea of The Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend **How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?**
 - Determining This Velocity Is Challenging When We Realize that **The Body Constantly Moves Faster and Faster ...**

⁶Exact 'Time-keeping' Is As Fascinating As It Is Challenging.

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its **Instantaneous Position as A Function of Time**.
 - We Understand The Idea of The **Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel** as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend *How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?*
 - Determining This Velocity Is Challenging When We Realize that The Body Constantly Moves Faster and Faster ...

⁶Exact 'Time-keeping' Is As Fascinating As It Is Challenging.

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated...
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its **Instantaneous Position as A Function of Time**.
 - We Understand The Idea of The **Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel** as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend **How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?**
 - Determining This Velocity Is Challenging When We Realize that **The Body Constantly Moves Faster and Faster ...**

Analysis of The Motion of A Falling Body

2: The Concept of *Instantaneous* Velocity

- We Take This Routine Experience of A Falling Body for Granted. However, Its Motion is Complicated. . .
 - First, We Must Be Able to Tell (Somehow) When An Exact Time Interval (e.g., 1 Second) Elapses⁶.
 - We Can Then Think of Measuring The Precise Distance Traveled by A Body Falling Freely (Under Gravity).
 - Then We Can Tell *Where in Air* The Body Was When $t = 1s, 2s, \dots$, i.e., We Can Describe Its **Instantaneous Position as A Function of Time**.
 - We Understand The Idea of The **Average Speed of A Moving/Moved Body over The Entire Duration of Its Travel** as $\frac{\text{Total Distance Traveled}}{\text{Total Time Elapsed since The Beginning}}$.
 - But Do We Really Comprehend **How Fast A Falling Body Was Moving at $t = 1s$, or $t = 2s$?**
 - Determining This Velocity Is Challenging When We Realize that **The Body Constantly Moves Faster and Faster . . .**

⁶Exact 'Time-keeping' Is As Fascinating As It Is Challenging.

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limit'ed Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limited Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limited Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limited Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limited Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limit'ed Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Analysis of The Motion of A Falling Body

3: Introducing The Infinitesimal Change

- To Determine The Instantaneous Velocity of A Body That Ever Moves Faster, Newton Made A Fair Assumption.
 - The Velocity Remains Constant During A Tiny (Infinitesimal) Period of Time.
 - The Continuous Change in Velocity Actually Comes in The Form of Tiny, Discrete Jumps.
 - Robinson Formalized This Change as A Number That Behaves Differently from Real Numbers.
- We'll First Study What Newton Proposed. Then We'll Go through Bishop Berkeley's Objections And Weierstrass's 'Limit'ed Remedy Resulting in Standard Analysis. Finally, We Will Encounter Robinson's Revival of Infinitesimals Resulting in Nonstandard Analysis!
 - This Will Compare And Contrast Robinson's Nonstandard Analysis with Weierstrass's Standard One.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So **Drop The Infinitesimal!**
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So Drop The Infinitesimal!
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So Drop The Infinitesimal!
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So Drop The Infinitesimal!
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So Drop The Infinitesimal!
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So Drop The Infinitesimal!
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So Drop The Infinitesimal!
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So **Drop The Infinitesimal!**
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Analysis of The Motion of A Falling Body

4: Newton's Insight, Leibniz's Notation, And Robinson's Formalization

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- An Infinitesimal Time Interval dt Later, Its Instantaneous Position, $s' = 4.9(1 + dt)^2$.
- Correspondingly, The Change in Instantaneous Position, $ds = s' - s = 4.9((1 + dt)^2 - 1)$.
- $\therefore ds = 9.8dt + 4.9dt^2$.
- The Instantaneous Velocity, v_1 , at Time $t = 1$ Is $\frac{ds}{dt} = \frac{9.8dt + 4.9dt^2}{dt} = 9.8 + 4.9dt$.
- Since dt Is Infinitesimal, So Is $4.9dt$. We Only Entertain Real Quantities, So **Drop The Infinitesimal!**
- Therefore, Instantaneous Velocity, v_1 , Is The Real Part of $\frac{ds}{dt} = 9.8$.

Analysis of The Motion of A Falling Body

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as Obscure, Repugnant(!), and Precarious(!!) as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 Is Unintelligible.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}$!
 - "May We Not Call Them (Those *Fluxions*) The Ghosts of Departed Quantities?"
 - Even Newton, Finally Acknowledging The Lack of Rigor in His Formulation, Could Not Provide The Necessary Rigor!
 - In Rebus Mathematicis errores quam minimi non sunt contemnendi⁹.

⁷Newton Was Aware of It.

⁸That Is, Derivatives.

⁹In Mathematics, The Minutest Errors Are Not to be Contemned.

Analysis of The Motion of A Falling Body

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as Obscure, Repugnant(!), and Precarious(!!) as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 Is Unintelligible.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}!$
 - "May We Not Call Them (Those *Fluxions*) The Ghosts of Departed Quantities?"
- Even Newton, Finally Acknowledging The Lack of Rigor in His Formulation, Could Not Provide The Necessary Rigor!
 - In Rebus Mathematicis errores quam minimi non sunt contemnendi⁹.

⁷ Newton Was Aware of It.

⁸ That Is, Derivatives.

⁹ In Mathematics, The Minutest Errors Are Not to be Contemned⁹.

Analysis of The Motion of A Falling Body

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as **Obscure, Repugnant(!), and Precarious(!!)** as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 **Is Unintelligible**.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}!$
 - "May We Not Call Them (Those *Fluxions*) **The Ghosts of Departed Quantities?**"
 - Even Newton, Finally Acknowledging **The Lack of Rigor in His Formulation**, Could Not Provide The Necessary Rigor!
 - *In Rebus Mathematicis errores quam minimi non sunt contemnendi*⁹.

⁷ Newton Was Aware of It.

⁸ That Is, Derivatives.

⁹ In Mathematics, The Minutest Errors Are Not to be Contemned.

Analysis of The Motion of A Falling Body

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as **Obscure, Repugnant(!), and Precarious(!)** as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 **Is Unintelligible**.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}!$
 - "May We Not Call Them (Those *Fluxions*) The Ghosts of Departed Quantities?"
 - Even Newton, Finally Acknowledging The Lack of Rigor in His Formulation, Could Not Provide The Necessary Rigor!
 - In Rebus Mathematicis errores quam minimi non sunt contemnendi⁹.

⁷ Newton Was Aware of It.

⁸ That Is, Derivatives.

⁹ In Mathematics, The Minutest Errors Are Not to

Analysis of The Motion of A Falling Body

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as Obscure, Repugnant(!), and Precarious(!!) as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 Is Unintelligible.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}!$
 - "May We Not Call Them (Those *Fluxions*) The Ghosts of Departed Quantities?"
- Even Newton, Finally Acknowledging The Lack of Rigor in His Formulation, Could Not Provide The Necessary Rigor!
 - In Rebus Mathematicis errores quam minimi non sunt contemnendi⁹.

⁷ Newton Was Aware of It.

⁸ That Is, Derivatives.

⁹ In Mathematics, The Minutest Errors Are Not to be Contemned.

Analysis of The Motion of A Falling Body

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as **Obscure, Repugnant(!), and Precarious(!!)** as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 **Is Unintelligible**.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}!$
 - “May We Not Call Them (Those *Fluxions*) **The Ghosts of Departed Quantities?**”
- Even Newton, Finally Acknowledging **The Lack of Rigor in His Formulation**, Could Not Provide The Necessary Rigor!

• In Rebus Mathematicis errores quam minimi non sunt contemnendi⁹.

⁷ Newton Was Aware of It.

⁸ That Is, Derivatives.

⁹ In Mathematics, The Minutest Errors Are Not to be Contemned⁹.


5: Bishop Berkeley's Critique of An Infidel Mathematician!

5: Bishop Berkeley's Critique of An Infidel Mathematician!

- Bishop Berkeley Wrote *The Analyst*, A Brilliant and Devastating Critique of Newton-Leibniz's Infinitesimals, in 1734⁷.
 - Newton's *Fluxions*⁸ Are as **Obscure, Repugnant(!), and Precarious(!!)** as Any Point in Divinity.
 - Leibniz's *Procedure* That $9.8 + 4.9dt$ Simply Equals 9.8 **Is Unintelligible**.
 - $dt \neq 0 \implies 9.8 + 4.9dt \neq 9.8$.
 - $dt = 0 \implies ds = 0 \implies \frac{ds}{dt} = \frac{0}{0}!$
 - “May We Not Call Them (Those *Fluxions*) **The Ghosts of Departed Quantities?**”
- Even Newton, Finally Acknowledging **The Lack of Rigor in His Formulation**, Could Not Provide The Necessary Rigor!
 - In Rebus Mathematicis errores quam minimi non sunt contemnendi⁹.

⁷Newton Was Aware of It.

⁸That Is, Derivatives.

⁹In Mathematics, The Minutest Errors Are Not to Be Ignored. 

Analysis of The Motion of A Falling Body

6: Setting The Stage for Weierstrass, The Rigorous Analyst of 19th Century

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Newton's Influence (And Inexact Infinitesimal Calculus's Undeniable Effectiveness) Overcame Berkeley's Criticism.
- For Almost Two Centuries, Naturalists Used “The Inexact” Infinitesimal Calculus to Solve Many Practical Problems in Physics.
- Ultimately, A Pure Mathematician like Weierstrass Led The Efforts to Reinstall Rigor in Analysis (Calculus) in The 19th Century.
- Like Our Greek Ancestors, Weierstrass Formally Outlawed Infinitesimals by Perfecting an Intriguing Idea . . .

Analysis of The Motion of A Falling Body

6: Setting The Stage for Weierstrass, The Rigorous Analyst of 19th Century

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Newton's Influence (And Inexact Infinitesimal Calculus's Undeniable Effectiveness) Overcame Berkeley's Criticism.
- For Almost Two Centuries, Naturalists Used "The Inexact" Infinitesimal Calculus to Solve Many Practical Problems in Physics.
- Ultimately, A Pure Mathematician like Weierstrass Led The Efforts to Reinstall Rigor in Analysis (Calculus) in The 19th Century.
- Like Our Greek Ancestors, Weierstrass Formally Outlawed Infinitesimals by Perfecting an Intriguing Idea . . .

Analysis of The Motion of A Falling Body

6: Setting The Stage for Weierstrass, The Rigorous Analyst of 19th Century

- Newton's Influence (And Inexact Infinitesimal Calculus's Undeniable Effectiveness) Overcame Berkeley's Criticism.
- For Almost Two Centuries, Naturalists Used “The Inexact” Infinitesimal Calculus to Solve Many Practical Problems in Physics.
- Ultimately, A Pure Mathematician like Weierstrass Led The Efforts to Reinstate Rigor in Analysis (Calculus) in The 19th Century.
- Like Our Greek Ancestors, Weierstrass Formally Outlawed Infinitesimals by Perfecting an Intriguing Idea . . .

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

6: Setting The Stage for Weierstrass, The Rigorous Analyst of 19th Century

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Newton's Influence (And Inexact Infinitesimal Calculus's Undeniable Effectiveness) Overcame Berkeley's Criticism.
- For Almost Two Centuries, Naturalists Used “The Inexact” Infinitesimal Calculus to Solve Many Practical Problems in Physics.
- Ultimately, A Pure Mathematician like Weierstrass Led The Efforts to Reinstall Rigor in Analysis (Calculus) in The 19th Century.
- Like Our Greek Ancestors, Weierstrass Formally Outlawed Infinitesimals by Perfecting an Intriguing Idea . . .

Analysis of The Motion of A Falling Body

6: Setting The Stage for Weierstrass, The Rigorous Analyst of 19th Century

- Newton's Influence (And Inexact Infinitesimal Calculus's Undeniable Effectiveness) Overcame Berkeley's Criticism.
- For Almost Two Centuries, Naturalists Used “The Inexact” Infinitesimal Calculus to Solve Many Practical Problems in Physics.
- Ultimately, A Pure Mathematician like Weierstrass Led The Efforts to Reinstall Rigor in Analysis (Calculus) in The 19th Century.
- Like Our Greek Ancestors, Weierstrass Formally Outlawed Infinitesimals by Perfecting an Intriguing Idea ...

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A Finite Time Interval¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A *Limit* Reached by It.

¹⁰Only Familiar Real Numbers; No dt, ds Business. ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

Analysis of The Motion of A Falling Body

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A Finite Time Interval¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A *Limit* Reached by It.

¹⁰Only Familiar Real Numbers; No dt, ds Business. ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

Analysis of The Motion of A Falling Body

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A Finite Time Interval¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A *Limit* Reached by It.

¹⁰Only Familiar Real Numbers; No dt, ds Business. ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

Analysis of The Motion of A Falling Body

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time:
 $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A **Finite Time Interval**¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A *Limit* Reached by It.

¹⁰Only Familiar Real Numbers; No dt, ds Business. ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ ↻ ↻

Analysis of The Motion of A Falling Body

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time:
 $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A **Finite Time Interval**¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A *Limit* Reached by It.

¹⁰Only Familiar Real Numbers; No dt , ds Business. ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ≡ ≡ ≡

Analysis of The Motion of A Falling Body

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time:
 $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A **Finite Time Interval**¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A *Limit* Reached by It.

¹⁰Only Familiar Real Numbers; No dt , ds Business. ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ≡ ≡ ≡

7: Weierstrass's Rigorous Formalization. Look Ma! No Infinitesimals!

Introduction

- Like Before, The Instantaneous Position of A Falling Body (in Meters) is Given by A Function of Time: $s = 4.9t^2$ (Where Time t Is Measured in Seconds).
- Therefore, at $t = 1$, $s = 4.9$.
- A **Finite Time Interval**¹⁰ Δt Later, Its Instantaneous Position, $s' = 4.9(1 + \Delta t)^2$.
- Correspondingly, The Change in Instantaneous Position, $\Delta s = s' - s = 4.9((1 + \Delta t)^2 - 1)$.
- $\therefore \Delta s = 9.8\Delta t + 4.9(\Delta t)^2 \implies \frac{\Delta s}{\Delta t} = 9.8 + 4.9\Delta t$.
- In a Stark Contrast with Newton, We Define Instantaneous Velocity, Not as A Ratio of Distance and Time ($\frac{\Delta s}{\Delta t}$), But as A **Limit** Reached by It.

¹⁰ Only Familiar Real Numbers; No dt, ds Business.

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A **Positive Real Number**, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To **Choose Another Positive Real Number**, δ . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That **For Any Positive Value of Δt That Is Less Than δ** , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ **Must Be Less Than $\frac{\epsilon}{4.9}$** , That Is, ϵ .
- IOW, He Asserts That as δ *Approaches* 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ **Approaches No Other Number But 9.8**: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A Positive Real Number, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To Choose Another Positive Real Number, δ . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That For Any Positive Value of Δt That Is Less Than δ , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ Must Be Less Than $4.9\frac{\epsilon}{4.9}$, That Is, ϵ .
- IOW, He Asserts That as δ Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ Approaches No Other Number But 9.8: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A Positive Real Number, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To Choose Another Positive Real Number, δ . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That For Any Positive Value of Δt That Is Less Than δ , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ Must Be Less Than $4.9\frac{\epsilon}{4.9}$, That Is, ϵ .
- IOW, He Asserts That as δ Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ Approaches No Other Number But 9.8: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A **Positive Real Number**, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To Choose Another Positive Real Number, δ . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That For Any Positive Value of Δt That Is Less Than δ , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ Must Be Less Than $4.9\frac{\epsilon}{4.9}$, That Is, ϵ .
- IOW, He Asserts That as δ Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ Approaches No Other Number But 9.8: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A **Positive Real Number**, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To **Choose Another Positive Real Number**, δ . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That For Any Positive Value of Δt That Is Less Than δ , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ Must Be Less Than $4.9\frac{\epsilon}{4.9}$, That Is, ϵ .
- IOW, He Asserts That as δ Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ Approaches No Other Number But 9.8: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A **Positive Real Number**, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To **Choose Another Positive Real Number**, δ . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That **For Any Positive Value of Δt That Is Less Than δ** , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ **Must Be Less Than $\frac{\epsilon}{4.9}$** , That Is, ϵ .
- IOW, He Asserts That as δ Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ Approaches No Other Number But 9.8: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Analysis of The Motion of A Falling Body

8: *Limits* Achieve Limitless Rigor!

- Weierstrass Illustrates The Idea of A *Limit* Before Defining It Precisely ...
- He Tries to Show That However Close to Zero Δt Goes, $\frac{\Delta s}{\Delta t} = 9.8 + 4.9(\Delta t)$ May Go Closer to A Real Number. If He Succeeds, His Rigor Holds Strong!
- First, We Give Weierstrass A **Positive Real Number**, However Small, ϵ . Thus, $\epsilon > 0$.
- Weierstrass Is Free To **Choose Another Positive Real Number, δ** . In This Case, He Chooses $\delta = \frac{\epsilon}{4.9}$; $\delta > 0$.
- Then, Weierstrass Asserts That **For Any Positive Value of Δt That Is Less Than δ** , $(\frac{\Delta s}{\Delta t} - 9.8)$ Which Equals $4.9\Delta t$ **Must Be Less Than $\frac{\epsilon}{4.9}$** , That Is, ϵ .
- IOW, He Asserts That as δ Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0, i.e., $\frac{\Delta s}{\Delta t}$ **Approaches No Other Number But 9.8**: $\lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = 9.8$.

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
 - Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
 - Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
 - Weierstrass Succeeds!
 - Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
 - Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
 - Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches Exactly 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Analysis of The Motion of A Falling Body

9: An Illustration of Weierstrass's Limit

- We Challenge Weierstrass to Prove That $(\frac{\Delta s}{\Delta t} - 9.8) = 4.9\Delta t$ Is Less Than Some Positive Number ϵ We Pick, for Every Positive Number Δt That Is Less Than Some Positive Number δ He Picks.
- Let's Give Him $\epsilon = 0.00049$.
- Weierstrass Readily Picks $\delta = \frac{\epsilon}{4.9} = 0.0001$.
- Clearly, for Every Positive Δt Less Than $\delta = \frac{\epsilon}{4.9}$ (i.e., $0 < \Delta t < \frac{\epsilon}{4.9}$), $0 < 4.9\Delta t < \epsilon$.
- Weierstrass Succeeds!
- Therefore, as Δt Approaches 0, $(\frac{\Delta s}{\Delta t} - 9.8)$ Also Approaches 0.
- Therefore, as Δt Approaches 0, $\frac{\Delta s}{\Delta t}$ Approaches *Exactly* 9.8. The Instantaneous Velocity is $9.8m/s$ at $t = 1s$.
- Weierstrass Could Seamlessly Find the Exact Limits of Many Other Functions.

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of *Limits*.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹ Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of *Limits*.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹ Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of *Limits*.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹ Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of *Limits*.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹ Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of *Limits*.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹ Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of Limits.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹ Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

- On A First Reading, This Mere Replacement of dt , ds by Δt , Δs Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ , δ , A Surprisingly Subtle Notion of Limits.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹Power of Proper Training?

Analysis of The Motion of A Falling Body

10: Is All That Rigor Worth It?

Nonstandard
Analysis

MD,RH,KM

Introduction

Summary

- On A First Reading, This Mere Replacement of dt, ds by $\Delta t, \Delta s$ Feels Strange.
- Granted, Weierstrass Achieved Two Important Things:
 - Removal of All Non-finite Quantities; Sticking to Reals.
 - Avoiding Division by Zero; He Never Sets Δt in $\frac{\Delta s}{\Delta t}$ to 0.
- Addressed The Bishop's Concerns, But at What Price?
 - An Everyday, Intuitive (If Paradoxical) Quantity, Instantaneous Velocity, Is Subjected to Unrelated Real Numbers, ϵ, δ , A Surprisingly Subtle Notion of Limits.
 - Not Knowing Limits Didn't Keep The Bernoullis, Euler, ... From Finding Velocities in Complicated Cases.
 - We Knew The Answers Using Infinitesimals That Matched Answers Found By Limits. And Yet, For "Logical Consistency", We Embraced a Subtle Definition That Mathematicians Appreciate¹¹ Much More Than Common People. Did We Treat Infinitesimals Fairly?

¹¹Power of Proper Training?

Analysis of The Motion of A Falling Body

11: And, Finally, The Rigorous Limit Is Defined!

Here's That Rigorous Definition of **The Limit of A Function** in Its Modern Grandeur:

Definition (Limit of A Real-Valued Function)

Consider The Function, $f : \mathbb{R} \rightarrow \mathbb{R}$, And Two Real Numbers p, L . We Say, **The Limit of f of x , as x Approaches p , Exists And Equals L** , And Write, $\lim_{x \rightarrow p} f(x) = L$, If The Following

Property Holds:

For Every Real $\epsilon > 0$, There Exists A Real $\delta > 0$, Such That For All Real x , $0 < |x - p| < \delta$ Implies $|f(x) - L| < \epsilon$.

It Is More Cryptic When Expressed Symbolically:

$$(\forall \epsilon > 0)(\exists \delta > 0)(\forall x \in \mathbb{R})(0 < |x - p| < \delta) \implies |f(x) - L| < \epsilon$$

Summary of Part 1

Introducing Nonstandard Analysis

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary

References

Nonstan-
dard
Analysis

MD,RH,KM

Introduc-
tion

Summary



[Goldbach, 1742] Christian Goldbach.

A problem we should try to solve before the ISPN 43
deadline,

Letter to Leonhard Euler, 1742.