

# MATH 417 502

## Homework 2

Keegan Smith

September 4, 2024

### Problem 1

a.) The first 20 iterations of the Newton method are shown below:

Iteration	Approximation	Residual
0	1.000000e+00	7.182818e-01
1	5.819767e-01	2.075957e-01
2	3.190550e-01	5.677201e-02
3	1.679962e-01	1.493591e-02
4	8.634887e-02	3.837726e-03
5	4.379570e-02	9.731870e-04
6	2.205769e-02	2.450693e-04
7	1.106939e-02	6.149235e-05
8	5.544905e-03	1.540144e-05
9	2.775014e-03	3.853917e-06
10	1.388149e-03	9.639248e-07
11	6.942351e-04	2.410369e-07
12	3.471577e-04	6.026621e-08
13	1.735889e-04	1.506742e-08
14	8.679696e-05	3.766965e-09
15	4.339911e-05	9.417547e-10
16	2.169971e-05	2.354406e-10
17	1.084989e-05	5.886025e-11
18	5.424958e-06	1.471512e-11
19	2.712481e-06	3.678613e-12
20	1.356302e-06	9.199308e-13

From the above, we can see that Newton's method probably converges, but not as quickly as we would like.

Requirements to converge quadratically:

$$g'(x^*) = 0$$

$$g''(x) \leq M$$

In our case, from newton's method:

$$g(x) = x - \frac{e^x - x - 1}{e^x - 1}$$

Thus we have:

$$\begin{aligned} g'(x) &= 1 - ((-e^x)(e^x - 1)^{-2}(e^x - x - 1) + (e^x - 1)^{-1}(e^x - 1)) \\ g'(0) &= 1 - ((-1)(1 - 1)^{-2}(1 - 0 - 1) + \frac{0}{0}) \end{aligned}$$

$\frac{0}{0}$  tells us nothing, NOTHING about a function. Thus we will take the limit as  $x$  approaches 0:

$$\begin{aligned} \lim_{x \rightarrow 0} (1 - ((-e^x)(e^x - 1)^{-2}(e^x - x - 1) + (e^x - 1)^{-1}(e^x - 1))) &= \lim_{x \rightarrow 0} (1 - (\frac{(-e^{2x} + xe^x + e^x)}{(e^x - 1)^2} + 1)) \\ \lim_{x \rightarrow 0} (1 - (\frac{(-e^{2x} + xe^x + e^x)}{(e^x - 1)^2} + 1)) &= \lim_{x \rightarrow 0} (-\frac{(-e^{2x} + xe^x + e^x)}{(e^x - 1)^2}) \end{aligned}$$

Using L'Hopital's:

$$\lim_{x \rightarrow 0} (-\frac{(-e^{2x} + xe^x + e^x)}{(e^x - 1)^2}) = \lim_{x \rightarrow 0} (-\frac{(-2e^{2x} + e^x + xe^x + e^x)}{2e^x(e^x - 1)})$$

This is still  $\frac{0}{0}$ .....use L'Hopital's again.

$$\lim_{x \rightarrow 0} (-\frac{(-2e^{2x} + 2e^x + xe^x)}{2e^x(e^x - 1)}) = \lim_{x \rightarrow 0} (-\frac{(-4e^{2x} + 2e^x + e^x + xe^x)}{2e^x(e^x - 1) + 2e^{2x}})$$

This finally gives us  $\frac{1}{2}$ , so we can safely say that since  $g'(0) < 1$ ,  $g$  converges, but since  $g'(0) \neq 0$ ,  $g$  does not converge quadratically.

b.) Start with  $\mu(x)$  and  $\mu'(x)$ :

$$\mu(x) = \frac{f(x)}{f'(x)}$$

$$\begin{aligned} \mu'(x) &= \frac{d}{dx} (f(x)(f'(x))^{-1}) \\ &= \frac{f'(x)}{f'(x)} + (-1) \cdot f''(x) \cdot f'(x)^{-2} \cdot f(x) \\ &= \frac{f'(x)}{f'(x)} - \frac{f''(x) \cdot f(x)}{f'(x)^2} \end{aligned}$$

Plugging into  $g(x)$ :

$$\begin{aligned}
 g(x) &= x - \frac{\mu(x)}{\mu'(x)} \\
 &= x - \frac{f(x)}{f'(x) \cdot \left( \frac{f'(x)}{f'(x)} - \frac{f''(x) \cdot f(x)}{f'(x)^2} \right)} \\
 &= x - \frac{f(x)}{\frac{f'(x)^2}{f'(x)} - \frac{f''(x) \cdot f(x)}{f'(x)}} \\
 &= x - \frac{f(x)}{f'(x)^{-1}(f'(x)^2 - f''(x) \cdot f(x))} \\
 &= x - \frac{f(x) \cdot f'(x)}{f'(x)^2 - f''(x) \cdot f(x)}
 \end{aligned}$$

c.) We can see that the new  $g(x)$  is approximately correct within the first 5 iterations:

Iteration	Approximation	Residual
0	1.000000e+00	7.182818e-01
1	-2.342106e-01	2.540578e-02
2	-8.458280e-03	3.567061e-05
3	-1.189018e-05	7.068790e-11
4	-4.218591e-11	0.000000e+00
5	-4.218591e-11	0.000000e+00

From the results above, it would appear that the new  $g$  converges quadratically. I think I'm supposed to verify this with theory however, so I will attempt to do so:

$$\begin{aligned}
 g'(x) &= 1 - \frac{(f'(x)^2 + f(x)f''(x))(f'(x)^2 - f''(x)f(x))}{(f'(x)^2 - f''(x)f(x))^2} \\
 &\quad + \frac{f(x)f'(x)(2f'(x)f''(x) - (f(x)f'''(x) + f'(x)f''(x)))}{(f'(x)^2 - f''(x)f(x))^2} \\
 g'(0) &= 1 - \frac{0}{0}
 \end{aligned}$$

Using L'Hopital's:

$$\lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)f''(x))(f'(x)^2 - f''(x)f(x))}{(f'(x)^2 - f''(x)f(x))^2} \right)$$

$$\begin{aligned}
&= \lim_{x \rightarrow 0} \left( \frac{(2f''(x)f'(x) + f'(x)f''(x) + f(x)f'''(x))(f'(x)^2 - f''(x)f(x))}{(2f''(x)f'(x) - f'''(x)f(x) - f''(x)f'(x))(2)(f'(x)^2 - f''(x)f(x))} \right. \\
&\quad \left. + \frac{(f'(x)^2 + f(x)f''(x))(2f''(x)f'(x) - f'''(x)f(x) - f''(x)f'(x))}{(2f''(x)f'(x) - f'''(x)f(x) - f''(x)f'(x))(2)(f'(x)^2 - f''(x)f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(2e^x f'(x) + f'(x)e^x + f(x)e^x)(f'(x)^2 - e^x f(x))}{(2e^x f'(x) - e^x f(x) - e^x f'(x))(2)(f'(x)^2 - e^x f(x))} \right. \\
&\quad \left. + \frac{(f'(x)^2 + f(x)e^x)(2e^x f'(x) - e^x f(x) - e^x f'(x))}{(2e^x f'(x) - e^x f(x) - e^x f'(x))(2)(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{e^x(2f'(x) + f'(x) + f(x))(f'(x)^2 - e^x f(x))}{e^x(2f'(x) - f(x) - f'(x))(2)(f'(x)^2 - e^x f(x))} \right. \\
&\quad \left. + \frac{(f'(x)^2 + f(x)e^x)e^x(2f'(x) - f(x) - f'(x))}{e^x(2f'(x) - f(x) - f'(x))(2)(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(2f'(x) + f'(x) + f(x))}{(2f'(x) - f(x) - f'(x))(2)} \right. \\
&\quad \left. + \frac{(f'(x)^2 + f(x)e^x)}{(2)(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{3f'(x) + f(x)}{(f'(x) - f(x))(2)} \right. \\
&\quad \left. + \frac{(f'(x)^2 + f(x)e^x)}{(2)(f'(x)^2 - e^x f(x))} \right) \\
&= \frac{0}{0}
\end{aligned}$$

"When you try so hard, but it doesn't even matter". L'Hopital's again...

$$\lim_{x \rightarrow 0} \left( \frac{3f'(x) + f(x)}{(f'(x) - f(x))(2)} + \frac{(f'(x)^2 + f(x)e^x)}{(2)(f'(x)^2 - e^x f(x))} \right)$$

$$\begin{aligned}
&= \lim_{x \rightarrow 0} \left( \frac{3f''(x) + f'(x)}{(f''(x) - f'(x))(2)} + \frac{(2f''(x)f'(x) + f'(x)e^x + f(x)e^x)}{(2)(2f''(x)f'(x) - e^x f(x) - f'(x)e^x)} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{3e^x + f'(x)}{(e^x - f'(x))(2)} + \frac{(2e^x f'(x) + f'(x)e^x + f(x)e^x)}{(2)(2e^x f'(x) - e^x f(x) - f'(x)e^x)} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{3e^x + f'(x)}{(e^x - f'(x))(2)} + \frac{e^x(2f'(x) + f'(x) + f(x))}{(2e^x)(2f'(x) - f(x) - f'(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{3e^x + f'(x)}{(e^x - f'(x))(2)} + \frac{(2f'(x) + f'(x) + f(x))}{(2)(2f'(x) - f(x) - f'(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{3e^x + f'(x)}{(e^x - f'(x))(2)} + \frac{(3f'(x) + f(x))}{(2)(f'(x) - f(x))} \right) \\
&= \frac{3}{2} + \lim_{x \rightarrow 0} \left( \frac{(3f'(x) + f(x))}{(2)(f'(x) - f(x))} \right) \\
&= \frac{3}{2} + \lim_{x \rightarrow 0} \left( \frac{(3f''(x) + f'(x))}{(2)(f''(x) - f'(x))} \right) \\
&= \frac{3}{2} + \frac{3}{2} = 3
\end{aligned}$$

Yay, now for the other term in the original expression:

$$\begin{aligned}
&\lim_{x \rightarrow 0} \left( \frac{f(x)f'(x)(2f'(x)f''(x) - (f(x)f'''(x) + f'(x)f''(x)))}{(f'(x)^2 - f''(x)f(x))^2} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{f(x)f'(x)(2f'(x)e^x - f(x)e^x - f'(x)e^x)}{(f'(x)^2 - e^x f(x))^2} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{f(x)f'(x)e^x x}{(f'(x)^2 - e^x f(x))^2} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)f''(x))e^x x + f(x)f'(x)(e^x x + e^x)}{2(2f''(x)f'(x) - e^x f(x) - e^x f'(x))(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)e^x x + e^x f(x)f'(x)(x + 1)}{2(2e^x f'(x) - e^x f(x) - e^x f'(x))(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{e^x((f'(x)^2 + f(x)e^x)x + f(x)f'(x)(x + 1))}{2e^x(2f'(x) - f(x) - f'(x))(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)x + f(x)f'(x)(x + 1)}{2(f'(x) - f(x))(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)x + f(x)f'(x)(x + 1)}{2x(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)x}{2x(f'(x)^2 - e^x f(x))} + \frac{f(x)f'(x)(x + 1)}{2x(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)}{2(f'(x)^2 - e^x f(x))} + \frac{f(x)f'(x)(x + 1)}{2x(f'(x)^2 - e^x f(x))} \right)
\end{aligned}$$

$$\begin{aligned}
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)}{2(f'(x)^2 - e^x f(x))} + \frac{f(x)f'(x)x}{2x(f'(x)^2 - e^x f(x))} + \frac{f(x)f'(x)}{2x(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(f'(x)^2 + f(x)e^x)}{2(f'(x)^2 - e^x f(x))} + \frac{f(x)f'(x)}{2(f'(x)^2 - e^x f(x))} + \frac{f(x)f'(x)}{2x(f'(x)^2 - e^x f(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{2f''(x)f'(x) + f'(x)e^x + f(x)e^x}{2(2f''(x)f'(x) - e^x f(x) - e^x f'(x))} + \frac{f'(x)^2 + f(x)f''(x)}{2(2f''(x)f'(x) - e^x f(x) - e^x f'(x))} \right. \\
&\quad \left. + \frac{f'(x)^2 + f(x)f''(x)}{2(f'(x)^2 - e^x f(x)) + 2x(2f''(x)f'(x) - e^x f(x) - e^x f'(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{2e^x f'(x) + f'(x)e^x + f(x)e^x}{2(2e^x f'(x) - e^x f(x) - e^x f'(x))} + \frac{f'(x)^2 + f(x)e^x}{2(2e^x f'(x) - e^x f(x) - e^x f'(x))} \right. \\
&\quad \left. + \frac{f'(x)^2 + f(x)e^x}{2(f'(x)^2 - e^x f(x)) + 2x(2e^x f'(x) - e^x f(x) - e^x f'(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{e^x(2f'(x) + f'(x) + f(x))}{2e^x(2f'(x) - f(x) - f'(x))} + \frac{f'(x)^2 + f(x)e^x}{2e^x(2f'(x) - f(x) - f'(x))} \right. \\
&\quad \left. + \frac{f'(x)^2 + f(x)e^x}{2(f'(x)^2 - e^x f(x)) + 2xe^x(2f'(x) - f(x) - f'(x))} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(3f'(x) + f(x))}{2(x)} + \frac{f'(x)^2 + f(x)e^x}{2e^x(x)} \right. \\
&\quad \left. + \frac{f'(x)^2 + f(x)e^x}{2(f'(x)^2 - e^x f(x)) + 2xe^x(x)} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{(3f''(x) + f'(x))}{2} + \frac{2f''(x)f'(x) + f'(x)e^x + f(x)e^x}{2e^x(x) + 2e^x} \right. \\
&\quad \left. + \frac{2f''(x)f'(x) + f'(x)e^x + f(x)e^x}{2(2f''(x)f'(x) - e^x f(x) - e^x f'(x)) + 2(2xe^x + x^2e^x)} \right) \\
&= \frac{3}{2} + \lim_{x \rightarrow 0} \left( \frac{e^x(2f'(x) + f'(x) + f(x))}{e^x(2(x) + 2)} \right. \\
&\quad \left. + \frac{2e^x f'(x) + f'(x)e^x + f(x)e^x}{2(2e^x f'(x) - e^x f(x) - e^x f'(x)) + 2(2xe^x + x^2e^x)} \right) \\
&= \frac{3}{2} + \lim_{x \rightarrow 0} \left( \frac{(3f'(x) + f(x))}{(2(x) + 2)} \right. \\
&\quad \left. + \frac{e^x(2f'(x) + f'(x) + f(x))}{2e^x(2f'(x) - f(x) - f'(x)) + 2e^x(2x + x^2)} \right)
\end{aligned}$$

$$\begin{aligned}
 &= \frac{3}{2} + \lim_{x \rightarrow 0} \left( \frac{(3f'(x) + f(x))}{2(f'(x) - f(x)) + 2e^x(2x + x^2)} \right) \\
 &= \frac{3}{2} + \lim_{x \rightarrow 0} \left( \frac{(3f''(x) + f'(x))}{2(f''(x) - f'(x)) + 2e^x(2x + x^2) + 2e^x(2 + 2x)} \right) \\
 &= \frac{3}{2} + \frac{3}{6} \\
 &= 2
 \end{aligned}$$

Thus in total we have:

$$g'(0) = 1 - 3 + 2 = 0$$

So, indeed,  $g$  converges quadratically.

## Problem 2

1. We know that if  $|g'(x^*)| < 1$  then  $g(x)$  converges. So for  $a$  we have:

$$\begin{aligned}
 g(x) &= \frac{1}{21}(20x + 21x^{-2}) \\
 g'(x) &= \frac{1}{21}(20 - 42x^{-3}) \\
 g'(21^{\frac{1}{3}}) &= \frac{1}{21}(20 - 42(21^{\frac{1}{3}})^{-3}) \\
 &= \frac{1}{21}(20 - 42(21^{-1})) \\
 &= \frac{1}{21}(20 - 2) \\
 &= \frac{1}{21}(18) \\
 &= \frac{18}{21} < 1
 \end{aligned}$$

Thus  $a$  likely converges, but does not converge quadratically since  $g'(x) \neq 0$ .

for  $b$  we have:

$$\begin{aligned}
 g(x) &= x - \frac{1}{3}(x^3 - 21)x^{-2}, \\
 g'(x) &= 1 - \frac{1}{3}((3x^2)x^{-2} + (x^3 - 21)(-2)x^{-3}), \\
 g'(x) &= 1 - \frac{1}{3}(3 + (x^3 - 21)(-2)x^{-3}), \\
 g'(21^{\frac{1}{3}}) &= 1 - \frac{1}{3}(3 + (21 - 21)(-2)(21^{-1})) \\
 &= 1 - \frac{1}{3}(3) \\
 &= 1 - 1 \\
 &= 0
 \end{aligned}$$

so  $b$  converges, and  $b$  also likely quadratically converges since  $g'(x*) \equiv 0$   
For  $c$  we have:

$$\begin{aligned}
 g(x) &= x - (x^4 - 21x)(x^2 - 21)^{-1} \\
 g'(x) &= 1 - ((4x^3 - 21)(x^2 - 21)^{-1} + (x^4 - 21x)(-1)(2x)(x^2 - 21)^{-2}) \\
 g'(21^{\frac{1}{3}}) &= 1 - (4(21 - 21)(21^{\frac{2}{3}} - 21)^{-1} - (21^{\frac{4}{3}} - 21^{\frac{4}{3}})(2(21^{\frac{1}{3}}))(21^{\frac{2}{3}} - 21)^{-2}) \\
 &= 1 - (0 - 0) \\
 &= 1
 \end{aligned}$$

$g'(x*) \geq 1$  so  $c$  does not converge linearly or quadratically.

2. Table for  $a$  is below:



Iteration	Approximation	Error
0	1.000000e+00	1.758924e+00
1	1.952381e+00	8.065432e-01
2	2.121754e+00	6.371699e-01
3	2.242850e+00	5.160745e-01
4	2.334840e+00	4.240845e-01
5	2.407093e+00	3.518308e-01
6	2.465059e+00	2.938649e-01
7	2.512243e+00	2.466807e-01
8	2.551057e+00	2.078671e-01
9	2.583238e+00	1.756864e-01
10	2.610081e+00	1.488427e-01
11	2.632580e+00	1.263439e-01
12	2.651510e+00	1.074147e-01
13	2.667484e+00	9.143969e-02
14	2.681000e+00	7.792397e-02
15	2.692459e+00	6.646529e-02

Table for  $b$  is below:

Iteration	Approximation	Error
0	1.000000e+00	1.758924e+00
1	7.666667e+00	4.907742e+00
2	5.230204e+00	2.471280e+00
3	3.742697e+00	9.837727e-01
4	2.994854e+00	2.359294e-01
5	2.777022e+00	1.809805e-02
6	2.759042e+00	1.176900e-04
7	2.758924e+00	5.020131e-09
8	2.758924e+00	4.440892e-16
9	2.758924e+00	4.440892e-16
10	2.758924e+00	4.440892e-16
11	2.758924e+00	4.440892e-16
12	2.758924e+00	4.440892e-16
13	2.758924e+00	4.440892e-16
14	2.758924e+00	4.440892e-16
15	2.758924e+00	4.440892e-16

Table for  $c$  is below:

Iteration	Approximation	Error
0	1.000000e+00	1.758924e+00
1	0.000000e+00	2.758924e+00
2	0.000000e+00	2.758924e+00
3	0.000000e+00	2.758924e+00
4	0.000000e+00	2.758924e+00
5	0.000000e+00	2.758924e+00
6	0.000000e+00	2.758924e+00
7	0.000000e+00	2.758924e+00
8	0.000000e+00	2.758924e+00
9	0.000000e+00	2.758924e+00
10	0.000000e+00	2.758924e+00
11	0.000000e+00	2.758924e+00
12	0.000000e+00	2.758924e+00
13	0.000000e+00	2.758924e+00
14	0.000000e+00	2.758924e+00
15	0.000000e+00	2.758924e+00

We can see from the above that our hypotheses do hold since  $a$  does converge, but not quickly.  $b$  converges and does so much more quickly than  $a$ , and  $c$  does not converge to  $21^{\frac{1}{3}}$ . The python code for this is below:

```

1 import math;
2 NUM_ITER = 16
3 EXPECTED = 21**(1/3)
4 def f(x):
5     return math.e**x - x - 1;
6 def f_prime(x):
7     return math.e**x - 1;
8 def f_double_prime(x):
9     return math.e**x;
10 def g1(x):
11     return x - f(x) / f_prime(x);
12 def g2(x):
13     return x - (f(x)*f_prime(x)) / (f_prime(x)**2 - f(x)
14         * f_double_prime(x))
15 def a(x):
16     return (20 * x + 21 / (x**2)) / 21
17 def b(x):
18     return x - (x**3 - 21)/(3 * x**2)
19 def c(x):
20     return x - (x**4 - 21 * x) / (x**2 - 21)
21 def compute(initial_x, function, f_function = None):
22     result = "\\begin{tabular}{|c|c|c|c|}\n"
23     result += "\\hline\n"
24     result += "Iteration & Approximation & Error\\\\\n"
25     result += "\\hline\n"
26     curr_approx = initial_x

```

```
26     error = abs(curr_approx - EXPECTED)
27     for i in range(0, NUM_ITER):
28         result += f"{i} & {curr_approx:.6e} & {error:.6e}
29             \\\\\n"
30         result += "\\hline\n"
31         curr_approx = function(curr_approx)
32         error = abs(curr_approx - EXPECTED)
33         result += "\\end{tabular}\n"
34
35     return result
36     #return approx;
37 function_mapping = {
38     "f" : f,
39     "g1": g1,
40     "g2": g2,
41     "a" : a,
42     "b" : b,
43     "c" : c,
44 }
45 if __name__ == "__main__":
46     initial_num = float(input("enter initial num: "))
47     my_func = input("enter the name of the function: ")
48     #my_func_2 = input("enter the name of the function (
49         f): ")
50     result = compute(initial_num, function_mapping[
51         my_func])
52     print(result)
```