

# (sub A2) Cell wall (CW)

(intro to A2.2 and A2.3)


### Purpose of the lecture:

- What are the general functions of the CW?
- What is the Gram stain?
- How do the structures of gram-positive and gram-negative cell walls differ?





# The Gram Stain



(a) Gram-Positive

(b) Gram-Negative


Differential stain, most NB in microbiology


Staining is related to or reflects differences in?

# A2.2 Gram-positive cell wall

- Thick homogeneous CW
- Primarily peptidoglycan (PG)

\_





Gustav Bouwer - 2018

#### **Techoic acids**

- CW contains large amounts
- Polymers

Connected to?

\_

Extend to surface

What are lipotechoic acids?

### Techoic acid functions


Maintain structure of cell wall

Negative charge

# A2.3 Gram-negative cell wall

- More complex
- Thin PG layer


\_\_\_\_



#### • OM contains:

\_\_\_


\_\_\_



## Braun's lipoprotein

- What does it do?

- How?



Gustav Bouwer - 2018

## Lipopolysaccharides (LPS)

Composition?


OM is asymmetric

\_\_\_

\_\_\_

• e.g. Salmonella typhimurium

#### **Structure**



#### **LPS functions**

- Lipid A is major component of OM, helps stability
- Protect cells permeability barrier
- Core polysaccharide
- O side chain
- LPS can acts as endotoxin

# **2.3.3.1 OM functions**

(component functions)

Prevent loss of constituents

Important protective barrier

#### Porins

Transmembrane proteins

- Channels for entrance/exit

### 2.3.2 Periplasmic space

Occupied by periplasm

Many proteins that participate in nutrient acquisition

ET and Energy

Chemoreceptors (chemotaxis)

Enzymes for PG synthesis (also Gram+)

Modification?

- Gram+?
  - Have what instead?