

Genetics

Course outline

- 1) extension of Mendelian genetics
- 2) Chromosomes structure and changes
- 3) Vertebrate development

Interactions between Genes, Alleles, and Phenotypes

Incomplete dominance

Genetic/ molecular cause

The recessive allele does not produce any functioning protein. The dominant allele does function protein but not in high enough quantities to result in the same phenotypic effect as a heterozygous dominant

phenotypic effect

A heterozygous individual will display a phenotype intermediate to that of homozygous dominant and homozygous recessive

Dominance

Genetic/molecular cause

The recessive allele does not produce any functioning protein. The dominant allele does function protein, which it produces in comparative amount to a homozygous dominant individual

Phenotypic consequence.

the heterozygous phenotype is identical to that of the heterozygous dominant individual.

Co-dominance

Genetic/ molecular cause

Both alleles are transcribed/translated to form a functioning protein

phenotypic effect

The effect comes from the full expression of both alleles

Dominance hierarchies.

It is often the case that there will be a number of alleles for one gene. In which case a dominance hierarchy is usually present such that allele A is Dominant to B is dominant to C and so forth (ie dominance in a hierarchy is transitive)

Pleiotropy

Genetic/molecular cause One gene locus affects the expression of several characteristics. Phenotypic cause Phenotype of one characteristic will change in conjunction with the phenotype of another characteristic

Notes about colour

As many genetic examples are based on colour it is worth understanding the basics of colouring in animals. colour results from pigments which are proteins, individual genes can either code directly for the production of these proteins or as an intermediate/enzyme which aids in the production of these proteins.

White

white does not result from a specific pigment but rather from a lack of pigment.

Co-dominance and Incomplete dominance

a co-dominant colour combination can look like an intermediate if it is the combination of two pigments, however in the case where one of the phenotypes is white then a intermediate can only be a result of incomplete dominance

Sex linked traits

Sex limited traits,

Important examples

BG 2032

13/02/2108

Phenylketonuria

single mutation can affect brain development however if the correct diet is followed then all detrimental health consequences can be ignored.

diabetes type II

lifestyle disease, the genetic component is estimated to change the likelihood of developing the disease by 2%, but the main causative factors of the disease relate to diet.

Lactose intolerance.

Individuals with lactose intolerance cant digest lactose, however the bacteria in their gut van which leads to the build up of gas. this can be very painful for the person in question. In humans lactose intolerance would naturally occur after weening however as cow milk is a staple part of most human diabetes the genes controlling lactase production are never switched off and the ability to digest lactose is not normally lost.

Himalayan Phenotype.

Cancer

basics

A cancer is a group of cells whose proliferation is uncontrolled, and which can spread to other locations in the body which are normally populated bu other cell types.

benigin tumours

grow but do not spread.

malignant tumours

over proliferate and invade other body tissues/ areas in the body.

types of cancers

Carcinomas

derived from epithelial cells

Sarcomas

derived from connective or muscle tissues. Known as Osteosarcoma in the case of bone tissue and Kaposi sarcoma in the case of soft connective tissue.

Lymphoma and leukaemias

1. Cancers of the hematopoietic system.
2. Lymphoma (solid tumour)
3. Leukaemia excess of circulating immature blood cell precursors. #####
Cancer of the nervous system brain and central nervous system.

NOTE: most common in children

Germ-line cancer.

Cancers of breast, prostate, lung, pancreas, and colon.

Causes of cancer.

In most cases cancer is not inherited. In breast cancer/ovarian cancer BRCA1, BRCA2 play some role in genetic disposition, perhaps 5-10%

hereditary.

Cancer normally sets in during old age as cells must accumulate a series of a specific set of mutations, (exactly which set of mutations accumulated lead to cancer is highly individual)

Environmental factors

1. UV
2. X rays
3. Alcohol
4. Overcooked food (heterocyclic amines, polycyclic aromatic hydrocarbons- meat. acrylamide- potatoes)
5. Azo dyes
6. tartrazine (food colourants)
7. Nitrate cured foods
8. Pesticides

Cancer initiators

agents which cause DNA damage (mutagens). these factors may be chemical, biological (such as HPV- human papilloma virus cause latent genetic damage- Cancer predisposition.

Cancer Promoters

Promotes excessive proliferation (does not directly damage DNA).

Examples

1. wounding
2. phorbol esters
3. HRT/ oestrogen (breast cancer)
4. hepatitis B (promotes stomach cancer)
5. HIV (Kaposi sarcoma, this disease is always present but only manifests itself in immunocompromised individuals)

stages of cancer development

1. normal cells (mutation occurs)
2. increased ability to proliferate (mutation)
3. 1-2mm tumour of rapidly proliferating cells which do not undergo apoptosis of differentiation. (mutation)
4. vascularized growing tumour (mutation)
5. large tumour capable of invading near by tissue (metastasis)

NOTE: cancer results from a series of somatic mutations, affecting the same cell, and different cancers are genetically heterogeneous.

Genetic instability in cancers.

deficient local DNA repair leads to the accumulation of point mutations increased chromosomal instability and gross genome abnormalities.

summary of key properties

1. disregards ex/in growth regulation signals.
2. avoid apoptosis, differentiation and replicative senescence
3. genetically unstable.
4. invasive
5. metastatic (survive and proliferate in foreign) sites.

Penetrance

percentage of the population who demonstrate at least some degree of phenotypic expression.

Expressivity

reflects the range of expressions of the gene/allele present in the population.

Genetic complementation

two different mutations in heterozygous condition affecting the same protein/pathway can complement each other to cause a novel phenotypic effect.

examples

rare form of albinism: both mother and father were normally pigmented and there families had minimal instances of albinism. however both children were albino's. Analysis found that this albinism stemmed not from those genes and mutation most commonly associated with albinism but rather from a combination of recessives (heterozygous) from both father and mother (mutations in TYRP1 gene), which on their own would have little or no effect on the pigment production pathway, but when combined reduced pigment production significantly.

NOTE: Rufus albinism leads to a phenotype with reddish hair, lighter skin, and blue grey eyes.

Forked line probability method.

##chromosome level (revise mitosis and meiosis) copy slides chromosome basic structure #### chromosome groups

meta-centric

centromere is half way up the length of the chromosome

submetacentric

centromere is more to the one side of the chromosome than the other,

Acrocentric

the centromere is very far to the one side of the chromosome, with a long arm containing most of the genes and a short arm containing predominantly telomeric DNA.

Telocentric

humans do not possess any telocentric chromosomes, but certain insect and crustacean species do.

Holocentric

Centromere like structure exist along the entire length of the chromosome. this may decrease the chances of faulty division/ segregation.

chromosome banding

bands were named and used to locate specific genes. the banding patterns are due to uneven DNA densities in the coiled structure of the chromosome.

Size and shape of different chromosomes.

Chromosome level mutations

Aneuploidy (Spelling)

Each cell has at least one extra chromosome or is missing at least one chromosome

Trisomy

when an individual inherits three copies of a particular chromosome. In Humans only three of all the possible trisomies are viable, (as in individuals with these mutations will still be born and not terminated during pregnancy)

of these three (Trisomy 13, 18, and 21) only individuals with trisomy 21 can survive past the first few years of childhood. Individuals with Trisomy 21 have down syndrome.

Down syndrome (Trisomy 21)

Frequency: 1/1000 Effects 1. mental retardation 2. short stature 3. heart disease 4. shortened life span.

Edward Syndrome (Trisomy 18)

Effects: 1. kidney and intestinal malformation 3. heart defects. 4. mental retardation

NOTE: only 8% survival past the first year.

Patau syndrome (Trisomy 13)

Effects: 1. Kidney and hear defects 2. polidactily (too many digits) 3. Nervous system abnormalities 4. Death within the 1st year.

NOTE: the smaller relative size of chromosome 21, meaning that it contains less important genes may be related to its increased viability .

Monosomy

when an individual inherits only one copy of a particular chromosome.

Nullisomy

having no copies of a particular chromosome.

Polyploidy

Structural rearrangements

1. deletion
2. duplication
3. translocation
4. inversion
5. fission
6. fusion

non disjuncture

(review mitosis and meiosis) If non disjuncture happens in meiosis it is generally worse than if it happens in mitosis as it will affect the whole organism.

Down Syndrome

occurs in about 1 of a thousand live births. originates from non disjunction in the egg cell rather than the sperm cell because the arrested development of the egg cells leads to decay of the separating and marking proteins as well as the spindle fibre leaving the whole process open to more error. 8% of individuals survive for one year ?

turner Syndrome monosomy X

Frequency 1/20 000 (only occurs in females) Effects: 1. webbed neck 2. short stature 3. underdeveloped ovaries and lack of secondary sex characteristics.

Klinefelter syndrome XXY

Too many X's ,XXY , XXXY, XXXXY, XXXXXY. (XXYY) Effects: 1. Decrease in testosterone levels 2. the more X's the higher the chance of brain damage.

XXY

XXY has the phenotype of a normal male. men with 2 Y chromosomes tend to be tall as there is a cumulative effect adding to height. (they may also exhibit increased aggression but this is unclear)

XXX

Frequency: 1/1000 Effects: 1. Normal female with normal fertility.

NOTE: this condition is seldom diagnosed.

Polyploidy

Three or more complete sets of chromosomes present in somatic cells. ####
2n-20n caused by non reduced gamete formation, which is very rare in males and non viable.

most often flowering plants are polyploid, they are even specifically bred to have more chromosome sets as this usually increases fruit and flower size.

Examples

water melon must be bred from a tetraploid(?) and a diploid to get a triploid infertile plant.

Kiwi fruit 12-16 copies

Strawberries 4 copies (tetraploid)

Frogs *Xenopus levis* 4n *Xenopus tropicalis* 2n

NOTE: plants are better adapted to polyploid because: 1. they are not as confined to a set physical form so different in growth and development genes operation levels are not so important 2. they can reproduce vegetatively so polyploid individuals aren't as severely evolutionarily disadvantaged. 3. less precise sperm targeting is necessary.

Deletions

deletions often occur by chromosome breakage during the cell cycle. These deletions may be terminal or interstitial. If the centromere is lost, the entire chromosome will be lost.

Piece of Chrome

sometimes multiple genes are lost when both strands break and a part of the chromosome is permanently lost. A specific deletion on chromosome 5. (where the entire p arm is deleted leads to a serious syndrome)

Cri du chat (5p-)

Frequency: 1/20000-50000 Effects: 1. Intellectual disability, 2. low muscle tone 3. microcephaly 4. distinctive facial features. 5. normal life expectancy

Duplications

Cause: retro transposition, or non-allelic homologous recombination in repeat rich areas.

A small region of a chromosome is repeated, and placed next to the existing copy (in tandem). duplications tend to happen in high repetitive sequences of DNA. These repeated regions mean that the chromosome will not line up very well with its homologous pair.

passed on by unequal order, and may or may not have a phenotypic effect.

Huntington Porea

Repeats alter gene function, which can lead to brain damage. Repeats may have a positive effect because they allow one copy of a gene to evolve independently, specialise in a different function while the original function is still conserved by the other copy of the gene.

Inversions.

Cause: non-allelic homologous recombination Effects: 1. pairing in meiosis occurs via looping 2. cross over in the looped region can cause major deletions or duplications.

Translocation

Effects: 1. novel chromosomes are generated 2. individuals are usually infertile. 3. can result in aneuploidy (such as 14q21q inherited down syndrome) 4.

terminal

A piece breaks off a chromosome and sticks to the end of a non-homologous chromosome.

reciprocal

Exchange of pieces between non-homologous chromosomes ##### Robertsonian
two Acrocentric chromosomes lose their short arms and get stuck together.

Fission

deletion (loss of acentric piece)'

Fusion

bicentric chromosomes (breakage in mitosis)

breakage fusion bridge cycles in cancer.

Copy image. 1. end of a chromosome breaks off (during meiosis) 2. one daughter cell inherits the chromosome lacking a telomere. 3. the new cell enters S phase and replicates its DNA (it now has two sticky DNA ends right next to each other.) 4. loose DNA strands bind to form continuous DNA loop. 5. fused sister chromatids are pulled apart during mitosis, causing breakage at a new sight 6. one daughter cell inherits chromosome with duplicated genes and again a free sticky end. this cycle can then begin again leading to even further chromosome distortion.

Imprinting

Hinny vs mule

Mule

horse mother donkey father hardy and obedient

Hinny

donkey mother, horse father. temperamental, untrainable.

Intersistial deletion of chromosome 15

Prader -Willi syndrome

Cause: deletion from father, ZNF127 and IPW inactivated on the maternal copy.
effects: 1. mental retardation 2. obesity 3. diminished growth.

Angelman syndrome

1. deletion from mother: UBE3A inactivated on paternal copy.

2. mental retardation
3. epilepsy
4. lack of motor development.

X inactivation

the inactivation of one of two X chromosomes in females via DNA methylation.

NOTE: which X chromosome is inactivated varies randomly from cell to cell. this inactivation process turns the inactive X into a bar body, (highly methylated and rolled up in histone proteins).

Frog colour

details

the skin colour of a certain from is controlled by one allele. There are three different phenotypes observed, these are red, blue, and purple.

explanation

This is an example of co-dominance as the purple colour comes from a combination of the red and blue pigments. that is both red and blue pigments have been produced. We can conclude that the purple from is heterozygous, the blue frog is homozygous blue and the red frog is homozygous red.

Fruit flies eye colour.

details.

the colour of fruit flies eyes is determined by two gene loci. a fly with dominant alleles at the red locus will be red eyed, while a fly with dominant alleles at the brown locus will be brown eyed. However a fly with both dominant brown and cinnabar will be white eyed.

explanation

the white eyed phenotype is due to a form of inhibition similar to epistasis stops either of the pigments from being produced if both pigment genes are present.

Rabbit Skin Colour

details

the skin colour of rabbits is determined by one gene locus, which has four possible Alleles. All alleles code for the enzyme tyrosinase which acts to convert tyrosine into dopamine a critical first set in the production of hair pigment.

#1 non functional

The most recessive Allele codes for a nonfunctioning version of tyrosinase leading to albino rabbit.

#2 heat sensitive

The next most recessive allele codes for a heat inactivated version of tyrosine meaning that the final colour of the rabbits fur will depend on the temperature conditions it was raised in, and also that within a certain temperature range there will be a distinct colour difference between the warmer and colder parts of the animal.

#3 localised functionality

The next most recessive allele codes for a version of tyrosinase does not fold quite “correctly” and as such does not glycosylate in the same way as the dominant version. melanocytes which produce the grey pigment will still accept this tyrosinase into their cytoplasm and as such grey pigment will be produced. however as intracellular transport is affected the tyrosinase either cannot reach or cannot enter melanocytes capable of producing the brown pigments and so they rabbits are full grey

#4 fully functional

the most dominant allele codes for a fully functional form of tyrosinase and the combination of grey and orange pigments produced gives the rabbits a full brown coat.

human Blood types.

details

human blood types are determined by one gene locus with three alleles. allele O is recessive, and codes for no functioning protein. allele A and B are co dominant and each code for different sugars which are attached to the outside of red blood cells.

Bombay phenotype

details

parents A and AB, child O, mates with A, results B, AB and O.

explanation

A and B sugars are attached to red blood cells by appending them onto the end of glycoproteins called Substance H which are already attached to the cell. If an individual is homozygous recessive for substance H genes, none will be produced and so even if they have A or B alleles they will have a functionally O phenotype.

Alleles and genes A gene is a collection/ group of alleles which could occur at the same locus and which are associated with the same traits.

lethal alleles alleles which are incompatible with the organism's continued survival

dominance

most lethal alleles are recessive because dominant lethal alleles produce non-functioning proteins only so any individual with even one of the dominant allele would be unlikely to survive to reproductive age, and so the Allele would very quickly be bred out of the population.

sex chromosomes

lethal alleles, or alleles linked with low fertility located on the Y chromosome in humans are very rare. disease related to homozygous XX are more likely to be suffered by woman, as they will have one working copy of the given gene so they can survive but one defective copy which leads to the illness. males on the other hand will either survive and be entirely healthy with a working copy or die, if they are even born if they have only a defective copy.

lethal dominants

lethal dominants may occur in one of two cases. First, if the onset of the disease/ defect related to the Allele only occurs after reproductive age. Secondly if a De Novo (random, once off) DNA mutation occurs in a given individual. lethal dominants can also result from an Allele which interferes with the formation of aggregates of the protein which the gene codes for, or receptors of that protein, or finally if it has a detrimental effect on another protein or gene (such as the yellow mice)

examples

Yellow Mice.

details

Y allele codes for the yellow coloured pigment. y codes for other pigment/no pigment. Y is a lethal Dominant(?) Allele. As a YY individual with not develop past the early stages of implantation into the uterine wall.

molecular/genetic explanation

When the Y allele is present at the gene loci it causes a deletion in which, its own gene promoter, as well as the coding sequence of the gene upstream of it (the MERC gene necessary in RNA processing) are removed. the net result being that proteins relating to the gene upstream are no longer removed, and the pigment gene is now promoter by the MERC promoter instead of its own promoter

Result and Phenotype

the result is that Yy individual still produce enough RNA processing proteins to survive/ avoid miscarriage, but they are more susceptible to diseases such as obesity and cancer. Also their yellow pigment gene is promoted heavily during early development leading to yellow coloured fur. YY miscarriages early in pregnancy as it can't produce necessary RNA processing proteins. yy is healthy individual without yellow colour. birth ratios will be skewed by the fact that YY individuals are not actually born. NOTE: this is an example of incomplete dominance of Y with respect to MERC.

Effective Of the environment on genetic expression.

examples

Myopia (short sightedness):

Populations where book learning is not enforced/encouraged from an early age do not show high rates of myopia even though the genetic disposition between the two populations is very similar.

Phenylketonuria

single mutation can affect brain development however if the correct diet is followed then all detrimental health consequences can be ignored.

Diabetes type II

lifestyle disease, the genetic component is estimated to change the likelihood of developing the disease by 2%, but the main causative factors of the disease

relate to diet.

Lactose intolerance.

Individuals with lactose intolerance cant digest lactose, however the bacteria in their gut can which leads to the build up of gas. this can be very painful for the person in question. In humans lactose intolerance would naturally occur after weaning however as cow milk is a staple part of most human diet the genes controlling lactase production are never switched off and the ability to digest lactose is not normally lost.

Himalayan Phenotype.

Cancer

basics

A cancer is a group of cells whose proliferation is uncontrolled, and which can spread to other locations in the body which are normally populated by other cell types.

benign tumours

grow but do not spread.

malignant tumours

over proliferate and invade other body tissues/ areas in the body.

types of cancers

Carcinomas

derived from epithelial cells

Sarcomas

derived from connective or muscle tissues. Known as Osteosarcoma in the case of bone tissue and Kaposi sarcoma in the case of soft connective tissue.

Lymphoma and leukaemias

1. Cancers of the hematopoietic system.
2. Lymphoma (solid tumour)
3. Leukaemia excess of circulating immature blood cell precursors. #####
Cancer of the nervous system brain and central nervous system.

NOTE: most common in children

Germ-line cancer.

Cancers of breast, prostate, lung, pancreas, and colon.

Causes of cancer.

In most cases cancer is not inherited. In breast cancer/ovarian cancer BS1, BS2 play some role in genetic disposition, perhaps 5-10%

Hereditary.

Cancer normally sets in during old age as cells must accumulate a series of a specific set of notations, (exactly which set of mutations accumulated lead to cancer is highly individual) ##### Environmental factors 1. UV 2. X rays 3. Alcohol 4. Overcooked food (heterocyclic amines, polycyclic aromatic hydrocarbons-meat. acrylamide- potatoes 5. Azo dyes 6. tartrazine (food colourants) 7. Nitrate cured foods 8. Pesticides

Cancer initiators

agents which cause DNA damage (mutagens). these factors may be chemical, biological (such as HPV- human papilloma virus cause latent genetic damage- Cancer predisposition.

Cancer Promoters

Promotes excessive proliferation (does not directly damage DNA).

Examples

1. wounding
2. phorbol esters
3. HRT/ oestrogen (breast cancer)
4. hepatitis B (promotes stomach cancer)

5. HIV (Kaposi sarcoma, this disease is always present but only manifests itself in immunocompromised individuals)

stages of cancer development

1. normal cells (mutation occurs)
2. increased ability to proliferate (mutation)
3. 1-2mm tumour of rapidly proliferating cells which do not undergo apoptosis of differentiation. (mutation)
4. vascularized growing tumour (mutation)
5. large tumour capable of invading near by tissue (metastasis)

NOTE: cancer results from a series of somatic mutations, affecting the same cell, and different cancers are genetically heterogeneous.

Genetic instability in cancers.

deficient local DNA repair leads to the accumulation of point mutations increased chromosomal instability and gross genome abnormalities.

summary of key properties

1. disregards ex/in growth regulation signals.
2. avoid apoptosis, differentiation and replicative senescence
3. genetically unstable.
4. invasive
5. metastatic (survive and proliferate in foreign) sites.

Penetrance

percentage of the population who demonstrate at least some degree of phenotypic expression.

Expressivity

reflects the range of expressions of the gene/allele present in the population.

Genetic complementation

two different mutations in heterozygous condition affecting the same protein/pathway can complement each other to cause a novel phenotypic effect.

examples

rare form of albinism: both mother and father were normally pigmented and their families had minimal instances of albinism. however both children were albino's. Analysis found that this albinism stemmed not from those genes and mutation most commonly associated with albinism but rather from a combination of recessives (heterozygous) from both father and mother (mutations in TYRP1 gene), which on their own would have little or no effect on the pigment production pathway, but when combined reduced pigment production significantly.

NOTE: Rufus albinism leads to a phenotype with reddish hair, lighter skin, and blue grey eyes.

Forked line probability method.

##Chromosome level (revise mitosis and meiosis) copy slides chromosome basic structure #### chromosome groups

meta-centric

centromere is half way up the length of the chromosome

submetacentric

centromere is more to the one side of the chromosome than the other,

Acrocentric

the centromere is very far to the one side of the chromosome, with a long arm containing most of the genes and a short arm containing predominantly telomeric DNA.

Telocentric

humans do not possess any telocentric chromosomes, but certain insect or crustacean species do.

Holocentric

Centromere like structure exist along the entire length of the chromosome. this may decrease the chances of faulty division/ segregation.

chromosome banding

bands were named and used to locate specific genes. the banding patterns are due to uneven DNA densities in the coiled structure of the chromosome.

Size and shape of different chromosomes.

Chromosome level mutations

Aneuploidy (Spelling)

Each cell is has at least one extra chromosome or is missing at least one chromosome

Trisomy

when an individual inherits three copies of a particular chromosome. In Humans only three of all the possible trisomies are viable, (as in individuals with these mutations will still be born and not terminated during pregnancy)

of these three (Trisomy 13, 18, and 21) only individuals with trisomy 21 can survive past the first few years of childhood. Individuals with Trisomy 21 have down syndrome.

Down syndrome (Trisomy 21)

Frequency: 1/1000 Effects 1. mental retardation 2. short stature 3. heart disease 4. shortened life span.

Edward Syndrome (Trisomy 18)

Effects: 1. kidney and intestinal malformation 3. heart defects. 4. mental retardation

NOTE: only 8% survival past the first year.

Patua syndrome (Trisomy 13)

Effects: 1. Kidney and hear defects 2. polidactily (too many digits) 3. Nervous system abnormalities 4. Death within the 1st year.

NOTE: the smaller relative size of chromosome 21, meaning that it contains less important genes may be related to its increased viability .

Monosomy

when an individual inherits only one copy of a particular chromosome.

Nullisomy

having no copies of a particular chromosome.

Polyploidy

Structural rearrangements

1. deletion
2. duplication
3. translocation
4. inversion
5. fission
6. fusion

non disjuncture

(review mitosis and meiosis) If non disjuncture happens in meiosis it is generally worse than if it happens in mitosis as it will affect the whole organism.

Down Syndrome

occurs in about 1 of a thousand live births. originates from non disjunction in the egg cell rather than the sperm cell because the arrested development of the egg cells leads to decay of the separating and marking proteins as well as the spindle fibre leaving the whole process open to more error. 8% of individuals survive for one year ?

turner Syndrome monosomy X

Frequency 1/20 000 (only occurs in females) Effects: 1. webbed neck 2. short stature 3. underdeveloped ovaries and lack of secondary sex characteristics.

Klinefelter syndrome XXY

Too many X's ,XXY , XXXY, XXXXY, XXXXXY. (XXYY) Effects: 1. Decrease in testosterone levels 2. the more X's the higher the chance of brain damage.

XYY

XYY has the phenotype of a normal male. men with 2 Y chromosomes tend to be tall as there is a cumulative effect adding to height. (they may also exhibit increased aggression but this is unclear)

XXX

Frequency: 1/1000 Effects: 1. Normal female with normal fertility.

NOTE: this condition is seldom diagnosed.

Polyploidy

Three or more complete sets of chromosomes present in somatic cells. ####
 $2n-20n$ caused by non reduced gamete formation, which is very rare in males and non viable.

most often flowering plants are polyploid, they are even specifically bred to have more chromosome sets as this usually increases fruit and flower size.

Examples

water melon must be bred from a tetraploid(?) and a diploid to get a triploid infertile plant.

Kiwi fruit 12-16 copies

Strawberries 4 copies (tetraploid)

Frogs *Xenopus levis* $4n$ *Xenopus tropicalis* $2n$

NOTE: plants are better adapted to polyploid because: 1. they are not as confined to a set physical form so different in growth and development genes operation levels are not so important 2. they can reproduce vegetatively so polyploid individuals aren't as severely evolutionarily disadvantaged. 3. less precise sperm targeting is necessary.

Deletions

deletions often occur by chromosome breakage during the cell cycle. These deletions may be terminal or interstitial. If the centromere is lost, the entire chromosome will be lost.

Piece of Chrome

sometimes multiple genes are lost when both strands break and a part of the chromosome is permanently lost. A specific deletion on chromosome 5. (where the entire p arm is deleted leads to a serious syndrome)

Cri du chat (5p-)

Frequency: 1/20000-50000 Effects: 1. Intellectual disability, 2. low muscle tone 3. microcephaly 4. distinctive facial features. 5. normal life expectancy

Duplications

Cause: retro transposition, or non-allelic homologous recombination in repeat rich areas.

A small region of a chromosome is repeated, and placed next to the existing copy (in tandem). duplications tend to happen in high repetitive sequences of DNA. These repeated regions mean that the chromosome will not line up very well with its homologous pair.

passed on by unequal order, and may or may not have a phenotypic effect.

Huntington Porea

Repeats alter gene function, which can lead to brain damage. Repeats may have a positive effect because they allow one copy of a gene to evolve independently, specialise in a different function while the original function is still conserved by the other copy of the gene.

Inversions.

Cause: non-allelic homologous recombination Effects: 1. pairing in meiosis occurs via looping 2. cross over in the looped region can cause major deletions or duplications.

Translocation

Effects: 1. novel chromosomes are generated 2. individuals are usually infertile. 3. can result in aneuploidy (such as 14q21q inherited down syndrome) 4.

terminal

A piece breaks off a chromosome and sticks to the end of a non-homologous chromosome.

reciprocal

Exchange of pieces between non-homologous chromosomes #### Robertsonian two Acrocentric chromosomes lose their short arms and get stuck together.

Fission

deletion (loss of acentric piece)'

Fusion

bicentric chromosomes (breakage in mitosis)

breakage fusion bridge cycles in cancer.

Copy image. 1. end of a chromosome breaks off (during meiosis) 2. one daughter cell inherits the chromosome lacking a telomere. 3. the new cell enters S phase and replicates its DNA (it now has two sticky DNA ends right next to each other.) 4. loose DNA strands bind to form continuous DNA loop. 5. fused sister chromatids are pulled apart during mitosis, causing breakage at a new sight 6. one daughter cell inherits chromosome with duplicated genes and again a free sticky end. this cycle can then begin again leading to even further chromosome distortion.

Imprinting

Hinny vs mule

Mule

horse mother donkey father hardy and obedient

Hinny

donkey mother, horse father. temperamental, untrainable.

Intersistial deletion of chromosome 15

Prader -Willi syndrome

Cause: deletion from father, ZNF127 and IPW inactivated on the maternal copy.
effects: 1. mental retardation 2. obesity 3. diminished growth.

Angelman syndrome

1. deletion from mother: UBE3A inactivated on paternal copy.
2. mental retardation
3. epilepsy
4. lack of motor development.

X inactivation

the inactivation of one of two X chromosomes in females via DNA methylation.

NOTE: which X chromosome is inactivated varies randomly from cell to cell.this inactivation process turns the inactive X into a bar body, (highly methylated and rolled up in histone proteins).