

Inflammatory Stress Contributes to Tau Pathology in the PS19 Mouse Model of Alzheimer's Disease

Keene, J.1-3, Desjarlais, T.1, Jackman, B.1, Srinageshwar, B.2-4, Finneran, D. 1, Suresh, S.3, Henry, L.1, Gordon, M.1, Rossignol, J.2-4, Morgan, D.1 & Dunbar G.L.2,3,5

¹Department of Translational Neuroscience, Michigan State University, Grand Rapids, MI; and ²Field Neurosciences Inst. Lab. for Restorative Neurol.; ³Program in Neuroscience; ⁴Col. of Medicine; ⁵Dept. of Psychology, Central Michigan Univ., Mount Pleasant, MI.

Introduction

Alzheimer's disease (AD) is a complex neurodegenerative disease that causes behavioral deficits and pathophysiological hallmarks, including, tau and amyloid- β dysfunction and accumulation, neuroinflammation and subsequent synaptic loss and neuronal death. Neuroinflammation contributes to tau neurofibrillary tangle (NFT) formation (1). Microglia phagocytose debris and astrocytes form the blood brain barrier (2). The type, severity, and duration of inflammation contribute to neurodegeneration (3). The goal of this study is to determine the relationship and function between peripheral chronic inflammatory stress and behavioral and pathological hallmarks of AD—tau accumulation and immune system activation. We hypothesize that inflammatory stress increases pathological tau burden and contributes to the pro-inflammatory state characteristic of AD.

Methods

This study used PS19 mice—harboring a microtubule associated protein tau (MAPT) mutation (P301S). Non-Tg (C3H and C57BL/6) littermate counterparts were used as controls.

Mice received intraperitoneal injections (I.P.) of either infection mimetics (I.F.) (pro-inflammatory compounds: Zymosan A, Poly I:C, ODN2395, LPS) or mammalian ringers (control). Injections were administered starting at 6 months of age, spaced 1 month apart per injection, with a total of 4 injections per mouse.




FIGURE 1: Experimental Design




FIGURE 2: Experimental timeline, including I.P. injections, behavioral testing, hemi-brain collections, microscopy, and quantification.

P-Tau (AT8) and Total Human Tau (HT7)




FIGURE 3: Tau IHC results, 20x objective. AT8 (p-tau) A. PS19 Ringers, B. PS19 I.F. HT7 (total human tau) C. PS19 Ringers, and D. PS19 I.F.

Astrocyte Reactivity (GFAP) IHC Results




FIGURE 4: GFAP IHC results, 20x objective. A. PS19 Ringers, B. PS19 I.F. C. Non-Tg Ringers, and D. Non-Tg I.F.

Tau IHC and Behavior




FIGURE 5: AT8 was greater in I.F. mice compared to Ringers in the A. whole-hemisphere (WH) ($p = 0.0063$), and in the B. hippocampus ($p = 0.049$). Total tau did not sig differ (HT7) C in the WH or D. the hippocampus. WH GFAP was significantly elevated in the E. I.F. group compared to the PS19 Ringers groups ($p = 0.0226$). F. I.F. mice showed elevated basic movement ($p = 0.0367$) G. and SOR ($P = 0.0173$).

Summary

Peripheral and chronic inflammatory stress contribute to tau dysfunction—hyperphosphorylation and aggregation in the PS19 mouse model of AD. Inflammatory stimuli did not significantly alter total tau (HT7), suggesting that the immune system plays a specific role affecting only pathological forms of tau. The pro-inflammatory response may contribute to microtubule dissociation and accumulation in AD. These findings demonstrate that the immune system plays an important role in the onset and progression of the disease.

References

- Didonna, A. (2020). Tau at the interface between neurodegeneration and neuroinflammation. *Genes & Immunity*, 21(5), 288-300.
 - Kwon, H. S., & Koh, S. H. (2020). Neuroinflammation in neurodegenerative disorders: The roles of microglia and astrocytes. *Translational neurodegeneration*, 9, 1-12.
 - Walker, K. A., Fieck, B. N., & Westbrook, R. (2019). Understanding the role of systemic inflammation in Alzheimer's disease. *ACS Chemical Neuroscience*, 10(8), 3340-3342.
- Support for this study was provided by NIH R01AG051500 to Dr. David Morgan at Michigan State University and the Neuroscience Program, College of Medicine, the John G. Kulhavi Professorship in Neuroscience, and the E. Malcolm Field and Gary Leo Dunbar Endowed Chair in Neuroscience at Central Michigan University.

