RAJALAKSHMI ENGINEERING COLLEGE

Keerthana K 251801135-2025-AIDS

K2

Started on	Thursday, 24 September 2026, 11:24 AM
State	Finished
Completed on	Thursday, 24 September 2026, 11:24 AM
Time taken	47 secs
Marks	1.00/1.00
Grade	**10.00** out of 10.00 (**100**%)

Question 1 | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that A[j] - A[i] = k, i != j.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as 5 - 1 = 4

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```
#include <stdio.h>

int main() {
    int n, k, i = 0, j = 1;
    scanf("%d", &n);

    int a[n];
    for(int x = 0; x < n; x++)
        scanf("%d", &a[x]);

    scanf("%d", &k);

    while(j < n) {
        if(a[j] - a[i] == k) {
            printf("1");
            return 0;
        }
        if(a[j] - a[i] < k)
            j++;
        else
            i++;

        if(i == j)
            j++;
    }

    printf("0");
    return 0;
}
```

	Input	Expected	Got	
✔	3 1 3 5 4	1	1	✔
✔	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✔
✔	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✔
✔	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✔

Passed all tests! ✔

Correct

Marks for this submission: 1.00/1.00.

Back to Course