

Math 521 Homework 4

Due on Tuesday, April 17, 2018

Kristin Holmbeck

Contents

Theory	2
1.	2
2.	2
3.	2
Computing	3
1.	3
Code	4
Gram-Schmidt	4

List of Figures

Theory

1.

Consider the two eigenvector problems

$$C_x \mathbf{u} = \lambda_x \mathbf{u}$$

and

$$C_s \mathbf{v} = \lambda_s \mathbf{v}$$

where the matrices are related by $C_s = C_x + \alpha I$, where α is a real number and I is the usual identity matrix. Show that if \mathbf{u} is an eigenvector of C_x , then it is also an eigenvector of C_s associated with eigenvalue $\lambda_s = \lambda_x - \alpha$.

$$\begin{aligned} C_x \mathbf{u} &= \lambda_x \mathbf{u} = C_s \mathbf{u} + \alpha I \mathbf{u} \\ C_s \mathbf{u} &= \lambda_x \mathbf{u} - \alpha I \mathbf{u} \\ C_s \mathbf{u} &= (\lambda_x - \alpha) \mathbf{u} \\ \text{but } C_s \mathbf{v} &= \lambda_s \mathbf{v} \\ \implies \lambda_s &= \lambda_x - \alpha \quad \text{is an eigenvalue with associated eigenvector } \mathbf{u} \end{aligned}$$

2.

Let $A \in \mathbb{R}^{m \times n}$. Show that the matrix M defined as

$$M = \alpha^2 I + AA^T, \quad \alpha \neq 0 \in \mathbb{R}$$

is nonsingular, where $I = I_m$ and α is a nonzero real number.

For the trivial case, suppose $A = 0_{m \times n}$. Then $M = \alpha^2 I$ which is clearly nonsingular since $\alpha \neq 0$.

3.

Show that the between-class scatter matrix, S_B , in the multi-class *Fisher Discriminant Analysis* is given by

$$S_B = \sum_{i=1}^M n_i (\mathbf{m}_i - \mathbf{m})(\mathbf{m}_i - \mathbf{m})^T,$$

where M is the total number of distinct classes, n_i is the number of data points in class i , \mathbf{m}_i is the class mean of the i^{th} class, and \mathbf{m} is the mean across all n data points. You may use the facts that

$$S_T = S_B + S_W, \quad S_W = \sum_{i=1}^M \sum_{x \in D_i} (x - \mathbf{m}_i)(x - \mathbf{m}_i)^T, \quad \text{and} \quad S_T = \sum_{i=1}^n (x_i - \mathbf{m})(x_i - \mathbf{m})^T$$

Computing

1.

This project concerns the application of the KL procedure for incomplete data [3]. Let the complete data set be translation-invariant:

$$f(x_m, t_\mu) = \frac{1}{N} \sum_{k=1}^N \frac{1}{k} \sin[k(x_m - t_\mu)],$$

where $m = 1, \dots, M$, with M dimension of the ambient space (size of the spatial grid), and $\mu = 1, \dots, P$, with P the number of points in the ensemble.

Let $x_m = \frac{2\pi(m-1)}{M}$ and $t_\mu = \frac{2\pi(\mu-1)}{P}$.

Select an ensemble of masks $\{\mathbf{m}^{(\mu)}\}$, $\mu = 1, \dots, P$. where 10% of the indices are selected to be zero for each mask. Each pattern in the incomplete ensemble may be written as

$$\tilde{\mathbf{x}}^{(\mu)} = \mathbf{m}^{(\mu)} \cdot \mathbf{f}^{(\mu)},$$

where $(\mathbf{f}^{(\mu)})_m = \frac{1}{N} \sum_{k=1}^N \frac{1}{k} \sin[k(x_m - t_\mu)]$. Let $P = M = 64$ and $N = 3$.

- Compute the eigenvectors of this ensemble using the gappy algorithm [3].
- Plot the eigenvalues as a function of the iteration, and continue until they converge.
- Plot your final eigenfunctions corresponding to the 10 largest eigenvalues.
- Plot the element $\tilde{\mathbf{x}}^{(1)}$ and the vector $\tilde{\mathbf{x}}_D$ repaired according to Equation

$$\tilde{\mathbf{x}} \approx \tilde{\mathbf{x}}_D = \sum_{n=1}^D \tilde{a}_n \phi^{(n)}. \quad (1)$$

Determine the value of D that provides the best approximation to the original non-gappy pattern vector.

Code

Gram-Schmidt

References

- [1] Chang, Jen-Mei. *Matrix Methods for Geometric Data Analysis and Recognition*. 2014.
- [2] P. N. Belhumeur, J. P. Hespanha and D. J. Kriegman, "Eigenfaces vs. Fisherfaces: recognition using class specific linear projection," in *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol. 19, no. 7, pp. 711-720, Jul 1997.
- [3] R. Everson and L. Sirovich. The karhunen-loeve transform for incomplete data. *J. Opt. Soc. Am., A*, 12(8):1657-1664, 1995.