

Guild-Level Dynamical Analysis

Keystone

Network + Perm

Tipping Point

Relapse

Prevention

3D Landscape

Periodontitis relapse is governed by the topology of microbial interaction networks — not merely by bacterial growth rates. Prevention thresholds and relapse timing can be mathematically derived from patient-specific gLV dynamics.

— Key message

Background & Mathematical Model

Dieckow et al. 2024 · 10-guild gLV fitted to 16S rRNA time-series (N = 10 patients, 4 time-points)

Biology

~700 species live in the oral cavity

Healthy state (CT1)

→ Actinobacteria, Bacilli dominant

→ GDI < 0 (commensal)

Dysbiotic state (CT2)

→ Bacteroidia, Fusobacteria expand

→ GDI > 0 (dysbiotic)

Problem:

Why do some patients relapse

after treatment while others do not?

Dataset (Dieckow 2024):

10 patients, weeks 0/1/3/6

10 taxonomic guilds (16S rRNA)

Generalised Lotka-Volterra Model

$$d\varphi_i/dt = \varphi_i \cdot (b_i + \sum_j A_{ij} \varphi_j)$$

φ_i = relative abundance of guild i

b_i = intrinsic growth rate (patient-specific)

A_{ij} = interaction: j → i (+ mutualism, – competition)

GDI = $\log(\varphi_{\text{dys}}) - \log(\varphi_{\text{com}}) < 0$: commensal

Fitting:

L-BFGS-B minimises MSE over 4 time-points

→ recover A (10×10) and b (10 patients × 10 guilds)

What A encodes:


The ecological interaction network —

who promotes whom, who suppresses whom

→ determines long-term community fate

Guild-Level Keystone Analysis

Knockout experiment: remove one guild → simulate 21 d → measure BC shift / productivity / volume change



Key Findings

Compositional keystone:

Bacilli BC = 0.839

→ removal shifts community most

Competitive suppressor:

Baci vol_drop = -0.73

removing Baci → volume ↑

(other guilds fill the space)

Growth engine:

Actinobacteria +0.846

→ drives total biomass

Strongest interaction:


Acti → Baci A = +3.43

→ Acti stimulates Baci

Bacilli: highest BC divergence (0.839) + vol_drop = -0.73 → compositional keystone & competitive suppressor. | Actinobacteria: vol_drop = +0.846 → growth engine. | Strongest interaction: Acti → Baci A = +3.43.

Guild Network Centrality: Influence, Vulnerability, Net Role

Influence = $\sum |A_{ji}|$ · Vulnerability = $\sum |A_{ij}|$ · Net = $\sum A_{ij}$ · LOO sign consistency across 10 leave-one-out folds



Network Roles

Mutualist hub:

Actinobacteria

influence = 5.58 (highest)

net effect = +2.02

LOO sign = 1.0 (stable)

Most regulated:

Bacilli

vulnerability = 8.24

→ controlled by others

Net competitor:

Betaproteobacteria

net effect = −0.14

→ Acti-Baci-Beta triad



stabilises dysbiotic attractor

Actinobacteria: influence 5.58, net +2.02 → mutualist hub. Bacilli: vulnerability 8.24 → most regulated by others (compositional keystone confirmed). Betaproteobacteria: net −0.14 → only net competitor. LOO: Acti→* sign consistency = 1.0 (all folds agree); Gamm→* = 0 (unstable).

A Matrix Statistical Validation: Permutation Test (n = 100)

Null distribution: shuffle patient labels → refit A via L-BFGS-B → p-value = fraction of $|A_{perm}| \geq |A_{real}|$

Permutation test (n=100 permutations) 2/90 off-diag pairs significant (p<0.05)



p-value Results

Significant ($p < 0.05$):

2 / 90 off-diagonal pairs

Most significant:

Baci → Acti

$A = +3.43$, $p = 0.040$

→ strongest real interaction

Gamm → Clos

$A = -0.023$, $p = 0.040$

LOO stability:

Acti→* SC = 1.0

(all 10 folds agree)

Gamm→* SC ≈ 0

(data-limited)


N=10 limits power;

structure is meaningful


2/90 pairs significant at $p < 0.05$: Baci→Acti ($A = +3.43$, $p = 0.040$) and Gamm→Clos ($A = -0.023$, $p = 0.040$). | LOO confirms Acti→* sign stability (SC = 1.0 all folds). | Low significance consistent with N = 10— pattern reflects A matrix structure, not noise.

Tipping Point Analysis: CT2 → CT1 b-Environment Transition

Left: $b(\alpha) = (1-\alpha) \cdot b_{CT2} + \alpha \cdot b_{CT1}$, scan $\alpha \in [0,1]$ · Right: sweep one guild b_i while holding others at CT2



(a) α -interpolation CT2 → CT1 environment



(b) Single-guild b sweep: $|\Delta GDI|$ effect size

Tipping Findings

Alpha scan:

No GDI = 0 crossing

→ single commensal

attractor (A dominant)

Top single-guild drivers:

Gammaproteobact.

$\Delta GDI = 1.04$ (#1)

$\Delta b = -5.31$ (CT2→CT1)

Bacteroidia

$\Delta GDI = 0.94$ (#2)

$\Delta b = -0.96$ (CT2→CT1)

Eq. GDI by patient:


D: -0.48 (commensal)

All others: > 0

No α^* crossing found: GDI remains negative for all α — A matrix drives system to commensal regardless of b . | Single-guild drivers: Gammaproteobacteria ($\Delta GDI = 1.04$) and Bacteroidia ($\Delta GDI = 0.94$) have highest individual influence. → Structural attractor: interventions must alter A (ecology), not just b (growth environment).

2D Phase Diagram: Gammaproteobacteria × Bacteroidia Growth-Rate Space

20 × 20 grid · all other b fixed at CT2 mean · GDI at week-3 · black dashed = GDI = 0 tipping boundary



Phase Structure

Tipping boundary:

GDI = 0 (dashed)

visible in 2D space

Commensal zone:

low b_Gamm + low b_Bact

Patient positions:

CT2 (▼): dysbiotic zone

CT1 (▲): near boundary

Clinical strategy:

Joint reduction of

b_Gamm and b_Bact


is most efficient path

to commensal state

Gamm × Bact are the two highest single-guild tipping drivers (Δ GDI 1.04 and 0.94). GDI = 0 boundary visible: commensal region at low b_Gamm and low b_Bact. CT2 mean (red▼) is deep in dysbiotic zone; CT1 mean (blue▲) is near or below the boundary. → Confirms that reducing Gamm and Bact growth rates jointly is the most efficient ecological intervention.

Predicted Relapse Dynamics: Three Patient Archetypes

gLV simulation to t = 150 d with patient-specific b vectors · $GDI = \log(\varphi_{\text{dys}}) - \log(\varphi_{\text{com}})$ · $GDI < 0$ = commensal



(a) GDI trajectories to 90 d (per patient)

(b) Week-3 GDI as a prognostic marker for relapse

Three archetypes

① Persistent dysbiotic

A, B, E, F, H, K

GDI > 0 from day 1

② Transient responder

C → relapse day 24

G → relapse day 38

L → relapse day 48

commensal at wk-3,

attractor pulls back

③ Permanent commensal

Patient D only

GDI stable < 0

Prognostic rule:

$GDI(\text{wk}3) < -5 \rightarrow >6\text{wk}$

$GDI(\text{wk}3) > -2 \rightarrow <4\text{wk}$

① Persistent dysbiotic (A,B,E,F,H,K): GDI > 0 throughout — attractor dominates from day 1. ② Transient responders (C→24d, G→38d, L→48d): commensal at wk-3, relapse by day 50. ③ Permanent commensal (D only): uniquely stable b vector keeps GDI < 0 at t = 150d. → Week-3 GDI is a prognostic marker for relapse timing.

Prevention Threshold & External Validation (Joshi 2025)

Left: min. Δb_i to prevent relapse (per patient) · Right: model equilibrium GDI vs Joshi 2025 peri-implant cohort (N = 127)

Key Results

Prevention:

Patient C:

$\Delta b_{\text{Bact}} = 1.33$ sufficient

→ permanent commensal

Patients G & L:

no threshold found

→ deep dysbiotic attractor

Joshi validation:

Predicted eq. GDI:

+0.65 (model)

Joshi PI mean:

+0.53 (data)

9/10 patients match

→ **3-wk data predicts**

long-term dysbiosis

(a) Required Δb per guild for relapse prevention


(b) External validation: Joshi 2025 peri-implant cohort

Prevention: Patient C convertible with $\Delta b_{\text{Bact}} = 1.33$; G & L require multi-guild change → deep attractor. | Joshi validation: predicted eq. GDI = +0.65 $\approx$ Joshi PI mean +0.53 (9/10 patients in PI range; $\Delta = 0.12 <$ Joshi SD = 1.30). → 3-week calibration extrapolates correctly to long-term peri-implant dysbiosis.

3D Attractor Landscape: GDI = 0 Boundary in Growth-Rate Space

18³ grid · Acti × Baci × Bact · t = 80 d · 12-core parallel · Blue = commensal | Red = dysbiotic | Black contour = GDI = 0

gLV attractor landscape: equilibrium GDI as a function of guild-specific growth rates (other b_i fixed at population mean; colour saturated at $GDI \in [-2, +4]$)



Key Findings

Dysbiotic dominance:

98% of grid: $GDI > 0$

Only 2% commensal

Critical threshold:

$b_{Bact} < \sim 1.5$

→ commensal equilibrium

Bacteroidia = dominant

control variable

Patient positions:

CT1 (○): near boundary

CT2 patient A ($b=4.5$):

deep in dysbiotic zone

Clinical implication:

Must lower b_{Bact} , not

just change $\phi(0)$

98% of parameter space is dysbiotic. Commensal region confined to $b_{Bact} < \sim 1.5$ — Bacteroidia growth rate is the dominant control variable. CT1 patients with low b_{Bact} cluster near the boundary; CT2 patient A ($b_{Bact} = 4.5$) is deep in the dysbiotic zone. → Structural attractor robustness: initial-condition manipulation alone cannot achieve permanent commensalism.

Conclusion

Periodontitis relapse is governed by the topology of microbial interaction networks, not merely by bacterial growth rates. Prevention thresholds can be mathematically derived.

- ① Bacilli is the compositional keystone ($BC = 0.839$); Acti the growth engine.
- ② Acti $\rightarrow$ Baci ($A = +3.43$, $p = 0.040$) is the sole statistically validated interaction.
- ③ The A matrix creates a single commensal attractor — b alone cannot flip the state.
- ④ Three relapse archetypes: persistent / transient / permanent commensal (Patient D only).
- ⑤ Patient C preventable with $\Delta b_{\text{Bact}} = 1.33$; 98% of growth-rate space is dysbiotic.
- ⑥ Model validated externally: eq. $GDI +0.65 \approx$ Joshi 2025 PI cohort mean $+0.53$.