A two-level biofilm model from one extended Hamilton principle

Keisuke Nishioka, working note, 15 October 2026 (not part of the thesis)

Aim. In the thesis the growth field of Klempt et al. 2024 and the point model of Klempt et al. 2026 are solved separately and joined at the Gauss points. This needs choices that no paper fixes: how the void of the point model relates to the void of the field, whether the point model is scaled to the field, what drives the growth, and how the two clocks are linked. Here all equations follow from one energy, one dissipation potential and one constraint.

1 Why one level is not enough

The direct generalisation takes every species as a field $\phi_i(\mathbf{X},t)$ with one void $\phi_0 = 1-\sum_i \phi_i$ and the point-model energy at every point. The point model keeps every ϕ_i away from zero by its barrier energy $K/[\phi_i^2(1-\phi_i)^2]$, and its production is local. A 1D solution with all fields in one Newton system shows the consequence: the outer liquid fills up from the barrier floor everywhere at once, so the model cannot represent a biofilm next to liquid. The two voids are therefore different physical objects: the liquid outside the biofilm, and the liquid inside it. This motivates two levels.

2 State, energy, dissipation

$\phi(\mathbf{X},t) \in [0,1]$	biofilm phase (Klempt et al. 2024): biofilm against outer liquid
$s_i,\ s_0$	composition of the biofilm: species $i = 1 \dots n$ and inner liquid, $\sum_i s_i + s_0 = 1$
ψ_i	living fraction of species i (Klempt et al. 2026)
α	local expansion, $\mathbf{F}_g = \alpha \mathbf{I}$, $\alpha(0) = 1$
$\phi s_i,\ \phi \sum_i s_i \psi_i$	absolute amount of species i, living biomass

With $w = \phi + \varepsilon$ ($\varepsilon > 0$ small, it keeps the composition equations regular in the outer liquid where they carry no weight), the energy density is

$$E = \frac{\beta}{2}|\nabla\phi|^2 + \lambda w \Psi_{\mathrm{pm}}(s, s_0, \psi) + w\frac{\beta_s}{2}\sum_i |\nabla s_i|^2 - k_\alpha(\alpha-1)\,\phi \sum_i s_i\psi_i + E_{\mathrm{mech}}, \tag{1}$$

with the point-model energy of Klempt et al. 2026

$$\Psi_{\mathrm{pm}} = K\Big[\sum_i B(s_i) + B(s_0) + \sum_i B(\psi_i)\Big] + \frac{\alpha^*}{2}\sum_i b_i\psi_i^2 - \frac{c^*}{2}\sum_{ij} A_{ij}\, s_i\psi_i\, s_j\psi_j, \qquad B(x) = \frac{1}{x^2(1-x)^2}, \tag{2}$$

and $E_{\mathrm{mech}} = \frac{\mu}{2}\phi^2(\mathbf{I}:\mathbf{C}_e - 3)$ as in Klempt et al. 2024 (dropped below for the same reason as there: its effect on growth is small). The dissipation potential is

$$D = \frac{\eta_\phi}{2}\dot\phi^2 + \lambda w D_{\mathrm{pm}} + \frac{\eta_\alpha}{2}\dot\alpha^2, \qquad D_{\mathrm{pm}} = \sum_i \Big[\frac{\eta}{2}\dot s_i^2 + \frac{\eta}{2}\big(\tfrac{\mathrm{d}}{\mathrm{d}t}(s_i\psi_i)\big)^2\Big] + \frac{1}{2}\dot s_0^2, \tag{3}$$

and the constraint is imposed with the multiplier field γ through $\lambda w\gamma(\sum_i s_i + s_0 - 1)$. The only new parameters are λ (the energy scale of the composition relative to the phase), β_s (spreading of the composition inside the biofilm) and the form of the growth energy.

3 Stationarity conditions

The extended Hamilton principle

$$\delta_q \int_\Omega E\,\mathrm{d}V + \int_\Omega \partial_{\dot q} D\,\delta q\,\mathrm{d}V = 0$$

for all variations δq of $q = \{\phi, s_i, s_0, \psi_i, \alpha, \gamma\}$, with $\mathbf{n}\cdot\nabla\phi = \mathbf{n}\cdot\nabla s_i = 0$ on the boundary, gives:

Composition (divided by $\lambda w > 0$):

$$KB'(s_i) + \gamma + \eta(1+\psi_i^2)\dot{s}_i + \eta s_i \psi_i \dot{\psi}_i - c^* \psi_i \Big(\sum_j A_{ij} s_j \psi_j + \frac{k_\alpha(\alpha-1)\phi}{\lambda w c^*} \Big) - \frac{\beta_s}{\lambda w} \nabla \cdot (w \nabla s_i) = 0, \quad (4)$$

$$KB'(s_0) + \gamma + \dot{s}_0 = 0, \quad (5)$$

$$KB'(\psi_i) + \alpha^* b_i \psi_i + \eta s_i \psi_i \dot{s}_i + \eta s_i^2 \dot{\psi}_i - c^* s_i \Big(\sum_j A_{ij} s_j \psi_j + \frac{k_\alpha(\alpha-1)\phi}{\lambda w c^*} \Big) = 0, \quad (6)$$

$$\textstyle\sum_i s_i + s_0 - 1 = 0. \quad (7)$$

Without the growth term and without composition gradients these are exactly the point model of Klempt et al. 2026 at every point. The growth energy enters with the same structure as the interactions.

Phase:

$$\eta_\phi \dot{\phi} - \beta \nabla^2 \phi + \lambda \Psi_{\mathrm{pm}} + \frac{\beta_s}{2} \sum_i |\nabla s_i|^2 - k_\alpha(\alpha-1) \sum_i s_i \psi_i = 0. \quad (8)$$

The composition acts on the phase through $\lambda \Psi_{\mathrm{pm}}$: where the composition energy is low (strong favourable interactions), the biofilm tends to grow, and the growth source is weighted by the living share. The dependence of the dissipation weight w on ϕ is not a force: the principle varies D with respect to the rates only.

Expansion:

$$\eta_\alpha \dot{\alpha} = k_\alpha \phi \sum_i s_i \psi_i. \quad (9)$$

4 Limits

1. $\lambda \to 0$, $\beta_s = 0$: the composition at every point is the point model on its own clock, started where the biofilm is, and (9) is the growth law of the element's scheme (c), $\dot{\alpha} = \kappa \phi_{3D} \sum_i \bar{\phi}_i$ with $\kappa = k_\alpha / \eta_\alpha$. Scheme (c) is thus the decoupled limit of this model, and its point model, which runs unscaled as in scheme (b), is the consistent one: its ϕ_0 is the inner liquid s_0.

2. With $\phi \sum_i s_i$ in place of the living biomass in the growth energy and one species, (9) is Eq. 36 of Klempt et al. 2024 up to the factor $\sum_i s_i$.

3. Scheme (a), which scales the point model to $\sum_i \phi_i = \phi_{3D}$, identifies the inner liquid with the outer one ($s_0 = 1 - \phi$). It is not a limit of the model.

5 Energy

Proposition 1. *Let E not depend on time explicitly and let no external power act. Along solutions,*

$$\frac{\mathrm{d}}{\mathrm{d}t} \int_\Omega E \, \mathrm{d}V = -2 \int_\Omega D \, \mathrm{d}V \leq 0.$$

Proof. Every stationarity condition has the form $\partial_q E + (\text{gradient terms}) + \partial_{\dot{q}} D + (\text{constraint terms}) = 0$. Multiply each by $\dot{q}$, integrate over Ω and add. The gradient terms integrate by parts to the rate of the gradient energies (the boundary terms vanish by the no-flux conditions); the constraint terms give $\lambda w \gamma \frac{\mathrm{d}}{\mathrm{d}t}(\sum_i s_i + s_0) = 0$; and D is quadratic in the rates, so $\sum_q \dot{q} \, \partial_{\dot{q}} D = 2D$. What remains is $\frac{\mathrm{d}}{\mathrm{d}t} \int E = -\int 2D$. □

Remark 1. *Switching an antibiotic on (α^* changing in time) adds $\int \partial_t E = \int \lambda w \frac{\dot{\alpha}^*}{2} \sum_i b_i \psi_i^2$, an external input; this is the jump seen at the switch in the 1D runs. Klempt et al. 2024's front term h_ϕ^* is not derived from a potential, so it is left out here; a model that includes it is no longer a gradient flow.*

6 Discretisation and checks

1D, nodes with mirrored ends, all unknowns of all nodes in one residual, implicit Euler, full Newton to $|R| < 10^{-10}$ with damping that keeps $s, \psi \in (0,1)$ and $\phi > 0$ (`JAXFEM/unified_two_level_1d.py`). For the one-level model the same scheme reproduces the converged point model to 1.6×10^{-13} and the discrete energy decreases at every step (`unified_1d_monolithic.py`).

7 What remains assumed

The form of the growth energy (living biomass, and $\alpha - 1$ so that it vanishes before growth); λ and β_s; the clock of the composition relative to the phase, which is now the ratio of the dissipation coefficients η/η_ϕ and can be fixed by the measured volume; the stiffness; detachment.