Master's Thesis: Status and Plan

Biofilm growth coupled into the IKM FEM element, following Klempt et al.

Keisuke Nishioka

Institute of Continuum Mechanics (IKM), Leibniz Universität Hannover
Keio University


20 October 2026

Since the last meeting: the scope of the FEM part is fixed

	content	reference	coupling
thesis (LUH)			
one species	growth law, Eq. 36	Klempt et al. 2024	field → growth
two species	composition from the point model	Klempt et al. 2026	field → composition (one-way)
continuation (Keio)			
five species	calibrated point model; first runs in Appendix G	none yet	two-way: composition and stress act back

One and two species can be checked against published examples; five species have no reference yet.

Verification, step by step




Each step is checked before the next one builds on it.

- point model: two misprints in Klempt et al. 2026 identified (no $+\gamma$ in Eq. 17, no $\frac{1}{2}$ in Eq. 19)
- same point model in Fritsch et al. 2025: their figures reproduced within 0.013

The coupling: what the element does and what is added


The element solves the biofilm field of Klempt et al. 2024 (Eq. 34):

$$\dot{\phi} - \beta\,\Delta\phi - k_\alpha\,\alpha + |\nabla\phi|\,r\,\frac{c}{k+c}\,\mathbf{n}_\phi \cdot \mathbf{n}_c = 0$$




Klempt 2024 Table 2: $k_\alpha = 10^{-3}$, $E = 10$ Pa, $\nu = 0.49$. β, r, k, d, g: the framework's example input, not from the paper.

Model and boundary conditions



- 2 mm cube, 512 elements; only rigid-body motion fixed
- nutrient $c = 1$ on the face $y = -1$ mm; seed of 32 elements with $\phi = 1$
- uniform growth is stress free; stress comes from differences in growth
- used for Checks 1–3 and the two-species runs (ANSYS); the Klempt 2024 reproduction uses the paper's 20 μm cube (Python)

What I implemented



speed single-element run 357 s → **5.6 s**

check every ANSYS call replayed in Python, difference 0


used as it is

- IKM element: field, stiffness
- point-model equations (Klempt et al. 2026)

written in this work

- Eq. 36 and the composition step in the material call
- Fortran → Python bridge, sub-stepping
- point-model solver (JAX): all 14 paper examples
- checks: closed form in ANSYS, call replay, about 440 tests

Second version: the point model in Fortran




- replaces the bridge and the Python process; the material call is unchanged
- per sub-step: Newton on the 12 unknowns $\phi_1..\phi_5,\ \phi_0,\ \psi_1..\psi_5,\ \gamma$ (6 iterations), exact Jacobian written out, LU with pivoting

check	**result**
against the Python solver (16 states, 4 cases)	largest relative difference 8×10^{-15}
ANSYS, 8^3, two species, one week	stress, nutrient identical; point model $\leq 1.6 \times 10^{-12}$
run time, same model	156 s $\rightarrow$ **36 s**

No server or port: one executable, so runs on a cluster or in parallel need no extra process.


Checks 1 and 2: exact solution and whole model

1. Exact solution




α matches to 3×10^{-11}

2. Whole model, 512 elements



same pattern (cosine 1.0000); factor $0.455 = \frac{1}{2} \cdot \frac{10}{11}$

Check 3: the seeded element and its neighbours




- $\alpha - 1 = 1.1 \times 10^{-3} = k_\alpha \phi t$, as Eq. 36 says
- seed compressed; neighbours carry about 25× its von Mises stress
- von Mises/$E \leq 10^{-5}$: slow growth (stresses scale with E)

Four things found along the way (partner element)

	what I found	effect	what I did
1	growth variable is the mean of both species fields	growth halved with one species (unused field stays at its start value)	asked Oliver whether intended
2	start values must satisfy $\phi_1 + \phi_2 \leq 1$	example input starts both at 1	one species: second field starts at 0
3	first result set not in equilibrium	zero stress while the element sets up its operators	first result set excluded
4	front-growth term inactive (reading slip); $\lvert\Delta\phi\rvert$ where Eq. 34 has $\lvert\nabla\phi\rvert$	seed does not spread	corrected in a test copy: then ϕ exceeds 1

Checks 1–3 do not depend on these four points.

Two species: composition from the point model (Klempt et al. 2026)



Point model run two ways: A) rescaled every step to ϕ_{3D}; B) started when ϕ_{3D} first reaches ϕ_{min}, then independent. Assumed, not from a paper: $s = 0.15$, $\phi_{cap} = 0.9$, $\phi_{min} = 0.01$.

- the field sets the **amount**, the point model the **composition**
- one-way: the composition does not act on the stress


Two species: results in the packed seed (ANSYS, $T^* = 1$)

case (Klempt et al. 2026)	paper	coupled, $\phi_{\mathrm{cap}} = 0.9$	$\phi_{\mathrm{cap}} = 0.85$ / 0.95
3: coexistence	0.60	0.667	0.668 / 0.664
6: one species wins	≈ 0 ($\psi_1 = 0.05$)	0.021 ($\psi_1 = 0.048$)	–

share $\phi_1/(\phi_1 + \phi_2)$; paper: point model alone; parameters $c^* = 100$, $\alpha^* = 10$

- same type of outcome; the ratio shifts because the field sets the amount
- converged in the coupling time step

Added since 5 October: growth from the living biomass



Lysis or an antibiotic in the point model lowers ψ_i and so the growth. Assumed, not from a paper: κ (mean growth of the seed at $T^* = 1$ as Eq. 36), $s = 0.2375$ ($T^* = 1$ = Day 21).


Eq. 36 uses the amount: $\dot{\alpha} = k_\alpha \phi$.
Here: $\dot{\alpha}/\alpha = \kappa\, \phi_{3D} \sum_i \phi_i \psi_i$, so lysis in the point model lowers the growth.

κ chosen so that the mean growth of the seed equals Eq. 36 (an assumption, not from a paper; Appendix E).

seed, $T^* = 1$	Eq. 36	living	diff.
$\alpha - 1$, 8^3	$6.752 \cdot 10^{-4}$	$6.752 \cdot 10^{-4}$	0.0 %
von Mises/E, 8^3	$4.684 \cdot 10^{-5}$	$4.357 \cdot 10^{-5}$	−7.0 %
von Mises/E, 16^3; 24^3			−7.1; −7.0 %
antibiotic vs. none, 8^3	0 %	−32 %	

five species calibrated (CH), $E = 10$ Pa; the antibiotic case with the previous calibration

Growth from the living biomass with synthetic data




Same seed problem; point model: four-species case of Fritsch et al. 2025 with its true parameters (no measured data), own void, start $\phi_i = 0.02$. $T^* = 1$ spans their 1500-step validation run ($s = 0.15$); antibiotic $\alpha^* = 50$ from $T^* = 0.5$ as in their Fig. 18. κ as before (same seed growth at $T^* = 1$, an assumption). Local nutrient: $c^* = 25\,c(x)$, $c = 1$ on one face, $\Lambda^2 = gL^2/d = 16$. (c): $\alpha - 1$ on the mid-plane with the local nutrient.

seed, $T^* = 1$, 32^3 field, 16^3 stress	$\alpha - 1$	von Mises
Eq. 36	$6.56 \cdot 10^{-4}$	$6.07 \cdot 10^{-4}$ Pa
living biomass, uniform c	same	+12 %
antibiotic from $T^* = 0.5$	−12 %	−13 %
local nutrient, $\Lambda = 4$	−58 %	−68 %

The sign of the stress change follows the time course of the living biomass: here it rises from 0.08 (growth comes late, +12 %), in the calibrated run it falls (−7 %).

Two modelling assumptions, each with a sensitivity study


1. time link $s = 0.15$



each step: point model advances $s\,\Delta t$
(no paper links the two clocks)

case (Klempt et al. 2026)	effect of s
6: one species wins	winner appears for $s \gtrsim 0.15$
3: coexistence	share moves by 0.03–0.06

2. cap $\phi_{\mathrm{cap}} = 0.9$




the amount given to the point model
is $\min(\phi_{3D}, \phi_{\mathrm{cap}})$

ϕ_{cap}	0.85	0.90	0.95
case 3 share	0.668	0.667	0.664

All other parameters are the published values.


Limits

- composition is local, not carried with the front: effect on the mean ≤ 0.001 (right, Python)
- s has no source in a paper
- Eq. 34 as printed moves the colony instead of growing it (Appendix D)
- two α: the element's own and Eq. 36 are integrated separately (Check 2)



Composition at $T^* = 1$ of the points first reached at each time; Klempt 2024 test case 4.1 field (Python, 20 μm cube), Klempt 2026 cases 3 and 6.

Status and timeline



- examiners Prof. Junker (IKM), Dr. Wangenheim (IDS); supervisors Dr. Soleimani, Dr. Geisler
- draft sent on 9 October; **submission 13 November**
- colloquium 16 December, 13:00, IKM room 320
- growth from the living biomass: not in the thesis, continued at Keio

Questions for this meeting

1. **Your coupling ideas:** which direction would you take first?
2. **Growth law:** amount of the field (Eq. 36) or living biomass of the point model?
3. **Two-way coupling:** should the stress act back on the growth or the composition?
4. **Rate κ:** fixed by the mean growth of Eq. 36, or by the measured volume?
5. **Next steps:** what should be done here, and what at Keio?

References

- Klempt, Soleimani, Junker (2023), PAMM: precursor spatial model
- Klempt, Soleimani, Wriggers, Junker (2024), Biomech. Model. Mechanobiol.: growth field, Eq. 34/36, Table 2
- Klempt, Geisler, Soleimani, Junker (2026), Arch. Appl. Mech.: multi-species point model, Table 1 cases
- Fritsch et al. (2025), arXiv:2512.15145: Bayesian updating, same model

Appendix A: notation

Growth field and mechanics (Klempt et al. 2024)

ϕ	biofilm volume fraction (0 = void, 1 = dense)
c	nutrient concentration
α	expansion variable, $\mathbf{F}_g = \alpha \mathbf{I}$, $\alpha(0) = 1$; growth is $\alpha - 1$
$\mathbf{F} = \mathbf{F}_e \mathbf{F}_g$	elastic / growth split
k_α	growth factor (10^{-3} per T^*)
β	phase-field regularisation
r, k	front growth rate, Monod constant
T^*	normalised time, $T^* \in [0, 1]$

Point model (Klempt et al. 2026)

ϕ_i	volume fraction of species i
ϕ_0	void, $\phi_0 = 1 - \sum_i \phi_i$
ψ_i	living fraction (viability) of species i
γ	Lagrange multiplier for $\sum \phi_i + \phi_0 = 1$
A_{ij}	interaction matrix (self, cooperation, competition)
b_i	sensitivity to the antibiotic
c^*, α^*	nutrient, antibiotic concentration
η_i	viscosity (time scale) of species i

Coupling (this work)

ϕ_{3D}	amount from the element field
$\phi_i / \sum_j \phi_j$	composition (share of species i)
s	time link: point model advances $s\,\Delta t$
ϕ_{cap}	largest amount the point model sees

Appendix B: background, growth field and mechanics

Kinematics. Growth is a volume change: $\mathbf{F} = \mathbf{F}_e\mathbf{F}_g$ with isotropic $\mathbf{F}_g = \alpha\mathbf{I}$. Stress comes only from $\mathbf{F}_e$ (neo-Hookean). A body that can grow freely stays stress free; stress appears where growth is constrained.

Material (Klempt et al. 2024, Eq. 20 and Table 2): $E = 10$ Pa, $\nu = 0.49$, stiffness weighted with ϕ^2 (void: floor $10^{-3}E$); no viscosity. Stresses scale linearly with E (the example input's 1000 MPa gives $10^8\times$ larger values).

Field equations (Klempt et al. 2024, from an extended Hamilton principle):

$$\dot{\phi} = \underbrace{\beta\,\Delta\phi}_{\text{diffusion}} + \underbrace{k_\alpha\,\alpha}_{\text{local source}} - \underbrace{r\,\frac{c}{k+c}\,\nabla\phi\cdot\frac{\nabla c}{|\nabla c|}}_{\text{front growth towards nutrient}}\,, \qquad \dot{\alpha} = k_\alpha\,\phi$$

- ϕ and c are nodal fields of the element; α is an internal variable at the Gauss points
- The front term moves the biofilm boundary up the nutrient gradient; it vanishes where ϕ or c is uniform
- The nutrient diffuses much faster than the biofilm grows (quasi-static c)

Appendix C: point model and one substep of the coupled run

Point model (Klempt et al. 2026), per species i, no space dependence, implicit, $\Delta t = 10^{-4}$:


$$(\eta_{\phi,i} + \eta_i \psi_i^2)\,\dot{\phi}_i + \eta_i \phi_i \psi_i\,\dot{\psi}_i = c^* \psi_i \textstyle\sum_j A_{ij}\phi_j\psi_j - \gamma - (\text{barrier})$$

$$\eta_i\,(\phi_i \psi_i\,\dot{\phi}_i + \phi_i^2\,\dot{\psi}_i) = c^* \phi_i \textstyle\sum_j A_{ij}\phi_j\psi_j - \alpha^* b_i\,\psi_i - (\text{barrier}), \qquad \textstyle\sum_i \phi_i + \phi_0 = 1$$

One substep $t_n \to t_{n+1}$: ANSYS solves only $\operatorname{div}\boldsymbol{\sigma} = \mathbf{0}$; the material routine turns $\mathbf{F}$ into $\boldsymbol{\sigma}$ after removing the growth.

repeat (Newton iteration of ANSYS)
  $\mathbf{F}$ at every Gauss point from the current displacements
  for each Gauss point
    $\phi_{3D} = \phi_1 + \phi_2$ from the element; restore the converged state of t_n
    $\alpha_{n+1} = \alpha_n + k_\alpha\,\phi_{3D}\,\Delta t$ Eq. 36
    $\mathbf{F}_e = \mathbf{F}\,\mathbf{F}_g^{-1}$, $\mathbf{F}_g = \alpha_{n+1}\mathbf{I}$ $\to$ $\boldsymbol{\sigma}$, tangent
    if two species and $\phi_{3D} \geq \phi_{\min}$:
      scale the point model to $\min(\phi_{3D}, \phi_{\mathrm{cap}})$
      bridge (Fortran $\to$ C $\to$ socket $\to$ Python): n_{sub} steps of $s\Delta t/n_{\mathrm{sub}}$
      scale back; store ϕ_i, ψ_i, γ as state variables
    return $\boldsymbol{\sigma}$ and the tangent to ANSYS
  assemble the residual, update the displacements
until converged
the element advances ϕ_1, ϕ_2, c to t_{n+1} (its own solver)

Appendix D: Klempt et al. 2024 test cases, reproduced in Python



test case	time scale	RMS ϕ	RMS c
4.1 (Fig. 4)	2	0.03	0.02
4.2 “high” (Fig. 7)	10	0.04	0.04
4.2 “low” (Fig. 7)	4–5	0.09	0.05

Independent solver on the paper's 20 μm cube, not the ANSYS model. Table 2 unchanged; growth on both faces and consumption $g\phi c$ (two departures from Eq. 34/35 as printed); nutrient in a 2 μm corner block; one time scale per run ($T^* = t/t_{\mathrm{ref}}$, Sec. 4). Curves agree; the shape agrees in kind (egg towards the corner) but is more slender, also with other diffusion and a 2nd-order scheme. Table 3's pressure fits μ in MPa.

Appendix E: growth from the living biomass, implementation and runs


$$\ln \alpha_{n+1} = \ln \alpha_n + \kappa\, \phi_{3D} \int_{t_n}^{t_{n+1}} \sum_i \phi_i \psi_i \, dt \qquad \text{(Appendix C, line 5 only)}$$

item	how
form	increment of $\ln \alpha$, as Klempt 2024 Table 1
$\phi_{3D} \in [0, 1]$	only marks where the biofilm is
point model	one per Gauss point, on its own (not scaled to ϕ_{3D})
start	when $\phi_{3D} \geq \phi_{\min}$, from the calibrated Day-1 state
integral	over its own substeps, clock s per unit T^*
storage	state variables of the Gauss point; element unchanged
κ	$8.14 \cdot 10^{-3}$: mean $\alpha - 1$ as Eq. 36 ($-0.003\,\%$); assumed

seed, $T^* = 1$	Eq. 36	living	diff.
$\alpha - 1$, 8^3	$6.752 \cdot 10^{-4}$	$6.752 \cdot 10^{-4}$	0.0 %
von Mises/E, 8^3	$4.684 \cdot 10^{-5}$	$4.357 \cdot 10^{-5}$	$-7.0\,\%$
vM, 16^3; 24^3			-7.1; $-7.0\,\%$
vM/($E(\alpha - 1)$)	0.069	0.064	
antibiotic, 8^3	0 %	$-32\,\%$	


CH, $E = 10$ Pa, $\nu = 0.49$; the antibiotic case with the previous calibration. Same amount of growth, different spread inside the seed; $\alpha - 1$ and stress scale with κ.

Appendix F: how the run is implemented, and the tools



Blue: added in this work; grey: used as it is. One line of the material routine changes compared with Eq. 36.

Appendix G: calibrated five-species model in the element



	start	$T^* = 1$	growth $\alpha - 1$	seed von Mises
shares (Eq. 36)	0.2 each	An 0.38, Vei 0.23, So 0.23, Fn 0.09, Pg 0.08	$6.752 \cdot 10^{-4}$, same as two species	$\mathrm{vM}/E = 4.68 \cdot 10^{-5}$
living biomass $\sum_i \phi_i \psi_i$	0.90	0.44 (mostly So lost)	same mean	$-7.0\,\%$ (8^3), $-7.1\,\%$ (16^3)


commensal flow condition (CH), calibration to the measured data, $E = 10$ Pa, 8^3

Appendix H: two and five species, implementation

	two species (5 October)	five species (since 10 October)
point model	Python, through a material server (Fortran $\rightarrow$ C $\rightarrow$ socket $\rightarrow$ Python)	Fortran, in the same executable (no server, about 4 times faster, same results)
parameters	Klempt et al. 2026 cases 3 and 6 ($c^* = 100$, $\alpha^* = 10$)	calibration (TMCMC, MAP, 5×5 interactions), $c^* = 25$, $\alpha^* = 0$
start	equal shares	equal shares; calibrated Day-1 state for the living-biomass growth
stiffness	partner's example input, $E = 1000$ MPa	Klempt 2024, $E = 10$ Pa
coupling (Eq. 36)	same code: amount scaled to ϕ_{3D}, $\phi_{\mathrm{cap}} = 0.9$, time link $s = 0.15$	
growth from the living biomass	–	yes; point model runs on its own, $s = 0.2375$ ($T^* = 1$ is Day 21)

The number of species is a parameter (1 to 5) of the same routine.

Appendix I: fields on the mid-plane over time




Plane $z = 0$ seen from above (x right, y up), 16^3, CH calibrated to the measured data. Growth by Eq. 36. The phase spreads and thins, the nutrient is used around the seed, and the stress forms a ring at the seed edge.

Appendix J: analysis conditions

geometry, mesh	2 mm cube; 8^3 (512), 16^3, 24^3 hexahedral elements; seed of 32 elements at the centre (8^3, refined with the mesh)
mechanical BC	only rigid-body motion fixed (corner nodes); otherwise traction free
nutrient BC	$c = 1$ on the face $y = -1$ mm; no flux elsewhere
initial state	$\phi = 1$ in the seed, ≈ 0 outside; $\alpha = 1$; $c = 1$ everywhere
time	$T^* = 1$ is Day 21; $\Delta T^* = 0.025$ (40 steps; living-biomass runs: 1/95); ANSYS 2022 R2, static, nonlinear geometry
material	neo-Hooke, $E = 10$ Pa, $\nu = 0.49$, $E \propto \phi^2$ (Klempt 2024 Table 2); $\mathbf{F}_g = \alpha \mathbf{I}$
growth field	$k_\alpha = 10^{-3}$, $\beta = 0.02$ mm$^2/T^*$ (Table 2 scaled to the 2 mm cube, assumed)
point model	two species: Klempt 2026 cases 3 and 6; five species: calibrated MAP (CH), $c^* = 25$
coupling	$\phi_{\mathrm{cap}} = 0.9$, $\phi_{\mathrm{min}} = 0.01$, time link $s = 0.15$ (assumed); living-biomass growth: $s = 0.2375$, $\kappa = 8.14 \cdot 10^{-3}$ (assumed)
output	seed means of $\alpha - 1$ and von Mises$/E$ at $T^* = 1$

Values not from a paper are marked as assumed.

Appendix K: mesh convergence



mesh	elements	nodes	integration points	seed elements
8^3	512	729	4 096	32
16^3	4 096	4 913	32 768	256
24^3	13 824	15 625	110 592	864

8 integration points per hexahedral element ($2 \times 2 \times 2$)

- seed stress: $+23\,\%$ from 8^3 to 16^3, $+3.4\,\%$ from 16^3 to 24^3; converging
- seed growth changes by less than $3.5\,\%$
- difference of the growth laws: -7 to $-8\,\%$ on every mesh
- measured-data calibration: $-7.0\,\%$ at 8^3 and 24^3, $-7.1\,\%$ at 16^3

Appendix L: finite element formulation

balance	$\operatorname{div}\boldsymbol{\sigma} = \mathbf{0}$ in the current configuration, no body force; weak form by ANSYS (static, large deformation)
kinematics	$\mathbf{F} = \mathbf{F}_e\mathbf{F}_g$, $\mathbf{F}_g = \alpha\mathbf{I}$; Cauchy stress from $\mathbf{F}_e$
material	neo-Hooke in $\mathbf{F}_e$, stiffness $(\phi^2 + f)E$ with a small void part f; tangent returned by the material routine
element	SOLID185, 8 nodes, $2 \times 2 \times 2$ integration with B-bar (default), for $\nu = 0.49$
nonlinear solver	full Newton–Raphson, fixed steps (no automatic time stepping), up to 50 iterations, default force tolerance
coupling	staggered: ANSYS solves the mechanics with ϕ fixed; at the end of each step the partner element updates ϕ explicitly and solves the nutrient (linear system, PARDISO) on its neighbour operator
stability	explicit update of ϕ: $\Delta t \leq h^2/(2\beta)$; met for all meshes used
supports	3-2-1: one corner node u_x, u_y, u_z, one u_y, u_z, one u_z (6 dofs, only rigid-body motion)

Time step (one species, 8^3, $T^* = 1.1$): $\Delta T^* = 0.1 \to 0.05 \to 0.025$ changes the seed von Mises/E by $-3.0\,\%$ and $-1.5\,\%$ ($\alpha - 1$: $-0.7\,\%$, $-0.3\,\%$), first order; the runs use $\Delta T^* = 0.025$, the living-biomass runs $1/95 \approx 0.011$.