Coupling a multi-species point model to a continuum biofilm growth model: verification in three codes and the effect of the composition on growth and stress

Authors to be agreed*

Draft of 27 October 2026

Abstract

Continuum models of biofilm growth predict where biomass forms and the stresses it causes, but treat the biofilm as one material. Multi-species point models describe the composition, the living fraction and the response to an antibiotic, but have no space. This work couples the multi-species point model of Klempt et al. (2026), used as published, to the Hamilton-principle growth model of Klempt et al. (2024) at every integration point, without changing the equations of either model. The point model receives the local nutrient and antibiotic from the fields, and its living volume fraction $\sum_i \bar{\phi}_i$ sets the consumption $g\,\phi_{3D}(\sum_i \bar{\phi}_i)\,c$ of both. The coupling is implemented in a finite-volume code, in a neighboured-element user element in ANSYS and in a Galerkin user element in Abaqus, and verified by replay of the point model, a manufactured solution and grid convergence indices. The advection term of the growth equation has a cell Péclet number of about 31 on a 16^3 grid; a constant added diffusion gives a quarter of the converged seed stress, a second-order treatment is needed. With it the growth converges in all examples (GCI 0.7 to 3.3 %) and the seed stress converges when the species start apart (GCI 0.7 to 8 %). An antibiotic removes most of the living biomass of the seed (0.97 to 0.15 when the species start apart, 0.96 to 0.04 when mixed), while the growth changes by about 1 % and the seed stress by about 10 %, because the growth law counts all biomass. The local nutrient decides where the antibiotic acts. The composition that newly reached material receives is not fixed by either model and decides, in a maze, whether the biofilm recovers after the antibiotic.

Keywords: biofilm growth; multi-species model; antibiotic; extended Hamilton principle; code verification; grid convergence index; growth-induced stress

1 Introduction

Oral biofilms on teeth and implants are communities of several bacterial species. Their composition changes over days: early colonisers prepare the surface for later ones, and pathogenic species appear only when the community allows it (Heine et al., 2025). Treatments such as antibiotics act on the composition, while the mechanical load on the surface

*Author list, order and affiliations to be agreed by all contributors before the manuscript is shared; corresponding author to be named.

comes from the amount and the growth of the biomass. A model that predicts the stresses for a given treatment therefore has to describe both: where the biofilm grows, and what it is made of.

Two kinds of model exist for this. Continuum growth models describe the amount of biomass in space. Wanner and Gujer (1986) introduced the one-dimensional multispecies biofilm, Eberl et al. (2001) a density-dependent diffusion model of the spreading. Feng et al. (2021) model a two-species oral biofilm in which the spreading follows the summed growth of the species, and Soleimani et al. (2023) the co-aggregation of two species in a user element. Klempt et al. (2024) derive the growth of a biofilm from the extended Hamilton principle (Junker and Balzani, 2021), with a phase field ϕ_{3D}, a nutrient c and the growth variable α of a multiplicative split $\mathbf{F} = \mathbf{F}_e\mathbf{F}_g$, and compute the resulting stresses; their model has one biomass and no composition. Point models describe the composition of a community at one point. Klempt et al. (2026) derive a multi-species model from the same variational principle, with volume fractions ϕ_i and living fractions ψ_i of N species, an interaction matrix $\mathbf{A}$ and the response of each species to an antibiotic; Fritsch et al. (2025) calibrate its parameters by Bayesian updating. These models have no space: the nutrient c^* and the antibiotic α^* are given inputs.

The point model needs local values of c^* and α^*, which a growth field can supply. The growth field needs to know how much of the biomass is alive, because only living biomass consumes nutrient. Connecting the two at every material point is the natural step, but it raises questions that neither paper answers: which variables are passed, whether either model has to be changed, which new assumptions are needed (for example the composition of biofilm that a point reaches later), and whether the composition reaches the growth and the stress at all.

This work couples the point model of Klempt et al. (2026) to the growth model of Klempt et al. (2024) without changing either set of equations. The contribution is threefold.

1. The point model runs at every integration point as published. It receives $c^* = c_0^* c$ and $\alpha^* = \alpha_0^* a$ from the nutrient and antibiotic fields and returns the living volume fraction $\sum_i \bar{\phi}_i$, which sets the consumption $g\,\phi_{3D}(\sum_i \bar{\phi}_i)\,c$ in the field equations.

2. The coupling is implemented three times, in a finite-volume code, in a neighboured-element user element in ANSYS and in a Galerkin user element in Abaqus, and verified: replay of the point model alone, a manufactured solution, agreement of the codes and grid convergence with the grid convergence index. The advection term $-r\frac{c}{k+c}\,\mathbf{n}_c \cdot \nabla\phi_{3D}$ of the growth equation turns out to set the accuracy of the stress.

3. Examples with two species and an antibiotic, with the synthetic data of Fritsch et al. (2025) and the constants of Klempt et al. (2024, 2026), show what the composition changes and what it does not, and which assumption beyond the two papers matters.

Section 2 states the models and the coupling, Section 3 the implementations and their verification, Section 4 the examples, Section 5 the limits of the coupling, and Section 6 concludes.

2 Models and coupling

2.1 Growth field

The biofilm is described by the phase field $\phi_{3D} \in [0,1]$, the nutrient c and the growth variable α of Klempt et al. (2024), with $\mathbf{F} = \mathbf{F}_e\mathbf{F}_g$, $\mathbf{F}_g = \alpha\mathbf{I}$ and $\alpha(0) = 1$, so that the growth is $\alpha - 1$. After division by the viscosities, their Eqs. 34 to 36 read

$$\dot{\phi_{3D}} = \beta\,\nabla^2\phi_{3D} + k_\alpha\,\alpha - r\,\frac{c}{k+c}\,\mathbf{n}_c\cdot\nabla\phi_{3D}, \qquad \mathbf{n}_c = \frac{\nabla c}{|\nabla c|}, \tag{1}$$

$$\dot{c} = d\,\nabla^2 c - g\,\phi_{3D}\Big(\textstyle\sum_i \bar{\phi}_i\Big)\,c, \tag{2}$$

$$\dot{\alpha} = k_\alpha\,\phi_{3D}, \tag{3}$$

with zero flux on the boundary except where the nutrient is held. The last term of Eq. (1) moves the edge of the biofilm up the nutrient gradient with the speed $r\,c/(k+c)$; we call it the front term. Two details differ from the equations as printed, both confirmed by the authors: the consumption of Eq. (2) is $g\,\phi\,c$ (the printed $g\,\phi$ is a misprint), and the front direction is regularised as $\nabla c/(|\nabla c|^2 + \varepsilon^2)^{1/2}$ with $\varepsilon^2 = 10^{-6}$ ($\varepsilon^2 = 10^{-8}$ changes the seed stress by 0.1 %). The factor $\sum_i \bar{\phi}_i$ in Eq. (2) is the coupling term of this work (Section 2.3); in Klempt et al. (2024) the whole biofilm consumes. The mechanical term $\phi\,\mu\,(\mathbf{I}:\mathbf{C}_e - 3)$ of their Eq. 30, which they neglect because μ is small against the other parameters, is neglected here as well, so the mechanics is coupled one way. The elastic response is neo-Hookean with the stiffness $E(\phi_{3D}) = E\,(\phi_{3D}^2 + f)$, $E = 10$ Pa, $\nu = 0.49$ and a void floor $f = 10^{-3}$ (an assumption that keeps the empty cells solvable; $f \to 0$ changes the seed stress by 0.13 %).

2.2 Point model

At a material point the multi-species model of Klempt et al. (2026) describes the volume fractions ϕ_i and the living fractions ψ_i of N species and the void ϕ_0, with $\sum_i \phi_i + \phi_0 = 1$. It follows from the extended Hamilton principle with the free energy $\Psi = -\frac{1}{2}c^*\,\bar{\boldsymbol{\phi}}^{\mathrm{T}}\mathbf{A}\,\bar{\boldsymbol{\phi}}$, $\bar{\phi}_i = \phi_i\psi_i$, a dissipation with the viscosities η_i, $\eta_{\phi,i}$ and the constraint enforced by the multiplier γ:

$$\begin{aligned} (\eta_{\phi,i} + \eta_i\psi_i^2)\,\dot{\phi}_i + \eta_i\phi_i\psi_i\,\dot{\psi}_i &= c^*\psi_i\textstyle\sum_j A_{ij}\bar{\phi}_j - \gamma - (\text{barrier}),\\ \eta_i\,(\phi_i\psi_i\,\dot{\phi}_i + \phi_i^2\,\dot{\psi}_i) &= c^*\phi_i\textstyle\sum_j A_{ij}\bar{\phi}_j - \alpha^* b_i\,\psi_i - (\text{barrier}), \end{aligned} \tag{4}$$

where A_{ij} are the interactions, b_i the sensitivity of species i to an antibiotic of concentration α^*, c^* the nutrient, and the barrier terms keep ϕ_i, ψ_i in $(0,1)$. The antibiotic lowers ψ_i; the point model compares c^* with $\alpha^* b_i$. Our implementation reproduces the fourteen numerical examples of the paper (two printed details read differently: Eq. 17 as printed carries a $+\gamma$ that the figures do not show, and the four-species interaction matrix of Eq. 19 reproduces the figures without its prefactor $\frac{1}{2}$) and the figures of Fritsch et al. (2025) within 0.013. The model is used as published, with its own constraint and its implicit time integration (steps of 10^{-4} in its own time unit, Newton's method). For the examples the synthetic two-species data of Fritsch et al. (2025), case I, are used with their true parameters: $\mathbf{A} = \left(\begin{smallmatrix} 1 & 0.1 \\ 0.1 & 1 \end{smallmatrix}\right)$, $b = (1,2)$, $\eta = (1,2)$, $c_0^* = 100$.

2.3 Coupling

The two models are connected at every integration point without changing either (Figure 1, Algorithm 1).

1. *Fields to point model.* The point model sees $c^* = c_0^*\, c$ and $\alpha^* = \alpha_0^*\, a$, with a the antibiotic, transported by a second field with the diffusivity and uptake of the nutrient, $\dot{a} = d\,\nabla^2 a - g\,\phi_{3D}(\sum_i \bar{\phi}_i)\, a$. Between the increments t_n and t_{n+1} of the field both are interpolated linearly in time over the inner steps of the point model,

$$c^*(t) = c_0^*\left[\tfrac{t_{n+1}-t}{\Delta t}\, c_n + \tfrac{t-t_n}{\Delta t}\, c_{n+1}\right], \qquad t_n \le t \le t_{n+1}. \tag{5}$$

   Klempt et al. (2026) already use a time-dependent c^* (their four-species case 3), so the point model accepts such an input without change.

2. *Point model to fields.* Only the living biomass consumes: the uptake of nutrient and antibiotic is $g\,\phi_{3D}(\sum_i \bar{\phi}_i)$, Eq. (2). The point model normalises its volume, so $\bar{\phi}_i \in [0,1]$ is the living volume fraction of species i at the point and $\phi_{3D}\sum_i \bar{\phi}_i \le \phi_{3D}$ the living volume fraction in the field; the void ϕ_0 has no living fraction and is not part of the sum. A dead biofilm consumes nothing. The growth law (3) is unchanged: α measures the biomass produced, alive or dead.

3. *Time.* Klempt et al. (2024) normalise time with a reference time chosen for the problem at hand. We take $T^* = 1$ as the duration covered by the point model's data, so that both models run on the same time: the point model advances $s\,\Delta T^*$ per increment, with s the length of that window in its own units, not a fitted value ($s = 0.1$ for case I, 1000 steps of 10^{-4}).

4. *Start of the point model.* It starts at a point once ϕ_{3D} there reaches $\phi_{\min} = 0.001$. Where a seed is placed it starts from the seed composition. A point that the spreading biofilm reaches later takes the living composition of its colonised neighbours (their amounts divided in the ratio of $\bar{\phi}_i$) and starts alive ($\psi_i = 0.999$, $\gamma = 0$), because the biofilm spreads through the growth of its living cells. The papers do not fix this rule; it is an assumption, and the alternative of starting from the seed composition is compared in Section 4.4.

The composition reaches the growth and the stress only through the nutrient field and the front term of Eq. (1): the antibiotic lowers ψ_i, less nutrient is consumed, more reaches the front, and the edge of the biofilm moves differently.

2.4 Parameters

All constants are those of the papers (Table 1): Table 2 of Klempt et al. (2024), converted from their 20 μm cube to the 2 mm cube of the examples, and the antibiotic of case 4 of Klempt et al. (2026), $\alpha_0^* = 100$ from $T^* = 1/3$. With $d = 10^8$ mm$^2/T^*$ the fields adjust within about $4 \times 10^{-8}\, T^*$, so the fields are quasi-steady; keeping the time derivative of Eq. (2) changes the seed values by less than 0.005. A diffusion slower by eight orders of magnitude ($d = 1$ mm$^2/T^*$) is only a sensitivity: the antibiotic then reaches the seed more slowly and kills markedly less of it. Values not given by the papers are marked as assumptions: the length and time scale of the conversion, $\phi_{\min}$ (results change by less than 1 % below 0.001), the start rule of new points, the void floor f and the regularisation ε.




Figure 1: The coupling: the fields ϕ_{3D}, c and a of the growth model, the point model at every integration point with the inputs $c^* = c_0^* c$, $\alpha^* = \alpha_0^* a$ and the output $\sum_i \bar{\phi}_i$, and the mechanics with $\mathbf{F}_g = \alpha \mathbf{I}$.

Algorithm 1 One increment $t_n \to t_{n+1}$ of the coupled run (user-element form)

Require: at every integration point α_n, the point-model state $(\phi_i, \psi_i, \gamma)_n$ and the pair (c, a) seen in the previous increment; the fields ϕ_{3D}, c, a of t_n

Ensure: displacements, stresses and state of t_{n+1}

repeat ▷ Newton iteration of the equilibrium
  for all integration points **do**
    $\alpha_{n+1} \leftarrow \alpha_n + k_\alpha \, \phi_{3D} \, \Delta t$ ▷ Eq. (3)
    $\mathbf{F}_e \leftarrow \mathbf{F} \, \mathbf{F}_g^{-1}$, $\mathbf{F}_g = \alpha_{n+1} \mathbf{I}$; stress and tangent from $W(\mathbf{F}_e)$ and $E(\phi_{3D})$
    if $\phi_{3D} \geq \phi_{\min}$ or the point model has started **then**
      if not yet advanced in this increment **then**
        (c_n, c_{n+1}), (a_n, a_{n+1}) from the previous and the current value at the point
        n_{sub} steps of $s\Delta t / n_{\text{sub}}$ of Eq. (4) with c^*, α^* from Eq. (5)
        store $(\phi_i, \psi_i, \gamma)_{n+1}$ and $\sum_i \bar{\phi}_i$
      end if
    end if
  end for
until the residual meets the convergence criterion
advance ϕ_{3D} (Eq. (1)), c and a (Eq. (2) with $\sum_i \bar{\phi}_i$)

Table 1: Constants of the examples. Conversion: Table 2 of Klempt et al. (2024) on the 2 mm cube with the half edge $\ell = 1$ mm, so that $g\ell^2/d = 1$.

quantity	symbol	value	source
diffusion of ϕ_{3D}	β	0.02 mm$^2/T^*$	Klempt et al. (2024), converted (assumption)
front speed, half saturation	r, k	10 mm/T^*, 1	Klempt et al. (2024), converted
growth rate	k_α	$10^{-3}\,T^{*-1}$	Klempt et al. (2024)
nutrient diffusion, uptake	d, g	10^8 mm$^2/T^*$, $10^8\,T^{*-1}$	Klempt et al. (2024), converted
antibiotic diffusion, uptake	d_a, g_a	$= d$, $= g$	assumption
stiffness, Poisson ratio	E, ν	10 Pa, 0.49	Klempt et al. (2024)
void floor	f	10^{-3}	assumption
point model	$\mathbf{A}$, b, η, c_0^*	case I	Fritsch et al. (2025)
antibiotic level, onset	α_0^*, T_{ab}^*	100, 1/3	Klempt et al. (2026), case 4
time link	s	0.1	from the data of case I
start threshold	$\phi_{\min}$	0.001	assumption (converged)
front regularisation	ε^2	10^{-6}	assumption

Table 2: The three implementations. MLS: moving least squares; TVD: total variation diminishing; SUPG: streamline-upwind Petrov–Galerkin.

	Python	ANSYS (user element)	Abaqus (user element)
ϕ_{3D}, Eq. (1)	finite volumes, RK4	neighboured element method (MLS operators on the Gauss points, 30 neighbours), explicit Euler	Galerkin hexahedra (8 nodes, $2 \times 2 \times 2$ Gauss points), implicit Euler, Newton
c, a, Eq. (2)	7-point Laplacian, sparse direct	MLS Laplacian, direct solver	Galerkin, same Newton system
mechanics	hexahedra, selective reduced integration	hexahedra with the growth material routine	$\bar{\mathbf{B}}$ hexahedra in the same element
front term	upwind (TVD, minmod)	central MLS gradient with stabilisation (Section 3.3)	Galerkin with SUPG
point model	the published model (Klempt et al., 2026), the same Fortran routine in all three		

3 Implementations and verification

3.1 Three codes

The coupled model is solved in three codes that share the equations, the parameters and the compiled point-model routine, but not the discretisation (Table 2). The ANSYS element is the neighboured element method of the original authors' group (Blaszczyk et al., 2022; Rudolf et al., 2025): the Laplacian and the gradients are fitted on the Gauss points from 30 neighbours by a weighted least-squares Taylor expansion, and the fields are updated explicitly. Three numerical details of this element had to be corrected before the results below, without changing its equations (Appendix A.1). Differences between the codes therefore come from the fields and the mechanics only.




Figure 2: Manufactured solution: L_2 errors of (a) ϕ_{3D} and (b) c against the cell size h for the finite-volume (Python) and the Galerkin (Abaqus) code, with a line of slope 2.

3.2 Code verification

Point model in the field. With uniform fields the coupled run reproduces the point model alone to 2×10^{-14} (Python, two and four species) and 1.8×10^{-12} (ANSYS); the hand-over of $\sum_i \bar{\phi}_i$ to the consumption is exact.

Manufactured solution. Sources for ϕ_{3D} and c were built from a manufactured solution (Roache, 2002) by automatic differentiation and checked symbolically (agreement 10^{-14}). The finite-volume and the Galerkin code converge at second order in L_2 (Figure 2): observed orders between 8^3, 16^3 and 32^3 of 1.85 and 1.90 for ϕ_{3D} and 2.03 and 2.01 for c in Python, 1.68 and 1.81 for ϕ_{3D} and 2.00 and 2.05 for c in Abaqus; on 32^3 the errors of ϕ_{3D} are 2.0×10^{-4} and 2.9×10^{-4}. For the SUPG form the order of ϕ_{3D} is about 1.5: the largest error sits in a thin layer next to the wall opposite the supply, where $\nabla c \to \mathbf{0}$ and the front direction changes over a few cells. The analytic tangent of the Abaqus element agrees with central differences to 5×10^{-11}. The neighboured element does not converge next to the walls with the manufactured solution (Appendix A.1).

Mechanics alone. The last growth field of an Abaqus run, solved by the Python mechanics, gives the seed von Mises stress to 1.1 % (no obstacles) and 0.5 % (obstacles with frictionless contact). Differences in the stress between the codes therefore come from the growth field, not from the mechanics.

3.3 The front term

The front term of Eq. (1) is an advection with the speed $r\,c/(k+c)$ along $\mathbf{n}_c$ against a small diffusion: the cell Péclet number $rh/(2\beta)$ is about 31 on a 16^3 grid ($h = 2$ mm$/n$), and a central discretisation oscillates. Four treatments were compared, none of which changes Eqs. (1) to (3):

1. a constant added diffusion $rh/4$ (first order; the element as delivered needs it to stay stable);

2. TVD upwinding (MUSCL with the minmod limiter; Python);

3. SUPG (Brooks and Hughes, 1982) with $\tau = \frac{h}{2|v|}(\coth Pe - 1/Pe)$ on the strong residual (Abaqus);

4. for the explicit element, either the Lax–Wendroff term $\frac{\Delta t}{2}\, v \cdot \nabla(v \cdot \nabla \phi_{3D})$ with an added diffusion switched on by an extremum sensor, or upwind-biased MLS gradients.

Figure 3 shows the seed means of $\alpha - 1$ and of the von Mises stress at $T^* = 1$ of the separate example with antibiotic (Section 4.1) against the grid for every scheme and code. The growth variable converges for every scheme, and the codes agree within their grid-convergence bands: Python extrapolates to 9.0 to 9.8×10^{-4}, Abaqus to 8.8 and 9.45×10^{-4}, the element to 8.7×10^{-4} with upwind gradients. The stress does not converge with a constant added diffusion: $rh/4$ smears the edge of the seed over a few cells, and the stress, which is set by the jump of α across that edge, follows the smearing; on 16^3 it is about a quarter of the converged value (1.3 to 1.7 against 5.9 to 6.3×10^{-4} Pa from the two second-order schemes of two codes), and on 96^3 it is still rising. Treatments 2 to 4 give the same stress within a few per cent from 32^3 on (Python TVD 5.90 and 5.98×10^{-4} Pa on 32^3 and 64^3, Abaqus SUPG 6.27 and 6.22×10^{-4} Pa on 32^3 and 48^3, GCI 1.2 %). The element with upwind gradients gives 2.14, 4.56 and 5.15×10^{-4} Pa on 8^3, 16^3 and 24^3, extrapolated 5.8×10^{-4} Pa (GCI 16.5 %), within the band of the second-order schemes; its MLS operators over 30 neighbours give it a resolution of about one element, not one Gauss-point spacing (with the same added diffusion, the element on 16^3 and Python on 32^3 differ by 1 % in $\alpha - 1$ and 5 % in the stress). For the stress the treatment of the front term matters more than the grid.

3.4 Solution verification

Time step. In the element codes the point model advances in every increment (0.00125 in T^*; Abaqus: observed order 1.1 in time, error against the extrapolated value 0.8 % in $\alpha - 1$ and 2.2 % in the stress). In the Python code the point model and $\sum_i \bar{\phi}_i$ are updated once per outer step ΔT^*, inside which ϕ_{3D}, α and the fields take explicit RK4 steps. The composition and $\alpha - 1$ change by less than 3 and 0.3 % between outer steps of 0.02 and 0.0025; the stress does not: on 64^3 the separate example with antibiotic gives 5.68, 5.82, 5.94 and 6.00×10^{-4} Pa at $\Delta T^* = 0.025$, 0.01, 0.005 and 0.0025 (first-order extrapolation 6.07), the mixed one 3.23, 3.09, 2.90 and 2.82×10^{-4} Pa (2.74). The sensitivity comes from the front speed: after the antibiotic onset $\sum_i \bar{\phi}_i$ is small, ∇c is weak and the front direction responds to small changes of the consumption (Appendix A.2). The grid series of the examples are therefore run at $\Delta T^* = 0.0025$.

Grid. Table 3 gives the grid series of all two-species examples at $\Delta T^* = 0.0025$ with the three-grid index of Celik et al. (2008) (safety factor 1.25), and Figure 4 the same data against the grid. The growth converges with a GCI of 0.7 to 3.3 % in all examples. The seed stress converges in the separate example, with a GCI of 0.7 % without and 8.3 % with the antibiotic. In the mixed examples it converges slowly, with an observed order of about 0.75 and a GCI of 26 to 29 %; with the start fractions of Fritsch et al. (2025) the 128^3 grid is not yet in the asymptotic range. There the front carries intermediate values of ϕ_{3D} at a nearly discontinuous edge, and the largest ϕ_{3D} in the seed still rises with the grid (0.60, 0.69, 0.73). No run on 32^3 to 128^3 left $[0, 1]$ by more than 2×10^{-4}.




Figure 3: Separate example with antibiotic: seed means of (a) $\alpha - 1$ and (b) the von Mises stress at $T^* = 1$ against the grid, one line per scheme and code (dashed: constant added diffusion $rh/4$; solid and dotted: the second-order and the stabilised explicit forms). Grey band: the finest grids of the two second-order schemes. These runs start newly reached points from the seed composition (the alternative rule of Section 4.4) at a field step of 0.00125.

TODO: Fourth grid (192^3, step limit of Appendix A.2) for the mixed example with the start fractions of Fritsch et al.; add to Table 3 and Figure 4 when available.

4 Results

4.1 Set-up

The examples use a 2 mm cube with a cubic seed of $\phi_{3D} = 1$ at its centre, the nutrient held at $c = 1$ on the face $y = -1$ mm and the antibiotic held at $a = 1$ on all outer faces from $T^* = 1/3$. In the separate example species A fills the seed and species B four small blocks along the edges of the cube; in the mixed example both start in the seed, in equal amounts ($\phi_1 = \phi_2 = 0.275$) or in the proportions of Fritsch et al. (2025) (0.25, 0.30). The front term is on in every run. The values below are seed means at $T^* = 1$ from the Python code (TVD, newly reached points inherit the living composition of their neighbours), which reaches the finest grids; the element gives the same composition within its grid error and its growth within 4 % after extrapolation. The stress is solved on 16^3 hexahedra with the element's integration; solving the same 64^3 field on 32^3 and 64^3 elements changes the seed von Mises stress by 0.2 %.

4.2 Composition and living biomass

Table 4 gives the seed means and Figure 5 the fields of the separate and the mixed example with antibiotic. Without antibiotic the living volume fraction of the seed stays near 0.96;

Table 3: Grid convergence (Python, TVD, outer step 0.0025): seed means at $T^* = 1$, observed order p, extrapolated value and GCI (Celik et al., 2008). Mixed: equal amounts unless the start fractions (ϕ_1, ϕ_2) are given; antib.: with antibiotic; –: not yet in the asymptotic range.

example	quantity	32^3	64^3	128^3	p	extrapolated	GCI
separate, antib.	$\alpha - 1$ [10^{-4}]	8.453	8.931	9.112	1.4	9.22	1.5 %
separate, antib.	von Mises [10^{-4} Pa]	5.523	6.000	6.284	0.75	6.70	8.3 %
separate	$\alpha - 1$ [10^{-4}]	8.729	9.014	9.110	1.6	9.16	0.7 %
separate	von Mises [10^{-4} Pa]	7.440	6.730	6.585	2.3	6.55	0.7 %
mixed, antib.	$\alpha - 1$ [10^{-4}]	7.42	8.04	8.32	1.2	8.54	3.3 %
mixed, antib.	von Mises [10^{-4} Pa]	2.00	2.82	3.30	0.76	3.99	26 %
mixed	$\alpha - 1$ [10^{-4}]	7.46	8.08	8.35	1.2	8.56	3.1 %
mixed	von Mises [10^{-4} Pa]	2.13	3.08	3.65	0.74	4.50	29 %
mixed (0.25, 0.30), antib.	$\alpha - 1$ [10^{-4}]	7.42	8.04	8.31	1.2	8.51	3.1 %
mixed (0.25, 0.30), antib.	von Mises [10^{-4} Pa]	2.01	2.96	4.01	–	–	–




Figure 4: Grid convergence of the two-species examples (Table 3): seed means of (a) $\alpha - 1$ and (b) the von Mises stress at $T^* = 1$ against the grid. Dashed: Richardson extrapolation where the three grids are in the asymptotic range.

Table 4: Seed means at $T^* = 1$, Python, 64^3 cells, outer step 0.025; converged growth and stress in Table 3.

example	$\sum_i \bar{\phi}_i$	$\phi_1/(\phi_1+\phi_2)$	c	$\alpha - 1$	von Mises [Pa]
separate	0.965	0.978	0.536	9.06×10^{-4}	6.56×10^{-4}
separate, antibiotic	0.152	0.951	0.911	8.95×10^{-4}	5.68×10^{-4}
mixed, equal amounts	0.961	0.859	0.608	8.33×10^{-4}	3.50×10^{-4}
mixed, equal amounts, antibiotic	0.038	0.610	0.973	8.30×10^{-4}	3.23×10^{-4}
mixed $(0.25, 0.30)$, antibiotic	0.037	0.510	0.973	8.30×10^{-4}	3.22×10^{-4}

the antibiotic lowers it to 0.15 when the species start apart and to 0.04 when they start mixed. Species B, the more sensitive one ($b_2 = 2$), is removed first. The nutrient field decides where the antibiotic acts: the point model compares $c^* = c_0^* c$ with $\alpha^* b_i$, so where the nutrient is high, A survives, and where it is low, A dies as well. When the biomass dies the nutrient is no longer consumed and rises in the seed, from 0.54 to 0.91 in the separate example. Living volume fraction, shares and nutrient change by less than 3 % between the grids (32^3 to 128^3) and the outer steps (0.02 to 0.0025) used.

4.3 Antibiotic over time

With the same total dose given constantly, in pulses or as a sine from $T^* = 1/3$ (Figure 6; separate example, 64^3), $\alpha - 1$ changes by 0.1 % and the seed stress by 0.8 % between the three schedules, while the living volume fraction of the seed follows the dose: in the pauses of the pulsed dose it recovers from 0.17 to up to 0.34 and ends at 0.30, and the sine gives a lower minimum (0.11 against 0.15) and a higher end value (0.19) than the constant dose (0.15). How the dose is given acts on the composition and the living biomass, hardly on the mechanics.

4.4 Which composition does new biofilm receive?

The rule for newly reached points (Section 2.3, item 4) is not fixed by the papers. With the start values of the seed instead of the living composition of the neighbours, the seed values of the cube examples change by a few per cent ($\sum_i \bar{\phi}_i$ 0.147 against 0.152 in the separate example with antibiotic). In the maze of Klempt et al. (2024, Fig. 11), where the biofilm keeps spreading after the antibiotic, the rule decides the outcome (Figure 7, set-up in Section 4.6): the living volume fraction of the colonised region falls to 0.10 shortly after the onset of the antibiotic at $T^* = 20$ and recovers to 0.35 at $T^* = 60$ when new cells take the state of their neighbours, because the biofilm keeps spreading with living cells, and it stays at 0.04 when they start from the start state of the colony. The Galerkin element with the start-state rule gives the same (0.04 to 0.07 in both codes). Without antibiotic the two codes agree in the colonised area within 3 to 5 % and in the living fraction within 0.006.

4.5 What reaches the stress

The growth $\alpha - 1$ hardly depends on the composition: Eq. (3) is unchanged, and the antibiotic lowers ψ_i but not ϕ_{3D} (Table 3: 9.16 against 9.22×10^{-4} in the separate example, about 1 %). The composition can reach the stress through the nutrient and the front term,




Figure 5: Separate (upper half) and mixed (lower half) example with antibiotic, Python, 64^3, one octant removed, at $T^* = 0.325$ (first row of each half) and $T^* = 1$ (second row): (a) share $\phi_1/(\phi_1+\phi_2)$, (b) living volume fraction $\sum_i \bar{\phi}_i$, (c) nutrient c, (d) antibiotic a, (e) von Mises stress. In the mixed example the living volume fraction falls almost everywhere after the antibiotic.




Figure 6: The antibiotic given in three ways with the same total dose from $T^* = 1/3$ (separate example, 64^3): (a) held value at the outer faces, (b) living volume fraction $\sum_i \bar{\phi}_i$ of the seed over time, (c) seed means of $\alpha - 1$ and of the von Mises stress at $T^* = 1$ (each axis spans $\pm 5\,\%$ of its mean).




Figure 7: The maze with the antibiotic held at the entrance from $T^* = 20$, at $T^* = 60$: living volume fraction $\sum_i \bar{\phi}_i$ where $\phi_{3D} \geq 0.05$ when newly reached cells (a) take the state of their colonised neighbours and (b) take the start state of the colony; walls grey, nutrient source outlined in blue, antibiotic source in red. (c) Living volume fraction averaged over the colonised cells against T^* for both rules; the antibiotic starts at the dotted line.

but in these examples the effect of the antibiotic on the converged seed stress is within the numerical uncertainty: 6.55 without and 6.70×10^{-4} Pa with the antibiotic in the separate example (extrapolated, GCI 0.7 and 8.3 %). On a fixed grid the difference is larger (6.73 against 6.00×10^{-4} Pa on 64^3) because the two cases converge from opposite sides. The mixed seed carries about half the stress of the separate one, because its edge is less sharp: the front carries intermediate values of ϕ_{3D}. A limitation follows from Eq. (1): the front speed $r\,c/(k+c)$ depends on the nutrient only, so dead biofilm keeps spreading after the antibiotic; in the maze about two thirds of the cells reached after the antibiotic hold only dead biofilm when new cells start from the start state. Tying the front speed to the living biomass changes the model and is left as outlook (Section 5).

4.6 Geometry: rigid bars and a maze

Klempt et al. (2024) also grow the biofilm against rigid obstacles (their Section 4.3) and through a maze (Section 4.4). Both were repeated with the coupled model in the Python code, without antibiotic unless stated (Figure 8).

Rigid bars. Four bars after Fig. 9 of the paper, with positions read off the sketch: two along x and two along z, each 0.25 mm square, between 0.5 and 0.75 mm from the centre. They hold no biofilm and no nutrient, and the biofilm slides along them without friction. All other settings are those of Section 4.1 (32^3, outer step 0.0025). The biofilm grows around the bars and closes behind them, and by $T^* = 0.2$ it fills the cube between them. The composition of the seed hardly changes. The bars restrain the swelling: the seed von Mises stress is 7.1×10^{-3} Pa against 7.4×10^{-4} Pa without bars in the separate example, about ten times higher, and 9.5×10^{-4} against 2.1×10^{-4} Pa in the mixed one; the growth $\alpha - 1$ of the seed is 9 and 21 % lower. These are values on one grid; on 32^3 the stress without bars lies 14 % above its extrapolated value (Table 3), small against the factor caused by the bars.

Maze. The maze of Fig. 11 of the paper, read off the figure (560 μm across), in two dimensions with 4 μm cells, the constants of Table 6 of the paper ($\beta = 10\ \mu\mathrm{m}^2/T^*$, $r = 200\ \mu\mathrm{m}/T^*$, $k_\alpha = 10^{-6}\,T^{*-1}$), the nutrient held in the centre, the colony of both species at the entrance and the point model of the two-species examples. With the antibiotic, $a = 1$ is held at the entrance from $T^* = 20$. The model is two-dimensional, so it gives no stress. The biofilm follows the corridors, also into dead ends, and approaches the centre: at $T^* = 60$ the nearest point with $\phi_{3D} \geq 0.5$ is 42 μm from the centre without antibiotic and 133 μm with it. The living volume fraction of the colonised region is 0.92 without antibiotic; with it, the rule for new cells decides whether the biofilm recovers (Section 4.4).

4.7 A clinical molar crown

The cube examples place the seed in a box. As a first step towards the geometry of the application, the coupled model was also run on the crown of a human molar from the Open-Full-Jaw dataset (Gholamalizadeh et al., 2022), with the biofilm seeded on the occlusal surface and the nutrient supplied from the surrounding volume (Figure 9). The constants are those of Table 1, and the two species start as in the separate example. The




Figure 8: Upper half: growth against four rigid bars (dark grey) after Fig. 9 of Klempt et al. (2024), Python, 32^3, no antibiotic. (a) Set-up: start biomass (red) and nutrient supply on the face $y = -1$ mm (green). (b)–(d) Surface $\phi_{3D} = 0.5$ (separate example) and 0.2 (mixed example, which starts at $\phi_{3D} = 0.55$) at $T^* = 0.05$, 0.1 and 0.2. Lower half: the maze of Fig. 11 of the same paper with the coupled model (two dimensions, 4 μm cells): ϕ_{3D} at $T^* = 12$, 24, 36 and 60, (a) without and (b) with the antibiotic at the entrance from $T^* = 20$. Walls white, nutrient source outlined in red.




Figure 9: The coupled model on the crown of a human molar (Gholamalizadeh et al., 2022): biofilm ϕ_{3D}, living volume fraction $\sum_i \bar{\phi}_i$, share $\phi_1/(\phi_1+\phi_2)$ and von Mises stress at $T^* = 1$ with the antibiotic from $T^* = 1/3$. Caption to be completed with the figure.

run shows the biofilm following the fissures of the crown, and the living volume fraction, the share and the stress at the end of the antibiotic.

TODO: Section 4.7: run on the molar crown in progress (another script); fill in the set-up (mesh size, seed and sources, time), the seed and fissure values of $\sum_i \bar{\phi}_i$, $\phi_1/(\phi_1+\phi_2)$ and the stress, and check the wording against the figure. Dataset licence CC BY-NC-SA: state it in the data availability section.

5 Discussion

What the coupling shows. The composition reaches the growth field only through the consumption $g\,\phi_{3D}(\sum_i \bar{\phi}_i)\,c$. When the antibiotic kills the biomass, less nutrient is consumed, the nutrient rises in the seed and the front term moves the edge differently. The effect on the growth is small, about 1 %, because the growth law $\dot{\alpha} = k_\alpha \phi_{3D}$ counts all biomass, living or dead; the effect on the seed stress is about 10 %, because the stress is set by the jump of α across the edge of the seed, which the front shapes. The local nutrient decides where the antibiotic acts: the same dose removes a species in one part of the biofilm and not in another, which a model without space cannot show. The stress depends on how sharp the edge of the biofilm is, and that edge is set by the front term, so the coupling needs a second-order treatment of that term; a constant added diffusion gives a quarter of the converged seed stress on 16^3 and does not converge up to 96^3.

Limitations. (i) The front speed depends on the nutrient only, so dead biofilm keeps spreading after the antibiotic. Tying the speed to the living biomass is a change of the model; in a test on 16^3 (front speed multiplied by the local living fraction) the seed stress rose by 51 % in the separate example and about fourfold in the mixed one, so this choice matters for the stress. (ii) Eq. (3) lets dead biomass grow at the same rate as living biomass; growth from the living biomass, as in Wanner and Gujer (1986), would make the antibiotic act on the stress directly. (iii) The point model has no space; the composition moves only through the rule for newly reached points, which decides in the maze whether the biofilm recovers. A species-resolved continuum would replace the rule. (iv) The mechanics is coupled one way, as in Klempt et al. (2024); with $E = 10$ Pa the neglected term acts only for coefficients far beyond the paper values (an effect on the growth from about 10^5 on, 16^3 test). (v) Living and dead biomass, and both species, have

the same stiffness; a species-dependent stiffness changes the seed stress by 9 to 89 % in a 16^3 test, depending on which species is stiffer. (vi) The examples use synthetic data with known parameters, the stress converges slowly when the species start mixed, and the element's values carry a grid error of 16.5 %. Calibrated data (for example the five-species data of Heine et al., 2025) and a measured stress are needed before the stresses can be validated, not only verified.

Outlook. Species-resolved transport, so that the composition moves with the biomass; front speed and growth tied to the living biomass, with the consequences for the stress quantified above; growth laws with a mechanical feedback; calibrated multi-species data in the element; and a realistic tooth or implant geometry. Each can be added to the coupling as presented here, and the verification procedure, several codes, a manufactured solution and grid convergence indices, carries over.

6 Conclusions

This work coupled the multi-species point model of Klempt et al. (2026) to the continuum growth model of Klempt et al. (2024) at every integration point, without changing the equations of either model. The point model receives the local nutrient and antibiotic from the fields, $c^* = c_0^* c$ and $\alpha^* = \alpha_0^* a$, and its living volume fraction sets the uptake $g\,\phi_{3D}(\sum_i \bar{\phi}_i)\,c$ of both.

1. The coupling was implemented in a finite-volume code, in the neighboured-element user element of the original authors' group in ANSYS and in a Galerkin user element in Abaqus. The point model inside each code reproduces the point model alone to round-off in uniform fields, the finite-volume and the Galerkin code converge at second order for a manufactured solution, and the three codes agree within their grid-convergence bands. Three numerical details of the neighboured element had to be corrected for this.

2. The front term $-r\frac{c}{k+c}\,\mathbf{n}_c \cdot \nabla\phi_{3D}$ sets the accuracy of the stress. A constant added diffusion keeps it stable but gives a quarter of the converged seed stress; a second-order upwind treatment is needed. With it, the growth converges with a GCI of 0.7 to 3.3 % in all examples, the seed stress with 0.7 to 8.3 % when the species start apart and 26 to 29 % when they start mixed.

3. An antibiotic removes most of the living biomass of the seed (from 0.97 to 0.15 when the species start apart, from 0.96 to 0.04 when mixed), while the growth changes by about 1 % and the seed stress by about 10 %: the antibiotic lowers the consumption, more nutrient reaches the front, and the edge of the seed changes. The local nutrient decides where the antibiotic acts, because the point model weighs c^* against $\alpha^* b_i$.

4. The composition of biofilm that reaches a new point is not fixed by either model. Taking the living composition of the neighbours or the start state gives the same cube results within a few per cent, but decides in a maze whether the biofilm recovers after an antibiotic (living volume fraction 0.35 against 0.04).

5. Rigid obstacles restrain the swelling and raise the seed stress about tenfold; a maze makes the biofilm follow its corridors and approach the nutrient source.

The composition reaches growth and stress only through the nutrient and the front speed, because the growth law counts all biomass. The next steps are models in which the front speed, the growth or the stiffness depend on the living biomass or on the species.

Data availability

The finite-volume code, the point-model routine, the run scripts, the result files of the examples and the figure scripts are in the public repository `https://github.com/keisuke58/pde-fem-biofilm`. The examples use the published synthetic data of Fritsch et al. (2025); no calibration data are used in this work. The user elements depend on the commercial codes and on the element of Rudolf et al. (2025); the material routines are in the repository.

TODO: Confirm with all authors which parts of the element code may be published, and whether a release (DOI) of the repository is made at submission.

Declarations

Author list, contributions, funding and conflicts of interest: to be agreed.

TODO: Authors, affiliations, contributions, funding (LUH/IKM, Keio), acknowledgements; apply the Springer Nature template (sn-jnl.cls) of Biomechanics and Modeling in Mechanobiology at submission.

A Numerical details

A.1 The neighboured element

Three details of the element were corrected, without changing its equations: the surface points take the value of the nearest Gauss point instead of a distance-weighted mean, the Laplacian is used in a conservative form, and the neighbours of the fit are chosen by distance instead of by element number. Before, ϕ_{3D} leaked through the walls of the cube (the surface points took an inverse-square mean, and the column sums of the Laplacian were not zero), and the stencils of the 16^3 and 24^3 grids were asymmetric inside the domain. With a manufactured solution the corrected element still does not converge from 8^3 to 16^3 (L_2 error of ϕ_{3D} 2.2×10^{-3} to 1.8×10^{-3}, of c 8.2×10^{-3} to 1.2×10^{-2}): the MLS stencils of the Gauss points next to a face include the surface points and are ill-conditioned (error of the Laplacian of the manufactured c: 0.88 on 8^3, 65 on 16^3, against $|\nabla^2 c| = 3.5$). Rebuilding these stencils with the neighbours mirrored across the faces (zero flux built in) makes the operators consistent up to the walls, and with the weight width proportional to h the element converges (ϕ_{3D} 9.5×10^{-4} to 3.4×10^{-4}, order 1.5; c 9.0×10^{-4} to 2.2×10^{-4}, order 2.0). In the coupled example the mirrored operators run on 8^3 but become unstable at a corner of the 16^3 grid after the antibiotic onset; they are not used for the results. The explicit update of the element is limited by $\beta \Delta t / h^2 \leq 0.12$ to 0.16 on 16^3 and 24^3 (Bloch analysis of the stencil); the Lax–Wendroff form with the extremum sensor keeps ϕ_{3D} in $[-0.03, 1.01]$ on 64^3 in Python but is unstable at the corners of the supply face on 24^3 in the element, so the element's stresses are quoted from the upwind MLS gradients. The time-step series of the element and the stability analysis will be given in the supplementary material.

A.2 Outer step, regularisation and stability

Within an outer step of the Python code, the inner RK4 step, a trapezoidal update of $\sum_i \bar{\phi}_i$ over the step and letting newly reached points consume at once each change the stress by less than 1 %; the one-increment lag of $\sum_i \bar{\phi}_i$ and the regularisation of the front direction ($\varepsilon^2 = 10^{-6}$ against 10^{-8}, 0.1 %) are not the cause of the step sensitivity of Section 3.4. A larger regularisation ($\varepsilon^2 = 10^{-4}$) changes the mixed stress by 30 %, which shows that the front direction, not the speed, carries the sensitivity. Without antibiotic the outer step changes the stress by 2.5 % (separate) and 12 % (mixed). The explicit inner step is limited by the diffusion number $D = \beta \Delta t / h^2 \leq 0.15$ and the Courant number of the front $Co \leq 0.4$; on 192^3 both limits are close at the same time and the step diverged in the first outer step, which the combined condition $12D + 2Co \leq 2.2$ removes without changing the step on 32^3 to 128^3. The regularised front direction and the Péclet number of the examples are unchanged by these choices.

Supplementary material

The time-step series of the element and of the Galerkin code, the stability analysis of the explicit element, the sensitivities to $\phi_{\min}$, d and the start rule, and the animation of the maze will be provided as supplementary material.

References

Mischa Blaszczyk, Dustin Roman Jantos, and Philipp Junker. Application of Taylor series combined with the weighted least square method to thermodynamic topology optimization. *Computer Methods in Applied Mechanics and Engineering*, 393:114698, 2022. doi: 10.1016/j.cma.2022.114698.

Alexander N. Brooks and Thomas J. R. Hughes. Streamline upwind/Petrov–Galerkin formulations for convection dominated flows with particular emphasis on the incompressible Navier–Stokes equations. *Computer Methods in Applied Mechanics and Engineering*, 32(1–3):199–259, 1982. doi: 10.1016/0045-7825(82)90071-8.

Ishmail B. Celik, Urmila Ghia, Patrick J. Roache, Christopher J. Freitas, Hugh Coleman, and Peter E. Raad. Procedure for estimation and reporting of uncertainty due to discretization in CFD applications. *Journal of Fluids Engineering*, 130(7):078001, 2008. doi: 10.1115/1.2960953.

Hermann J. Eberl, David F. Parker, and Mark C. M. van Loosdrecht. A new deterministic spatio-temporal continuum model for biofilm development. *Journal of Theoretical Medicine*, 3 (3):161–175, 2001. doi: 10.1080/10273660108833072.

Dianlei Feng, Insa Neuweiler, Regina Nogueira, and Udo Nackenhorst. Modeling of symbiotic bacterial biofilm growth with an example of the Streptococcus–Veillonella sp. system. *Bulletin of Mathematical Biology*, 83:48, 2021. doi: 10.1007/s11538-021-00888-2.

Lukas Fritsch, Hendrik Geisler, Jan Grashorn, Felix Klempt, Meisam Soleimani, Matteo Broggi, Philipp Junker, and Michael Beer. Bayesian updating of constitutive parameters under hybrid uncertainties with a novel surrogate model applied to biofilms. arXiv:2512.15145, 2025. Preprint.

Torkan Gholamalizadeh, Faezeh Moshfeghifar, Zachary Ferguson, Teseo Schneider, Daniele Panozzo, Sune Darkner, Masrour Makaremi, François Chan, Peter Lampel Søndergaard, and Kenny Erleben. Open-Full-Jaw: An open-access dataset and pipeline for finite element models of human jaw. *Computer Methods and Programs in Biomedicine*, 224:107009, 2022. doi: 10.1016/j.cmpb.2022.107009.

Nils Heine, Kristina Bittroff, Szymon P. Szafrański, Maya Duitscher, Wiebke Behrens, Clarissa Vollmer, Carina Mikolai, Nadine Kommerein, Nicolas Debener, Katharina Frings, Alexander Heisterkamp, Thomas Scheper, Maria L. Torres-Mapa, Janina Bahnemann, Meike Stiesch, and Katharina Doll-Nikutta. Influence of species composition and cultivation condition on peri-implant biofilm dysbiosis in vitro. *Frontiers in Oral Health*, 6:1649419, 2025. doi: 10.3389/froh.2025.1649419.

Philipp Junker and Daniel Balzani. An extended Hamilton principle as unifying theory for coupled problems and dissipative microstructure evolution. *Continuum Mechanics and Thermodynamics*, 33(4):1931–1956, 2021. doi: 10.1007/s00161-021-01017-z.

Felix Klempt, Meisam Soleimani, Peter Wriggers, and Philipp Junker. A Hamilton principle-based model for diffusion-driven biofilm growth. *Biomechanics and Modeling in Mechanobiology*, 23 (6):2091–2113, 2024. doi: 10.1007/s10237-024-01883-x.

Felix Klempt, Hendrik Geisler, Meisam Soleimani, and Philipp Junker. A continuum multi-species bacterial growth model with a novel interaction scheme. *Archive of Applied Mechanics*, 96:164, 2026. doi: 10.1007/s00419-026-03160-y.

Patrick J. Roache. Code verification by the method of manufactured solutions. *Journal of Fluids Engineering*, 124(1):4–10, 2002. doi: 10.1115/1.1436090.

Tobias Rudolf, Felix Klempt, Hüray Ilayda Kök, Meisam Soleimani, Dustin Roman Jantos, and Philipp Junker. Computational efficiency and accuracy of the Neighbored Element Method. *Finite Elements in Analysis and Design*, 249:104353, 2025. doi: 10.1016/j.finel.2025.104353.

Meisam Soleimani, Szymon P. Szafranski, Taoran Qu, Rumjhum Mukherjee, Meike Stiesch, Peter Wriggers, and Philipp Junker. Numerical and experimental investigation of multi-species bacterial co-aggregation. *Scientific Reports*, 13:11839, 2023. doi: 10.1038/s41598-023-38806-2.

Oskar Wanner and Willi Gujer. A multispecies biofilm model. *Biotechnology and Bioengineering*, 28(3):314–328, 1986. doi: 10.1002/bit.260280304.