Coupling the Point Model to the Growth Field

Key points of the meeting of 20 October 2026

Keisuke Nishioka


Institute of Continuum Mechanics (IKM), Leibniz Universität Hannover

20 October 2026

What changes

- Point model (Klempt et al. 2026) used as published
- No scaling to the field ($\phi_0 = 1 - \phi_{3D}$ dropped)
- Field $\rightarrow$ point model: c^*, α^* from 3D fields
- Point model $\rightarrow$ field: only living biomass consumes
- Eq. 34 with front term, Eq. 36 unchanged

The loop at one Gauss point




$\sum_i \bar{\phi}_i$ scales the consumption of c and a $\rightarrow$ c enters the front term of Eq. 34 $\rightarrow$ spreading of ϕ_{3D} $\rightarrow$ growth α (Eq. 36) $\rightarrow$ stress

Equations

$$
\begin{aligned}
c^*(t) &= c_0^*\left[\tfrac{t_{n+1}-t}{\Delta t}c_n + \tfrac{t-t_n}{\Delta t}c_{n+1}\right], \quad \alpha^* \text{ likewise} \\
0 &= d\,\nabla^2 c - g\,\phi_{3D}\,\textstyle\sum_i \bar{\phi}_i\, c, \qquad \bar{\phi}_i = \phi_i\psi_i \\
0 &= d_a\nabla^2 a - g_a\,\phi_{3D}\,\textstyle\sum_i \bar{\phi}_i\, a \\
\dot{\phi}_{3D} &= \beta\nabla^2\phi_{3D} + k_\alpha\alpha - \|\nabla\phi_{3D}\|\,\tfrac{r\,c}{k+c}\,\mathbf{n}_{\nabla\phi_{3D}}\cdot\mathbf{n}_{\nabla c}, \qquad \dot{\alpha} = k_\alpha\phi_{3D}
\end{aligned}
$$


$g\phi$ in Eq. 35 of the paper is a misprint; $g\phi_{3D}c$ is used. ϕ_{3D}: field, ϕ_i: point model.

Nutrient and antibiotic: from the nodes to the Gauss points



- As strains from displacements: the shape functions at the Gauss point
- The antibiotic in the second nutrient field takes the same path
- One increment behind: the field is solved after the material routine

First ANSYS run: what is checked



- (a) c_{n+1} of one increment is c_n of the next: the pair is passed on without a jump
- (b) point model in ANSYS = Python with the same inputs, 10^{-12} or better from the second increment
- First increment: no nutrient yet ($c = 0$); start from the steady field instead
- To do: linear field $c = (y + 1)/2$, values at each Gauss point (node → Gauss point)

Time: one clock for both models

- $T^* = 1$ = duration of the point model's data
- s follows from it, not fitted (synthetic: $s = 0.1$)
- Check (synthetic): error 7×10^{-7}; with scaling 0.019

Examples and checks

Examples (synthetic, two species)

- A centre, B edges
- A and B mixed
- with / without antibiotic

Checks

- equals point model alone
- ANSYS = Python
- $0 \leq \phi \leq 1$
- ϕ_{min}, step, mesh

Results to follow

Assumptions and open points

- Assumptions: linear interpolation, $\sum_i \bar{\phi}_i$ one step late, start values of new points, ϕ_{min}
- No transport of species between points
- Front term needs upwinding
- Antibiotic: synthetic data only ($b = 0$ in the calibration)

Appendix: notation

ϕ_{3D}, α	biofilm phase field (ϕ of Klempt et al. 2024) and growth ($\mathbf{F}_g = \alpha\mathbf{I}$, $\alpha(0) = 1$), Klempt et al
c, a	nutrient and antibiotic fields (the element's first and second nutrient)
d, g, d_a, g_a	diffusion and consumption of nutrient and antibiotic
β, k_α, r, k	diffusion of ϕ, growth rate, front growth and Monod constant (Eq. 34, 36)
ϕ_i, ψ_i, ϕ_0	volume fraction and living fraction of species i, auxiliary void (Klempt et al. 2026)
c^*, α^*, b_i	nutrient and antibiotic of the point model, antibiotic sensitivity
$\bar{\phi}_i$	living volume fraction of species i, $\phi_i\psi_i$ (Klempt et al. 2026)
T^*, s	normalised time; point-model time per unit T^*