

## 1 Plan for Class

1. Stability of an equilibrium point
2. Lyapunov functions
3. Linearization about an equilibrium

## 2 Example from last time

$$\begin{aligned}\dot{x} &= y \\ \dot{y} &= x - x^2 + \epsilon[\alpha y + \beta xy]\end{aligned}$$

If  $\epsilon = 0$ , system is Hamiltonian:

$$H(x, y) = \frac{y^2}{2} - \frac{x^2}{2} + \frac{x^3}{3}$$

Note that  $H$  is constant along trajectories. In a Hamiltonian system you follow level sets.

What if  $\epsilon > 0$ ? If  $\alpha, \beta < 0$ , then  $H$  always decreases.

$$\begin{aligned}\frac{d}{dt}H(x(t), y(t)) &= \frac{\partial H}{\partial x}\dot{x} + \frac{\partial H}{\partial y}\dot{y} \\ &= \epsilon y^2[\alpha + \beta x] < 0\end{aligned}$$

when  $x < 0$  given  $\alpha, \beta < 0$ . Visually, this makes the point  $(x, y) = (1, 0)$  a sink.

This example is nice because it allows us to develop and test a bunch of tools that classify if we get homoclinic orbits, limit cycles, sinks, sources, etc. In general, this is a nice system to see all the phenomenon one expects in dynamical systems.

## 3 Stability of an equilibrium point

Three classes of stability:

1. Asymptotically stable (perturbations get asymptotically closer to the fixed point)
2. Lyapunov stable (orbit around the fixed point)
3. Unstable stable (perturbations get arbitrarily far away)

### 3.1 Equilibrium point

For  $\dot{x} = f(x)$ , the equilibrium point is simply

$$f(x) = 0 \iff x_0 \text{ eq. pt}$$

or also called a fixed point. In other words,  $x_0$  is stable if "when you start close, you stay close" (also called "Lyapunov stable"). More formally,  $\forall \epsilon > 0, \exists \delta > 0$  such that  $\|x(t) - x_0\| < \epsilon$  whenever  $\|x(0) - x_0\| < \delta$  for all  $t > 0$ . In plain english, if you start in an epsilon ball, you stay in the epsilon ball.

$x_0$  is asymptotically stable if it is stable and in addition  $x(t) \rightarrow x_0$  as  $t \rightarrow \infty$ . Centers are an example of  $x_0$  being stable but not asymptotically stable. Sinks are example of asymptotically stable.

**Unstable:**  $x_0$  is unstable if it is not stable.

#### 3.1.1 Tricky example

$$\dot{\theta} = \sin \frac{\theta}{2}$$

$\theta_0 = 0$  is a fixed point, and things will always decay to it, but it is unstable because it violates our  $\epsilon$  ball definition of stability. However the following example

$$\dot{\theta} = \sin \theta$$

has a fixed point at  $\theta_0 = \pi$  that is asymptotically stable.

#### 3.1.2 Hamiltonian example

The origin is unstable because there exists at least one perturbation that blows up to infinity. The point  $(x, y) = (1, 0)$  is asymptotically stable if  $\epsilon = 0$ .

## 4 Lyapunov functions

Lyapunov functions are a way to show an equilibrium point is stable in some region  $U \subset \mathbb{R}^n$ . Say we have a  $C^1$  function  $V(x)$  such that:

- $V(x_0) = 0$
- $V(x) > 0$  for all  $x \neq x_0$
- $\dot{V}(x) \leq 0$  for  $x \neq x_0$  then  $x_0$  is stable
- Further, if  $\dot{V}(x) < 0$  for  $x \neq x_0$  then  $x_0$  is asymptotically stable

where the first two conditions imply that  $V$  is "positive definite".

Here  $\dot{V}(x)$  means

$$\frac{d}{dt} V(x(t)) = \sum_{i=1}^n \frac{\partial V}{\partial x_i} \dot{x}_i = \sum_{i=1}^n \frac{\partial V}{\partial x_i} f_i(x) = f(x) \cdot \nabla V(x)$$

## 4.1 Proof

Say we have such a function  $V$  such that  $\dot{V} < 0$ . Let  $\eta$  be the minimum value of  $V(x)$  on<sup>1</sup> the sphere of radius  $\epsilon$  about  $x_0$ . We know that the minimum value exists from analysis because  $V$  is a continuous function on a compact set, so it must have a maximum and minimum. We want to find a  $\delta$  to complete our proof.

Because  $V$  is continuous, and  $V(x_0) = 0$ ,  $\exists \delta > 0$  such that  $|V(x) - V(x_0)| < \eta$  where  $\|x - x_0\| < \delta$ .

So  $V(x) < \eta$  for all  $x \in B_\delta$  (where this is an open ball radius  $\delta$  about  $x_0$ ) and  $V(x)$  decreases along a trajectory  $x(t)$ . So  $x(t)$  can never leave  $B_\epsilon$ . If it did, it would need to intersect the sphere of radius  $\epsilon$ , and on that sphere,  $V(y) \geq \eta$ , which is a contradiction.

To get asymptotically stable, since  $V(x)$  strictly decreases,  $V(x(t))$  must go to 0 as  $t \rightarrow \infty$ .

## 4.2 Simple pendulum example

From Newton, we have

$$ml^2\ddot{\theta} = -mgl \sin \theta - \mu \dot{\theta}$$

where we can rewrite it as

$$\begin{aligned}\dot{\theta} &= \omega \\ \dot{\omega} &= \frac{1}{ml^2} (-mgl \sin \theta - \mu \omega)\end{aligned}$$

We can imagine the phase space as a cylinder.

Let's prove that  $(0, 0)$  is stable for  $\mu \geq 0$ . We will give a Lyapunov from the energy

$$V(\theta, \omega) = \frac{1}{2}ml^2\omega^2 + mgl(1 - \cos \theta)$$

Let's make sure that  $V(0, 0) = 0$ , and it does. Let's make sure that it is positive definite. Indeed it is since  $\omega^2 > 0$  and  $1 - \cos \theta$  when  $\theta, \omega \in [-\pi, \pi]$  excluding 0.

Is it decreasing?

$$\begin{aligned}\dot{V}(\theta, \omega) &= \frac{\partial V}{\partial \theta} \dot{\theta} + \frac{\partial V}{\partial \omega} \dot{\omega} \\ &= (mgl \sin \theta) \omega + \omega [-mgl \sin \theta - \mu \omega] \\ &= -\mu \omega^2 \leq 0\end{aligned}$$

for all  $\omega, \theta \neq 0$ . This proves stability, and to prove asymptotic stability, we can use La Salle's invariance principle to conclude that starting at  $\omega = 0$  for any  $\theta \neq 0$  is not invariant, so is not stable.

---

<sup>1</sup>Where "on the sphere" means that it is on the shell of the sphere, not inside.

## 5 Linearization about an equilibrium

Say that for  $\dot{x} = f(x)$ , we have an equilibrium point  $x_0$  such that  $f(x_0) = 0$ . Via the Taylor series, we can write

$$\dot{x} = f(x) = f(x_0) + Df(x_0) \cdot (x - x_0) + \text{Higher order terms}$$

For Linearization, we ignore the higher order terms which may or may not be justified. Now we get the linearized system  $\dot{\xi} = A \xi$  that is

$$\dot{\xi} = Df(x_0) \cdot \xi$$

where  $Df(x_0)$  is an  $n \times n$  matrix.

Let  $A = Df(x_0)$ . Many properties of this linear system are determined by eigenvalues of  $A$ .

$$Av = \lambda v$$

where  $\lambda$  is an eigenvalue and  $v$  is an eigenvector. To verify, we can write

$$\xi(t) = c_1 e^{\lambda_1 t} v_1 + \cdots + c_n e^{\lambda_n t} v_n$$

$$A\xi(t) = c_1 e^{\lambda_1 t} \lambda_1 v_1 + \cdots + c_n e^{\lambda_n t} \lambda_n v_n = \dot{\xi}(t)$$

If  $\lambda_1, \dots, \lambda_n$  all have real part strictly negative,  $\xi(t) \rightarrow 0$  as  $t \rightarrow \infty$ , which is asymptotically stable. If any of  $\lambda_1, \dots, \lambda_n$  have real part  $> 0$ , then we get something unstable.

### 5.1 Linearization theorem

(Special case of Hartman-Grobman theorem) Let  $x_0$  be an equilibrium point for  $\dot{x} = f(x)$ , and let  $A = Df(x_0)$ .

- If all eigenvalues of  $A$  have strictly negative real parts, then  $x_0$  is asymptotically stable.
- If any eigenvalues of  $A$  has strictly positive real parts, then  $x_0$  is unstable.
- Otherwise, inconclusive.

We could get cases where the real part is zero, but the imaginary parts are nonzero.

#### 5.1.1 Proof sketch

Suppose  $A$  "has only eigenvalues with negative real part" (also called Hurwitz). Construct a Lyapunov function.

Result from linear systems: If  $A$  is Hurwitz, then  $A^T P + PA = -Q$  for any positive definite symmetric  $Q$ , solution  $P$  is positive definite symmetric.

$$\text{Let } V(\xi) = \xi^T P \xi =$$