

Listing of Trials contained in the Centralized Open-Access Rehabilitation database for Stroke (SCOAR).

Keith Lohse, PhD

School of Kinesiology, Auburn University

e | rehabinformatics@gmail.com

*As of 2016-04-15, the database contains 215 independent references (223 total) consisting of 489 independent groups of participants (which equates to 12,847 total participants).

**Note that citations are listed in the order they appear in the SCOAR database, which is roughly alphabetical except for follow-up studies/secondary analyses (i.e., published later in a separate manuscript) which are listed immediately after the original publication.

References

1. Abdullah HA, Tarry C, Lambert C, Barreca S, Allen BO. Results of clinicians using a therapeutic robotic system in an inpatient stroke rehabilitation unit. *J Neuroengineering Rehabil.* 2011;8:50.
2. Ada L, Dean CM, Hall JM, et al. A treadmill and overground walking program improves walking in persons residing in the community after stroke: a placebo-controlled, randomized trial. *Arch Phys Med Rehabil.* 2003;84(October):1486-1491. doi:10.1053/S0003-9993(03)00349-6.
3. Ada L, Dean CM, Lindley R. Randomized trial of treadmill training to improve walking in community-dwelling people after stroke: the AMBULATE trial. *Int J Stroke.* 2013;8(6):436-444.
4. Aisen ML, Krebs HI, Hogan N, et al. The effect of robotassisted therapy and rehabilitative training on motor recovery following stroke. *Arch Neurol.* 1997;54(4):443-446. doi:10.1001/archneur.1997.00550160075019.
5. Volpe B, Krebs H, Hogan N, Edelsteinn L. Robot training enhanced motor outcome in patients with stroke maintained over 3 years. *Neurology.* 1999.
6. Arya KN, Verma R, Garg RK, Sharma VP, Agarwal M, Aggarwal GG. Meaningful Task-Specific Training (MTST) for Stroke Rehabilitation: A Randomized Controlled Trial. *Top Stroke Rehabil.* 2012;19(3):193-211.
7. Bale M, Strand LI. Does functional strength training of the leg in subacute stroke improve physical performance? A pilot randomized controlled trial. *Clin Rehabil.* 2008;22(10-11):911-921.
8. Barbeau H, Visintin M. Optimal outcomes obtained with body-weight support combined with treadmill training in stroke subjects. *Arch Phys Med Rehabil.* 2003;84(10):1458.
9. Blennerhassett J, Dite W. Additional task-related practice improves mobility and upper limb function early after stroke: a randomised controlled trial. *Aust J Physiother.* 2004;50(4):219-224.
10. Boake C, Noser EA, Ro T, et al. Constraint-induced movement therapy during early stroke rehabilitation. *Neurorehabil Neural Repair.* 2007;21(1):14-24. doi:10.1177/1545968306291858.
11. Bolognini N, Vallar G, Casati C, et al. Neurophysiological and behavioral effects of tDCS combined with constraint-induced movement therapy in poststroke patients. *Neurorehabilitation & Neural Repair.* 2011;25(9):819-829.
12. Brock K, Haase G, Rothacher G, Cotton S. Does physiotherapy based on the Bobath concept, in conjunction with a task practice, achieve greater improvement in walking ability in people with stroke compared to physiotherapy focused on structured task practice alone? *Clin Rehabilitation.* 2011;25(10):903-912. doi:10.1177/0269215511406557.
13. Broeren J, Claesson L, Goude D, Rydmark M, Sunnerhagen KS. Virtual rehabilitation in an activity centre for community-dwelling persons with stroke. The possibilities of 3-dimensional computer games. *Cerebrovasc Dis.* 2008;26(3):289-296. doi:10.1159/000149576.
14. Brokaw EB, Nichols D, Holley RJ, Lum PS. Robotic therapy provides a stimulus for upper limb motor recovery after stroke that is complementary to and distinct from conventional therapy. *Neurorehabil Neural Repair.* 2014;28(4):367-376.
15. Brunner IC, Skouen JS, Strand LI. Is modified constraint-induced movement therapy more effective than bimanual training in improving arm motor function in the subacute phase post

- stroke? A randomized controlled trial. *Clin Rehabil*. 2012;26(12):1078-1086.
16. Burgar C, Lum PS, Scremin AME, et al. Robot-assisted upper-limb therapy in acute rehabilitation setting following stroke: Department of Veterans Affairs multisite clinical trial. *J Rehabil Res Dev*. 2011;48(4):445-458.
 17. Carmeli E, Peleg S, Bartur G, Elbo E, Vatine J. HandTutor(TM) enhanced hand rehabilitation after stroke - a pilot study. *Physiother Res Int*. 2010;16:191-200.
 18. Cauraugh JH, Kim SB. Chronic stroke motor recovery: Duration of active neuromuscular stimulation. *J Neurol Sci*. 2003;215(1-2):13-19.
 19. Chan MK, Tong RK, Chung KY. Bilateral upper limb training with functional electric stimulation in patients with chronic stroke. *Neurorehabilitation & Neural Repair*. 2009;23(4):357-365.
 20. Chen JC, Lin CH, Wei YC, Hsiao J, Liang CC. Facilitation of motor and balance recovery by thermal intervention for the paretic lower limb of acute stroke: a single-blind randomized clinical trial. *Clin Rehabil*. 2011;25(9):823-832.
 21. Cho KH, Lee WH. Virtual walking training program using a real-world video recording for patients with chronic stroke: a pilot study. *Am J Phys Med Rehabil*. 2013;92(5):371-380. doi:10.1097/PHM.0b013e31828cd5d3.
 22. Conroy SS, Whittall J, Dipietro L, et al. Effect of gravity on robot-assisted motor training after chronic stroke: a randomized trial. *Arch Phys Med Rehabil*. 2011;92(11):1754-1761.
 23. Cooke E V, Tallis RC, Clark A, Pomeroy VM. Efficacy of functional strength training on restoration of lower-limb motor function early after stroke: phase I randomized controlled trial. *Neurorehabilitation & Neural Repair*. 2010;24(1):88-96. doi:10.1177/1545968309343216.
 24. Crosbie JH, Lennon S, McGoldrick MC, McNeill MDJ, McDonough SM. Virtual reality in the rehabilitation of the arm after hemiplegic stroke: a randomized controlled pilot study. *Clin Rehabil*. 2012;26(9):798-806. doi:10.1177/0269215511434575.
 25. da Silva Cameirão M, Bermúdez I Badia S, Duarte E, et al. Virtual reality based rehabilitation speeds up functional recovery of the upper extremities after stroke: a randomized controlled pilot study in the acute phase of stroke using the rehabilitation gaming system. *Restor Neurol Neurosci*. 2011;29(5):287-298. doi:10.3233/RNN-2011-0599.
 26. Dahl a E, Askim T, Stock R, et al. Short- and long-term outcome of constraint-induced movement therapy after stroke: a randomized controlled feasibility trial. *Clin Rehabil*. 2008;22(5):436-447. doi:10.1177/0269215507084581.
 27. Daly JJ, Hogan N, Perepezko EM, et al. Response to upper-limb robotics and functional neuromuscular stimulation following stroke. *J Rehabil Res Dev*. 2005;42(6):723-736.
 28. Danzl MM, Chelette KC, Lee K, Lykins D, Sawaki L. Brain stimulation paired with novel locomotor training with robotic gait orthosis in chronic stroke: A feasibility study. *NeuroRehabilitation*. 2013;33(1):67-76.
 29. de Diego C, Puig S, Navarro X. A sensorimotor stimulation program for rehabilitation of chronic stroke patients. *Restor Neurol Neurosci*. 2013;31(4):361-371.
 30. Dean CM, Richards CL, Malouin F. Task-related circuit training improves performance of

- locomotor tasks in chronic stroke: a randomized, controlled pilot trial. *Arch Phys Med Rehabil*. 2000;81(4):409-417. doi:10.1053/mr.2000.3839.
31. Dean CM, Ada L, Bampton J, Morris ME, Katrak PH, Potts S. Treadmill walking with body weight support in subacute non-ambulatory stroke improves walking capacity more than overground walking: a randomised trial. *J Physiother*. 2010;56(2):97. doi:10.1016/S1836-9553(10)70039-4.
 32. Ada L, Dean C, Morris D, Simpson J, Katrak PH. Randomized trial of treadmill walking with body weight support to establish walking in subacute stroke: the MOBILISE trial. *Stroke*. 2010;41:1237.
 33. Desrosiers J, Bourbonnais D, Corriveau H, et al. Effectiveness of unilateral and symmetrical bilateral task training for arm during the subacute phase after stroke: A randomized controlled trial. *Clin Rehabil*. 2005;19(6):581-593. doi:10.1191/0269215505cr896oa.
 34. Di Lauro A, Pellegrino L, Savastano G, et al. A randomized trial on the efficacy of intensive rehabilitation in the acute phase of ischemic stroke. *J Neurol*. 2003;250(10):1206-1208. doi:10.1007/s00415-003-0185-2.
 35. Dias D, Lains J, Pereira A, Nunes R, Caldas J. Can we improve gait skills in chronic hemiplegics? A randomised control trial with gait trainer. *Eura Medicophys*. 2007;43:499.
 36. Dobkin BH, Plummer-D'Amato P, Elashoff R, Lee J, SIRROWS G. International randomized clinical trial, stroke inpatient rehabilitation with reinforcement of walking speed (SIRROWS), improves outcomes. *Neurorehabil Neural Repair*. 2010;24(3):235.
 37. Donaldson C, Tallis R, Miller S, Sunderland A, Lemon R, Pomeroy V. Effects of conventional physical therapy and functional strength training on upper limb motor recovery after stroke: a randomized phase II study. *Neurorehabilitation & Neural Repair*. 2009;23(4):389-397. doi:10.1177/1545968308326635.
 38. Dromerick a. W, Edwards DF, Hahn M. Does the application of constraint-induced movement therapy during acute rehabilitation reduce arm impairment after ischemic stroke?. *Stroke*. 2000;31(12):2984-2988. doi:10.1161/01.STR.31.12.2984.
 39. Dromerick AW, Lang CE, Birkenmeier RL, et al. Very Early Constraint-Induced Movement during Stroke Rehabilitation (VECTORS): a single-center RCT. *Neurology*. 2009;73(3):195-201.
 40. Duncan P, Richards L, Wallace D, et al. A randomized, controlled pilot study of a home-based exercise program for individuals with mild and moderate stroke. *Stroke*. 1998;29(10):2055-2060. doi:10.1161/01.STR.29.10.2055.
 41. Duncan P, Studenski S, Richards L, et al. Randomized clinical trial of therapeutic exercise in subacute stroke. *Stroke*. 2003;34(9):2173-2180. doi:10.1161/01.STR.0000083699.95351.F2.
 42. Duncan PW, Sullivan KJ, Behrman AL, et al. Body-weight-supported treadmill rehabilitation after stroke. *N Engl J Med*. 2011;364(21):2026. doi:10.1056/NEJMoa1010790.
 43. Eich H-J, Mach H, Werner C, Hesse S. Aerobic treadmill plus Bobath walking training improves walking in subacute stroke: a randomized controlled trial. *Clin Rehabil*. 2004;18(6):640. doi:10.1191/0269215504cr779oa.
 44. Fang Y, Chen X, Li H, Lin J, Huang R, Zeng J. A study on additional early physiotherapy after stroke and factors affecting functional recovery. *Clin Rehabil*. 2003;17(6):608-617.

<http://www.ncbi.nlm.nih.gov/pubmed/12971705>.

45. Feys HM, De Weerdts WJ, Selz BE, et al. Effect of a therapeutic intervention for the hemiplegic upper limb in the acute phase after stroke: a single-blind, randomized, controlled multicenter trial. *Stroke*. 1998;29(4):785-792. doi:10.1161/01.STR.29.4.785.
46. Fisher S, Lucas L, Thrasher TA. Robot-assisted gait training for patients with hemiparesis due to stroke. *Top Stroke Rehabil*. 2011;18(3):269-276. doi:10.1310/tsr1803-269.
47. Franceschini M, Carda S, Agosti M, et al. Walking after stroke: what does treadmill training with body weight support add to overground gait training in patients early after stroke?: a single-blind, randomized, controlled trial. *Stroke*. 2009;40(9):3079-3085. doi:10.1161/STROKEAHA.109.555540.
48. Fuzaro AC, Guerreiro CT, Galetti FC, Jucá RBVM, Araujo JE De. Modified constraint-induced movement therapy and modified forced-use therapy for stroke patients are both effective to promote balance and gait improvements. *Rev Bras Fisioter*. 2012;16(2):157-165.
49. Galvin R, Cusack T, O'Grady E, Murphy TB, Stokes E. Family-mediated exercise intervention (FAME): evaluation of a novel form of exercise delivery after stroke. *Stroke*. 2011;42(3):681-686.
50. Glasgow Augmented Physiotherapy Study. Can augmented physiotherapy input enhance recovery of mobility after stroke? A randomized controlled trial. *Clin Rehabil*. 2004;18(5):529-537. doi:10.1191/0269215504cr768oa.
51. Gelber D a., Josefczyk B, Herrman D, et al. Comparison of two therapy approaches in the rehabilitation of the pure motor hemiparetic stroke patient. *Neurorehabilitation & Neural Repair*. 1995;9(4):191-196. doi:10.1177/154596839500900401.
52. Gilbertson L, Langhorne P, Walker A, Allen A, Murray GD. Domiciliary occupational therapy for patients with stroke discharged from hospital: randomised controlled trial. *Br Med J*. 2000;320(7235):603-606.
53. Gladman JR, Lincoln NB, Barer DH. A randomised controlled trial of domiciliary and hospital-based rehabilitation for stroke patients after discharge from hospital. *J Neurol Neurosurg Psychiatry*. 1993;56(9):960-966.
54. Gladman JR, Lincoln NB. Follow-up of a controlled trial of domiciliary stroke rehabilitation (DOMINO Study). *Age Ageing*. 1994;23(1):9-13. <http://www.ncbi.nlm.nih.gov/pubmed/8010180>.
55. Globas C, Becker C, Cerny J, et al. Chronic stroke survivors benefit from high-intensity aerobic treadmill exercise: a randomized control trial. *Neurorehabil Neural Repair*. 2012;26(1):85-95. doi:10.1177/1545968311418675.
56. Green J, Forster A, Bogle S, Young J. Physiotherapy for patients with mobility problems more than 1 year after stroke: a randomised controlled trial. *Lancet*. 2002;359(9302):199-203. doi:10.1016/S0140-6736(02)07443-3.
57. Han C, Wang Q, Meng PP, Qi MZ. Effects of intensity of arm training on hemiplegic upper extremity motor recovery in stroke patients: a randomized controlled trial. *Clin Rehabil*. 2013;27(1):75-81.
58. Harris JE, Eng JJ, Miller WC, Dawson AS. A self-administered Graded Repetitive Arm

- Supplementary Program (GRASP) improves arm function during inpatient stroke rehabilitation: a multi-site randomized controlled trial. *Stroke*. 2009;40(6):2123-2128.
59. Hayner K, Gibson G, Giles GM. Comparison of constraint-induced movement therapy and bilateral treatment of equal intensity in people with chronic upper-extremity dysfunction after cerebrovascular accident. *Am J Phys Med Rehabil*. 2010;64(4):528-539. doi:10.5014/ajot.2010.08027.
 60. Hesse S, Werner C, Pohl M, Rueckriem S, Mehrholz J, Lingnau ML. Computerized arm training improves the motor control of the severely affected arm after stroke: a single-blinded randomized trial in two centers. *Stroke*. 2005;36(9):1960-1966. doi:10.1161/01.STR.0000177865.37334.ce.
 61. Hesse S, Welz a, Werner C, Quentin B, Wissel J. Comparison of an intermittent high-intensity vs continuous low-intensity physiotherapy service over 12 months in community-dwelling people with stroke: a randomized trial. *Clin Rehabil*. 2011;25(2):146-156. doi:10.1177/0269215510382148.
 62. Hidler J, Nichols D, Pelliccio M, et al. Multicenter randomized clinical trial evaluating the effectiveness of the Lokomat in subacute stroke. *Neurorehabilitation & Neural Repair*. 2009;23(1):5-13. doi:10.1177/1545968308326632.
 63. Hornby TG, Campbell DD, Kahn JH, Demott T, Moore JL, Roth HR. Enhanced gait-related improvements after therapist- versus robotic-assisted locomotor training in subjects with chronic stroke: a randomized controlled study. *Stroke*. 2008;39(6):1786-1792. doi:10.1161/STROKEAHA.107.504779.
 64. Housman SJ, Scott KM, Reinkensmeyer DJ. A randomized controlled trial of gravity-supported, computer-enhanced arm exercise for individuals with severe hemiparesis. *Neurorehabilitation & Neural Repair*. 2009;23(5):505-514. doi:10.1177/1545968308331148.
 65. Hsieh Y, Wu C, Liao W, et al. Effects of treatment intensity in upper limb robot-assisted therapy for chronic stroke: a pilot randomized controlled trial. *Neurorehabil Neural Repair*. 2011;25(6):503-511. doi:10.1177/1545968310394871.
 66. Hsieh Y, Wu C, Lin K, Yao G, Wu K, Chang Y. Dose-response relationship of robot-assisted stroke motor rehabilitation: the impact of initial motor status. *Stroke*. 2012;43(10):2729-2734. doi:10.1161/STROKEAHA.112.658807.
 67. Hui-Chan CWY, Ng SSM, Mak MKY. Effectiveness of a home-based rehabilitation programme on lower limb functions after stroke. *Hong Kong Med J*. 2009;15(3 Suppl 4):42-46. <http://www.ncbi.nlm.nih.gov/pubmed/19509438>.
 68. Hunter SK, Hammett L, Ball S, Smith N, Anderson C. Dose-response study of mobilisation and tactile stimulation therapy for the upper extremity early after stroke: a phase I trial. *Neurorehabilitation & Neural Repair*. 2011;25:314-322.
 69. Husemann B, Müller F, Krewer C, et al. Effects of locomotion training with assistance of a robot-driven gait orthosis in hemiparetic patients after stroke: a randomized controlled pilot study. *Stroke*. 2007;38(349):349-354. doi:10.1161/01.STR.0000254607.48765.cb.
 70. In TS, Jung KS, Lee SW, et al. Virtual Reality Reflection Therapy Improves Motor Recovery and Motor Function in the Upper Extremities of People with Chronic Stroke. *J Phys Ther Sci*.

- 2012;24(4):339-343. doi:10.1589/jpts.24.339.
71. Indredavik B, Bakke F, Solberg R, Rokseth R, Haaheim LL, Holme I. Benefit of a stroke unit: a randomized controlled trial. *Stroke*. 1991;22(8):1026-1031. doi:10.1161/01.STR.22.8.1026.
 72. Janssen TW, Beltman JM, Elich P, et al. Effects of electric stimulation-assisted cycling training in people with chronic stroke. *Arch Phys Med Rehabil*. 2008;89(3):463-469.
 73. Jin H, Jiang Y, Wei Q, Wang B, Ma G. Intensive aerobic cycling training with lower limb weights in Chinese patients with chronic stroke: discordance between improved cardiovascular fitness and walking ability. *Disabil Rehabil*. 2012;34(19):1665-1671. doi:10.3109/09638288.2012.658952.
 74. Jin H, Jiang Y, Wei Q, Chen L, Ma G. Effects of aerobic cycling training on cardiovascular fitness and heart rate recovery in patients with chronic stroke. *NeuroRehabilitation*. 2013;32(2):327-335.
 75. Johannsen L, Wing AM, Pelton T, et al. Seated bilateral leg exercise effects on hemiparetic lower extremity function in chronic stroke. *Neurorehabilitation & Neural Repair*. 2010;24(3):243-253.
 76. Jongbloed L, Stacey S, Brighton C. Stroke rehabilitation: sensorimotor integrative treatment versus functional treatment. *Am J Occup Ther*. 1989;43(6):391-397.
 77. Jonsdottir J, Cattaneo D, Recalcati M, et al. Task-oriented biofeedback to improve gait in individuals with chronic stroke: motor learning approach. *Neurorehabil Neural Repair*. 2010;24(5):478-485. doi:10.1177/1545968309355986.
 78. Jung J. Effects of virtual-reality treadmill training on balance and balance self efficacy in stroke patients with a history of falling. *J Phys Ther Sci*. 2012;24(5):1133-1136.
 79. Kahn LE, Zygmant ML, Rymer WZ, Reinkensmeyer DJ. Robot-assisted reaching exercise promotes arm movement recovery in chronic hemiparetic stroke: a randomized controlled pilot study. *J Neuroengineering Rehabil*. 2006;3:12.
 80. Kang H-K, Kim Y, Chung Y, Hwang S. Effects of treadmill training with optic flow on balance and gait in individuals following stroke: randomized controlled trials. *Clin Rehabil*. 2012;26(3):246-255. doi:10.1177/0269215511419383.
 81. Katz-Leurer M, Sender I, Keren O, et al. The influence of early cycling training on balance in stroke patients at the subacute stage. Results of a preliminary trial. *Clin Rehabil*. 2006;20(5):398-405.
 82. Khan CM, Oesch PR, Gamper UN, Kool JP, Beer S. Potential effectiveness of three different treatment approaches to improve minimal to moderate arm and hand function after stroke--a pilot randomized clinical trial. *Clin Rehabil*. 2011;25(11):1032-1041.
 83. Kihoon J. Effects of virtual reality-based rehabilitation on upper extremity function and visual perception in stroke patients: A randomized control trial. *J Phys Ther Sci*. 2012;24:1205-1208.
 84. Kim JH, Jang SH, Kim CS, Jung JH, You JH. Use of virtual reality to enhance balance and ambulation in chronic stroke: a double-blind, randomized controlled study. *Am J Phys Med Rehabil*. 2009;88(9):693-701. doi:10.1097/PHM.0b013e3181b33350.
 85. Kim E, Kang J, Park J, Jung B. Clinical feasibility of interactive commercial Nintendo gaming for chronic stroke rehabilitation. *J Phys Ther Sci*. 2012;24:901-903.
 86. Kiper P, Piron L, Turolla A, Stozek J, Tonin P. The effectiveness of reinforced feedback in virtual

- environment in the first 12 months after stroke. *Neurol Neurochir Pol.* 2011;45(5):436-444.
87. Kuys SS, Brauer SG, Ada L. Higher-intensity treadmill walking during rehabilitation after stroke in feasible and not detrimental to walking pattern or quality: a pilot randomized trial. *Clin Rehabil.* 2011;25(4):316-326. doi:10.1177/0269215510382928.
 88. Kwakkel G, Wagenaar RC, Twisk JW, Lankhorst GJ, Koetsier JC. Intensity of leg and arm training after primary middle-cerebral-artery stroke: a randomised trial. *Lancet.* 1999;354(9174):191-196.
 89. Kwakkel G, Kollen BJ, Wagenaar RC. Long term effects of intensity of upper and lower limb training after stroke: a randomised trial. *J Neurol Neurosurg Psychiatry.* 2002;72(4):473-479.
 90. Kwon J-S, Park M-J, Yoon I-J, Park S-H. Effects of virtual reality on upper extremity function and activities of daily living performance in acute stroke: a double-blind randomized clinical trial. *NeuroRehabilitation.* 2012;31(4):379-385. doi:10.3233/NRE-2012-00807.
 91. Langhammer B, Lindmark B, Stanghelle JK. Stroke patients and long-term training: is it worthwhile? A randomized comparison of two different training strategies after rehabilitation. *Clin Rehabil.* 2007;21(6):495-510.
 92. Langhammer B, Stanghelle J. Bobath or motor relearning programme? A comparison of two different approaches of physiotherapy in stroke rehabilitation: a randomized controlled study. *Clin Rehabil.* 2000;14:361-369.
 93. Langhammer B, Stanghelle JK, Programme P, Stanghelle JK, Rehabilitation S. Exercise on a treadmill or walking outdoors? A randomized controlled trial comparing effectiveness of two walking exercise programmes late after stroke. *Clin Rehabil.* 2010;24:46-54.
 94. Lau K, Mak M. Speed-dependent treadmill training is effective to improve gait and balance performance in patients with sub-acute stroke. *J Rehabil Med.* 2011;43:709-713.
 95. Laufer Y, Dickstein R, Chefez Y, Marcovitz E. The effect of treadmill training on the ambulation of stroke survivors in the early stages of rehabilitation: a randomized study. *J Rehabil Res Dev.* 2001;38(1):69-78.
 96. Laver K, George S, Ratcliffe J, et al. Use of an interactive video gaming program compared with conventional physiotherapy for hospitalised older adults: a feasibility trial. *Disabil Rehabil.* 2012;(January):1-7. doi:10.3109/09638288.2012.662570.
 97. Lee MJ, Kilbreath SL, Singh MF, et al. Comparison of effect of aerobic cycle training and progressive resistance training on walking ability after stroke: a randomized sham exercise-controlled study. *J Am Geriatr Soc.* 2008;56(6):976-985. doi:10.1111/j.1532-5415.2008.01707.x.
 98. Lee M, Cho H, Song C. The mirror therapy program enhances upper-limb motor recovery and motor function in acute stroke patients. *Am J Phys Med Rehabil.* 2012;6:689-699.
 99. Lin K-C-C, Wu C-Y-Y, Wei T-H-H, Lee C-Y-Y, Liu J-S-S. Effects of modified constraint-induced movement therapy on reach-to-grasp movements and functional performance after chronic stroke: A randomized controlled study. *Clin Rehabil.* 2007;21(12):1075-1086. doi:10.1177/0269215507079843.
 100. Lin K, Chen Y, Chen C, Wu C, Chang Y. The effects of bilateral arm training on motor control and functional performance in chronic stroke: a randomized controlled study. *Neurorehabilitation &*

- Neural Repair*. 2010;24(1):42-51. doi:10.1177/1545968309345268.
101. Lincoln NB, Parry RH, Vass CD. Randomized, controlled trial to evaluate increased intensity of physiotherapy treatment of arm function after stroke. *Stroke*. 1999;30(3):573-579. doi:10.1161/01.STR.30.3.573.
 102. Lo A, Guarino P, Richards L, Haselkorn J, Wittenberg G. Robot-assisted therapy for long-term upper-limb impairment after stroke. *N Eng J Med*. 2010;362:1772-1783.
 103. Logan P, Ahern J, Gladman J, Lincoln N. A randomized controlled trial of enhanced Social Service occupational therapy for stroke patients. *Clin Rehabilitation*. 1997;11(2):107-113. doi:10.1177/026921559701100203.
 104. Luft AR, Macko RF, Forrester LW, et al. Treadmill exercise activates subcortical neural networks and improves walking after stroke: a randomized controlled trial. *Stroke*. 2008;39(12):3341-3350. doi:10.1161/STROKEAHA.108.527531.
 105. Lum PS, Burgar CG, Shor PC, Majmundar M, Van der Loos HFM, Van der Loos M. Robot-assisted movement training compared with conventional therapy techniques for the rehabilitation of upper-limb motor function after stroke. *Arch Phys Med Rehabil*. 2002;83(7):952-959. doi:10.1053/apmr.2001.33101.
 106. Macko RF, Ivey FM, Forrester LW, et al. Treadmill exercise rehabilitation improves ambulatory function and cardiovascular fitness in patients with chronic stroke: a randomized, controlled trial. *Stroke*. 2005;36(10):2206-2211. doi:10.1161/01.STR.0000181076.91805.89.
 107. Masiero S, Celia A, Rosati G, Armani M. Robotic-assisted rehabilitation of the upper limb after acute stroke. *Arch Phys Med Rehabil*. 2007;88(2):142-149. doi:10.1016/j.apmr.2006.10.032.
 108. Masiero S, Armani M, Ferlini G, Rosati G, Rossi A. Randomized trial of a robotic assistive device for the upper extremity during early inpatient stroke rehabilitation. *Neurorehabilitation & Neural Repair*. 2014;28(4):377-386. doi:http://dx.doi.org/10.1177/1545968313513073.
 109. McClellan R, Ada L. A six-week, resource-efficient mobility program after discharge from rehabilitation improves standing in people affected by stroke: placebo-controlled, randomised trial. *Aust J Physiother*. 2004;50:163.
 110. Michielsen ME, Selles RW, van der Geest JN, et al. Motor recovery and cortical reorganization after mirror therapy in chronic stroke patients: a phase II randomized controlled trial. *Neurorehabilitation & Neural Repair*. 2011;25(3):223-233. doi:10.1177/1545968310385127.
 111. Mirelman A, Bonato P, Deutsch JE. Effects of training with a robot-virtual reality system compared with a robot alone on the gait of individuals after stroke. *Stroke*. 2009;40(1):169-174. doi:10.1161/STROKEAHA.108.516328.
 112. Moore JL, Roth EJ, Killian C, Hornby TG. Locomotor training improves daily stepping activity and gait efficiency in individuals poststroke who have reached a "plateau" in recovery. *Stroke*. 2010;41(1):129-135. doi:10.1161/STROKEAHA.109.563247.
 113. Moreland JD, Goldsmith CH, Huijbregts MP, et al. Progressive resistance strengthening exercises after stroke: a single-blind randomized controlled trial. *Arch Phys Med Rehabil*. 2003;84(10):1433-1440.

114. Morone G, Bragoni M, Iosa M, et al. Who may benefit from robotic-assisted gait training? A randomized clinical trial in patients with subacute stroke. *Neurorehabilitation & Neural Repair*. 2011;25(7):636-644. doi:10.1177/1545968311401034.
115. Morone G, Tramontano M, Iosa M, et al. The efficacy of balance training with video game-based therapy in subacute stroke patients: A randomized controlled trial. *Biomed Res Int*. 2014;580861. doi:10.1155/2014/580861.
116. Morris JH, Van Wijck F, Joice S, Ogston SA, Cole I, MacWalter RS. A comparison of bilateral and unilateral upper-limb task training in early poststroke rehabilitation: a randomized controlled trial. *Arch Phys Med Rehabil*. 2008;89(7):1237-1245. doi:10.1016/j.apmr.2007.11.039.
117. Mudge S, Barber PA, Stott NS. Circuit-based rehabilitation improves gait endurance but not usual walking activity in chronic stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2009;90(12):1989-1996. doi:10.1016/j.apmr.2009.07.015.
118. Ng SS, Hui-Chan CW. Transcutaneous electrical nerve stimulation combined with task-related training improves lower limb functions in subjects with chronic stroke. *Stroke*. 2007;38(11):2953-2959.
119. Ng MF, Tong RK, Li LS. A pilot study of randomized clinical controlled trial of gait training in subacute stroke patients with partial body-weight support electromechanical gait trainer and functional electrical stimulation: six-month follow-up. *Stroke*. 2008;39(1):154-160.
120. Nilsson L, Fugl-Meyer KH, Kristensen L, Sjölund B, Sunnerhagen KS. Walking training of patients with hemiparesis at an early stage after stroke: a comparison of walking training on a treadmill with body weight support and walking training on the ground. 2001;15:515-527.
121. Noh D, Lim J, Shin H, Paik N. The effect of aquatic therapy on postural balance and muscle strength in stroke survivors—a randomized controlled pilot trial. *Clin Rehabil*. 2008;22:966-976.
122. Olawale OA, Jaja SI, Anigbogu CN, Appiah-Kubi KO, Jones-Okai D. Exercise training improves walking function in an African group of stroke survivors: a randomized controlled trial. *Clin Rehabil*. 2011;25(5):442-450.
123. Olney SJ, Nymark J, Brouwer B, et al. A randomized controlled trial of supervised versus unsupervised exercise programs for ambulatory stroke survivors. *Stroke*. 2006;37(2):476-481. doi:10.1161/01.STR.0000199061.85897.b7.
124. Ouellette MM, LeBrasseur NK, Bean JF, et al. High-intensity resistance training improves muscle strength, self-reported function, and disability in long-term stroke survivors. *Stroke*. 2004;35(6):1404-1409. doi:10.1161/01.STR.0000127785.73065.34.
125. Outermans JC, van Peppen RPS, Wittink H, Takken T, Kwakkel G. Effects of a high-intensity task-oriented training on gait performance early after stroke: a pilot study. *Clin Rehabil*. 2010;24(11):979-987. doi:10.1177/0269215509360647.
126. Page SJ, Sisto S, Levine P, McGrath RE. Efficacy of Modified Constraint-Induced Movement Therapy in Chronic Stroke: A Single-Blinded Randomized Controlled Trial. *Arch Phys Med Rehabil*. 2004;85(1):14-18. doi:10.1016/S0003-9993(03)00481-7.
127. Page SJ, Levine P, Leonard AC. Modified constraint-induced therapy in acute stroke: a randomized controlled pilot study. *Neurorehabil Neural Repair*. 2005;19(1):27-32.

doi:10.1177/1545968304272701.

128. Page SJ, Levine P, Leonard A, Szaflarski JP, Kissela BM. Modified constraint-induced therapy in chronic stroke: Results of a single-blinded randomized controlled trial. *Phys Ther.* 2008;88(3):333-340. doi:10.2522/ptj.20060029.
129. Pandian S, Arya KN, Davidson EW. Comparison of Brunnstrom movement therapy and Motor Relearning Program in rehabilitation of post-stroke hemiparetic hand: a randomized trial. *J Bodyw Mov Ther.* 2012;16(3):330-337.
130. Pang MY, Eng JJ, Dawson AS, McKay HA, Harris JE. A community-based fitness and mobility exercise program for older adults with chronic stroke: a randomized, controlled trial. *J Am Geriatr Soc.* 2005;53(10):1667-1674.
131. Pang MY, Harris JE, Eng JJ. A community-based upper-extremity group exercise program improves motor function and performance of functional activities in chronic stroke: a randomized controlled trial. *Arch Phys Med Rehabil.* 2006;87:1-9.
132. Park HJ, Oh DW, Kim SY, Choi JD. Effectiveness of community-based ambulation training for walking function of post-stroke hemiparesis: a randomized controlled pilot trial. *Clin Rehabil.* 2011;25(5):451-459.
133. Park I-M, Lee Y-S, Moon B-M, Sim S-M. A Comparison of the Effects of Overground Gait Training and Treadmill Gait Training According to Stroke Patients' Gait Velocity. *J Phys Ther Sci.* 2013;25(4):379-382.
134. Parker CJ, Gladman JRF, Drummond a. ER, et al. A multicentre randomized controlled trial of leisure therapy and conventional occupational therapy after stroke. *Clin Rehabil.* 2001;15(1):42-52. doi:10.1191/026921501666968247.
135. Parry R, Lincoln N, Vass C. Effect of severity of arm impairment on response to additional physiotherapy early after stroke. *Clin Rehabil.* 1999.
136. Partridge C, Mackenzie M, Edwards S, et al. Is dosage of physiotherapy a critical factor in deciding patterns of recovery from stroke: a pragmatic randomized controlled trial. *Physiother Res Int.* 2000;5(4):230-240.
137. Patten C, Condliffe EG, Dairaghi CA, Lum PS. Concurrent neuromechanical and functional gains following upper-extremity power training post-stroke. *J Neuroengineering Rehabil.* 2013;10:1.
138. Peurala SH, Tarkka IM, Pitkanen K, Sivenius J, Pitkänen K, Sivenius J. The effectiveness of body weight-supported gait training and floor walking in patients with chronic stroke. *Arch Phys Med Rehabil.* 2005;86(8):1557-1564. doi:10.1016/j.apmr.2005.02.005.
139. Piron L, Tombolini P, Turolla A, et al. Reinforced Feedback in Virtual Environment Facilitates the Arm Motor Recovery in Patients after a Recent Stroke. *2007 Virtual Rehabil.* 2007:121-123. doi:10.1109/ICVR.2007.4362151.
140. Piron L, Turolla A, Agostini M, et al. Exercises for paretic upper limb after stroke: a combined virtual-reality and telemedicine approach. *J Rehabil Med.* 2009;41(12):1016-1102. doi:10.2340/16501977-0459.
141. Piron L, Turolla A, Agostini M, et al. Motor learning principles for rehabilitation: a pilot

- randomized controlled study in poststroke patients. *Neurorehabilitation & Neural Repair*. 2010;24(6):501-508. doi:10.1177/1545968310362672.
142. Platz T, Eickhof C, Van Kaick S, et al. Impairment-oriented training or Bobath therapy for severe arm paresis after stroke: a single-blind, multicentre randomized controlled trial. *Clin Rehabil*. 2005;19(7):714-724. doi:10.1191/0269215505cr904oa.
 143. Platz T, van Kaick S, Mehrholz J, Leidner O, Eickhof C, Pohl M. Best conventional therapy versus modular impairment-oriented training for arm paresis after stroke: a single-blind, multicenter randomized controlled trial. *Neurorehabil Neural Repair*. 2009;23(7):706-716. doi:10.1177/1545968309335974.
 144. Pohl M, Mehrholz J, Ritschel C, Rückriem S, Ruckriem S. Speed-dependent treadmill training in ambulatory hemiparetic stroke patients: a randomized controlled trial. *Stroke*. 2002;33(2):553-558. doi:10.1161/hs0202.102365.
 145. Pohl M, Werner C, Holzgraefe M, et al. Repetitive locomotor training and physiotherapy improve walking and basic activities of daily living after stroke: a single-blind, randomized multicentre trial (DEutsche GAngtrainerStudie, DEGAS). *Clin Rehabil*. 2007;21(1):17-27. doi:10.1177/0269215506071281.
 146. Pomeroy VM, Evans B, Falconer M, Jones D, Hill E, Giakas G. An exploration of the effects of weighted garments on balance and gait of stroke patients with residual disability. *Clin Rehabil*. 2001;15(4):390-397.
 147. Popovic MB, Popovic DB, Sinkjaer T, Stefanovic A, Schwirtlich L. Clinical evaluation of Functional Electrical Therapy in acute hemiplegic subjects. *J Rehabil Res Dev*. 2003;40(5):443-453.
 148. Rabadi M, Galgano M, Lynch D. A pilot study of activity-based therapy in the arm motor recovery post stroke: a randomized controlled trial. *Clin Rehabilitation*. 2008;22:1071-1084. doi:10.1177/0269215508095358.
 149. Ribeiro T, Britto H, Oliveira D, Silva E, Galvao E, Lindquist A. Effects of treadmill training with partial body weight support and the proprioceptive neuromuscular facilitation method on hemiparetic gait: a randomized controlled study. *Eur J Phys Rehabil Med*. 2013;49(4):451-461.
 150. Riccio I, Iolascon G, Barillari MR, Gimigliano R, Gimigliano F. Mental practice is effective in upper limb recovery after stroke: a randomized single-blind cross-over study. *Eur J Phys Rehabil Med*. 2010;46(1):19-25.
 151. Richards CL, Malouin F, Bravo G, Dumas F, Wood-Dauphinee S. The role of technology in task-oriented training in persons with subacute stroke: a randomized controlled trial. *Neurorehabilitation & Neural Repair*. 2004;18(4):199-211. doi:10.1177/1545968304269397.
 152. Richards C, Malouin F, Wood-Dauphinee S, Williams J. JB. Task-specific physical therapy for optimization of gait recovery in acute stroke patients. *Arch Phys Med Rehabil*. 1993;74:612-620.
 153. Rodgers H, Mackintosh J, Price C, et al. Does an early increased-intensity interdisciplinary upper limb therapy programme following acute stroke improve outcome? *Clin Rehabil*. 2003;17(6):579-589. doi:10.1191/0269215503cr652oa.
 154. Rydwick E, Eliasson S, Akner G. The effect of exercise of the affected foot in stroke patients--a randomized controlled pilot trial. *Clin Rehabil*. 2006;20(8):645.

<http://www.ncbi.nlm.nih.gov/pubmed/16944822>.

155. Sackley C, Lincoln N. Single blind randomized controlled trial of visual feedback after stroke: effects on stance symmetry and function. *Disabil Rehabil.* 1997;19:536.
156. Saeys W, Vereeck L, Truijen S, Lafosse C, Wuyts FP, Heyning P Van de. Randomized controlled trial of truncal exercises early after stroke to improve balance and mobility. *Neurorehabil Neural Repair.* 2012;26(3):231-238. doi:10.1177/1545968311416822.
157. Salbach NM, Mayo NE, Hanley JA, et al. A task-orientated intervention enhances walking distance and speed in the first year post stroke: a randomized controlled trial. *Clin Rehabil.* 2004;18(5):509-519. doi:10.1191/0269215504cr763oa.
158. Higgins J, Salbach NM, Wood-Dauphinee S, Richards CL, Cote R, Mayo NE. The effect of a task-oriented intervention on arm function in people with stroke: a randomized controlled trial. *Clin Rehabil.* 2006;20(4):296-310.
159. Saposnik G, Teasell R, Mamdani M, et al. Effectiveness of Virtual Reality Using Wii Gaming Technology in Stroke Rehabilitation. 2010:1477-1484. doi:10.1161/STROKEAHA.110.584979.
160. Schauer M, Mauritz K-HH. Musical motor feedback (MMF) in walking hemiparetic stroke patients: randomized trials of gait improvement. *Clin Rehabil.* 2003;17(7):713-722. doi:10.1191/0269215503cr668oa.
161. Schmid AA, Van Puymbroeck M, Altenburger PA, et al. Poststroke balance improves with yoga: a pilot study. *Stroke.* 2012;43(9):2402-2407.
162. Schwartz I, Sajin A, Fisher I, et al. The effectiveness of locomotor therapy using robotic-assisted gait training in subacute stroke patients: a randomized controlled trial. *Phys Med Rehabil.* 2009;1(6):516-523. doi:10.1016/j.pmrj.2009.03.009.
163. Severinsen K, Jakobsen JK, Pedersen AR, Overgaard K, Andersen H. Effects of resistance training and aerobic training on ambulation in chronic stroke. *Am J Phys Med Rehabil.* 2014;93(1):29-42.
164. Shin WS. The effects of rehabilitation exercise using a home video game (PS2) on gait ability of chronic stroke patients. *J Korea Acad Soc.* 2010;11(1):368-374.
165. Sivenius S, Pyrrala K, Heinonen O, Salonen J, Riekkinen P. The significance of intensity of rehabilitation of stroke—a controlled trial. *Stroke.* 1985;16:928-931.
166. Smania N, Gandolfi M, Paolucci S, et al. Reduced-intensity modified constraint-induced movement therapy versus conventional therapy for upper extremity rehabilitation after stroke: A multicenter trial. *Neurorehabil Neural Repair.* 2012;26(9):1035-1045. doi:10.1177/1545968312446003.
167. Smith DS, Goldenberg E, Ashburn A, et al. Remedial therapy after stroke: a randomised controlled trial. *Br Med J Clin Res Ed.* 1981;282(6263):517-520.
168. Stein J, Krebs HI, Frontera WR, Fasoli S, Hughes R, Hogan N. Comparison of two techniques of robot-aided upper limb exercise training after stroke. *Am J Phys Med Rehabil.* 2004;83(9):720-728.
169. Stinear CM, Barber PA, Coxon JP, Fleming MK, Byblow WD. Priming the motor system enhances the effects of upper limb therapy in chronic stroke. *Brain.* 2008;131:1381-1390.

170. Subramanian SK, Lourenço CB, Chilingaryan G, Sveistrup H, Levin MF. Arm motor recovery using a virtual reality intervention in chronic stroke: randomized control trial. *Neurorehabil Neural Repair*. 2013;27(1):13-23. doi:10.1177/1545968312449695.
171. Sullivan KJ, Knowlton BJ, Dobkin BH. Step training with body weight support: effect of treadmill speed and practice paradigms on poststroke locomotor recovery. *Arch Phys Med Rehabil*. 2002;83(5):683-691. doi:10.1053/apmr.2002.32488.
172. Sullivan KJ, Brown D a, Klassen T, et al. Effects of task-specific locomotor and strength training in adults who were ambulatory after stroke: results of the STEPS randomized clinical trial. *Phys Ther*. 2007;87(12):1580-1607. doi:10.2522/ptj.20060310.
173. Sunderland A, Tinson DJ, Bradley EL, Fletcher D, Langton Hewer R, Wade DT. Enhanced physical therapy improves recovery of arm function after stroke. A randomised controlled trial. *J Neurol Neurosurg Psychiatry*. 1992;55(7):530-535.
174. Sungkarat S, Fisher BE, Kovindha A. Efficacy of an insole shoe wedge and augmented pressure sensor for gait training in individuals with stroke: a randomized controlled trial. *Clin Rehabil*. 2011;25(4):360-369. doi:10.1177/0269215510386125.
175. Sutbeyaz S, Yavuzer G, Sezer N, Koseoglu BF. Mirror therapy enhances lower-extremity motor recovery and motor functioning after stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2007;88(5):555.
176. Tanaka N, Saitou H, Takao T, et al. Effects of gait rehabilitation with a footpad-type locomotion interface in patients with chronic post-stroke hemiparesis: a pilot study. *Clin Rehabil*. 2012;26(8):686-695. doi:10.1177/0269215511432356.
177. Taub E, Uswatte G, King DK, Morris D, Crago JE, Chatterjee A. A placebo-controlled trial of constraint-induced movement therapy for upper extremity after stroke. *Stroke*. 2006;37(4):1045-1049.
178. Thaut M, Leins A, Rice R, Argstatter H, Kenyon G. Rhythmic auditory stimulation improves gait more than NDT/Bobath training in nearambulatory patients early poststroke: a single-blind, randomized trial. *Neurorehabilitation & Neural Repair*. 2007;21:455-459.
179. Timmermans AA, Verbunt JA, van Woerden R, Moennekens M, Pernot DH, Seelen HA. Effect of mental practice on the improvement of function and daily activity performance of the upper extremity in patients with subacute stroke: a randomized clinical trial. *J Am Med Dir Assoc*. 2013;14(3):204-212.
180. Timmermans AA, Lemmens RJ, Monfrance M, et al. Effects of task-oriented robot training on arm function, activity, and quality of life in chronic stroke patients: a randomized controlled trial. *J Neuroengineering Rehabil*. 2014;11:45. doi:http://dx.doi.org/10.1186/1743-0003-11-45.
181. Treger I, Aidinof L. Modified constraint-induced movement therapy improved upper limb function in subacute poststroke patients: a small-scale clinical trial. *Top Stroke Rehabil*. 2012. http://www.maneyonline.com/doi/abs/10.1310/tsr1904-287. Accessed April 15, 2016.
182. van de Port IGL, Wevers LEG, Lindeman E, Kwakkel G. Effects of circuit training as alternative to usual physiotherapy after stroke: randomised controlled trial. *BMJ*. 2012;344(may10 1):e2672. doi:10.1136/bmj.e2672.

183. van Delden AL, Peper CL, Nienhuys KN, Zijp NI, Beek PJ, Kwakkel G. Unilateral versus bilateral upper limb training after stroke: the Upper Limb Training After Stroke clinical trial. *Stroke*. 2013;44(9):2613-2616.
184. van der Lee JH, Wagenaar RC, Lankhorst GJ, Vogelaar TW, Deville WL, Bouter LM. Forced use of the upper extremity in chronic stroke patients: results from a single-blind randomized clinical trial. *Stroke*. 1999;30(11):2369-2375. doi:10.1161/01.STR.30.11.2369.
185. van Nes IJ, Latour H, Schils F, Meijer R, van Kuijk A, Geurts AC. Long-term effects of 6-week whole-body vibration on balance recovery and activities of daily living in the postacute phase of stroke: a randomized, controlled trial. *Stroke*. 2006;37(9):2331-2335.
186. Van Vliet PM, Lincoln NB, Foxall A. Comparison of Bobath based and movement science based treatment for stroke: a randomised controlled trial. *J Neurol Neurosurg Psychiatry*. 2005;76(4):503-508. doi:10.1136/jnnp.2004.040436.
187. Varoqui D, Froger J, Pelissier J, Bardy B. Effect of coordination biofeedback on (re)learning preferred postural patterns in post-stroke patients. *Motor Control*. 2011;15:187.
188. Verheyden G, Vereeck L, Truijen S, et al. Additional exercises improve trunk performance after stroke: a pilot randomized controlled trial. *Neurorehabilitation & Neural Repair*. 2009;23(3):281.
189. Verma R, Arya KN, Garg RK, Singh T. Task-oriented circuit class training program with motor imagery for gait rehabilitation in poststroke patients: a randomized controlled trial. *Top Stroke Rehabil*. 2011;18(Suppl 1):620-632.
190. Visintin M, Barbeau H, Korner-Bitensky N, Mayo NE. A New Approach to Retrain Gait in Stroke Patients Through Body Weight Support and Treadmill Stimulation. *Stroke*. 1998;29(6):1122-1128. doi:10.1161/01.STR.29.6.1122.
191. Volpe BT, Krebs HI, Hogan N, Edelstein OTR L, Diels C, Aisen M. A novel approach to stroke rehabilitation: robot-aided sensorimotor stimulation. 2000;54(10):1938-1944.
192. Wade DT, Collen FM, Robb GF, Warlow CP. Physiotherapy intervention late after stroke and mobility. *BMJ*. 1992;304(6827):609-613.
193. Waldman G, Yang C-YY, Ren Y, et al. Effects of robot-guided passive stretching and active movement training of ankle and mobility impairments in stroke. *NeuroRehabilitation*. 2013;32(3):625-634. doi:10.3233/NRE-130885.
194. Walker M, Gladman J, Lincoln N, Siemonsma P, Whiteley T. Occupational therapy for stroke patients not admitted to hospital: a randomised controlled trial. *Lancet*. 1999;354:278-280.
195. Walker C, Brouwer B, Culham E. Use of visual feedback in retraining balance following acute stroke. *Phys Ther*. 2000;80:886.
196. Wang RY, Chen HI, Chen CY, Yang YR. Efficacy of Bobath versus orthopaedic approach on impairment and function at different motor recovery stages after stroke: a randomized controlled study. *Clin Rehabil*. 2005;19(2):155-164.
197. Werner R, Kessler S. Effectiveness of an intensive outpatient rehabilitation program for postacute stroke patients. *Am J Phys Med Rehabil*. 1996;75:114-120.
198. Werner C, Von Frankenberg S, Treig T, Konrad M, Hesse S. Treadmill training with partial body

- weight support and an electromechanical gait trainer for restoration of gait in subacute stroke patients: a randomized crossover study. *Stroke*. 2002;33(12):2895.
199. Westlake KP, Patten C. Pilot study of Lokomat versus manual-assisted treadmill training for locomotor recovery post-stroke. *J Neuroengineering Rehabil*. 2009;6:18.
 200. Whittall J, Waller SM, Sorkin JD, et al. Bilateral and unilateral arm training improve motor function through differing neuroplastic mechanisms: a single-blinded randomized controlled trial. *Neurorehabilitation & Neural Repair*. 2011;25(2):118-129.
 201. Widen Holmqvist L, von Koch L, Kostulas V, et al. A randomized controlled trial of rehabilitation at home after stroke in southwest Stockholm. *Stroke*. 1998;29(3):591-597. doi:10.1161/01.STR.29.3.591.
 202. Winstein CJ, Rose DK, Tan SM, Lewthwaite R, Chui HC, Azen SP. A randomized controlled comparison of upper-extremity rehabilitation strategies in acute stroke: A pilot study of immediate and long-term outcomes. *Arch Phys Med Rehabil*. 2004;85(4):620-628. doi:10.1016/j.apmr.2003.06.027.
 203. Wittenberg G, Chen R, Ishii K, Bushara K, Eckloff S. Constraint-induced therapy in stroke: magnetic-stimulation motor maps and cerebral activation. *Neurorehabilitation & Neural Repair*. 2003;17:48-57.
 204. Wolf SL, Winstein CJ, Miller JP, et al. Effect of constraint-induced movement therapy on upper extremity function 3 to 9 months after stroke: the EXCITE randomized clinical trial. *Jama*. 2006;296(17):2095-2104.
 205. Wolf SL, Thompson P a, Winstein CJ, et al. The EXCITE stroke trial: comparing early and delayed constraint-induced movement therapy. *Stroke*. 2010;41(10):2309-2315. doi:10.1161/STROKEAHA.110.588723.
 206. Wolf SL, Winstein CJ, Miller JP, et al. Retention of upper limb function in stroke survivors who have received constraint-induced movement therapy: the EXCITE randomised trial. *Lancet Neurol*. 2008;7(1):33-40.
 207. Wu C, Chen C, Tang SF, Lin K, Huang Y. Kinematic and clinical analyses of upper-extremity movements after constraint-induced movement therapy in patients with stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2007;88(8):964-970. doi:10.1016/j.apmr.2007.05.012.
 208. Wu CY, Chen CL, Tsai WC, Lin KC, Chou SH. A randomized controlled trial of modified constraint-induced movement therapy for elderly stroke survivors: changes in motor impairment, daily functioning, and quality of life. *Arch Phys Med Rehabil*. 2007;88(3):273-278.
 209. Wu CY, Chuang LL, Lin KC, Chen HC, Tsay PK. Randomized trial of distributed constraint-induced therapy versus bilateral arm training for the rehabilitation of upper-limb motor control and function after stroke. *Neurorehabilitation & Neural Repair*. 2011;25(2):130-139.
 210. Wu CY, Chen YA, Lin KC, Chao CP, Chen YT. Constraint-induced therapy with trunk restraint for improving functional outcomes and trunk-arm control after stroke: a randomized controlled trial. *Phys Ther*. 2012;92(4):483-492.
 211. Yang Y, Yen J, Wang R-Y, Yen L, Lieu F-K. Gait outcomes after additional backward walking training in patients with stroke: a randomized controlled trial. *Clin Rehabil*. 2005;19(3):264-273.

doi:10.1191/0269215505cr860oa.

212. Yang Y, Wang R, Lin K, Chu M, Chan R. Task-oriented progressive resistance strength training improves muscle strength and functional performance in individuals with stroke. *Clin Rehabil.* 2006;20:860-870.
213. Yang Y, Tsai M-P, Chuang T, Sung W-H, Wang R-Y. Virtual reality-based training improves community ambulation in individuals with stroke: a randomized controlled trial. *Gait Posture.* 2008;28(2):201-206. doi:10.1016/j.gaitpost.2007.11.007.
214. Yang Y, Chen I, Liao K, Huang C, Wang R. Cortical reorganization induced by body weight-supported treadmill training in patients with hemiparesis of different stroke durations. *Arch Phys Med Rehabil.* 2010;91:513.
215. Yang HC, Lee CL, Lin R, et al. Effect of biofeedback cycling training on functional recovery and walking ability of lower extremity in patients with stroke. *Kaohsiung J Med Sci.* 2014;30(1):35-42.
216. Yavuzer G, Eser F, Karakus D, Karaoglan B, Stam HJ. The effects of balance training on gait late after stroke: a randomized controlled trial. *Clin Rehabil.* 2006;20(11):960-969. doi:10.1177/0269215506070315.
217. Yavuzer G, Senel A, Atay MB, Stam HJ. Playstation eyetoy games'' improve upper extremity-related motor functioning in subacute stroke: a randomized controlled clinical trial. *Eur J Phys Rehabil Med.* 2008;44(3):237-244.
218. Yelnik AP, Le Breton F, Colle FM, et al. Rehabilitation of balance after stroke with multisensorial training: a single-blind randomized controlled study. *Neurorehabilitation & Neural Repair.* 2008;22(5):468. doi:10.1177/1545968308315996.
219. Yen J-G, Wang R-Y, Chen H-H, Hong C-T. Effectiveness of modified constraint-induced movement therapy on upper limb function in stroke subjects. *Acta Neurol Taiwan.* 2005;14(1):16-20.
220. Yen C, Wang R, Liao K, Huang C, Yang Y. Gait training induced change in corticomotor excitability in patients with chronic stroke. *Neurorehabilitation & Neural Repair.* 2008;22:22.
221. You SH, Jang SH, Kim Y-H, et al. Virtual reality-induced cortical reorganization and associated locomotor recovery in chronic stroke: an experimenter-blind randomized study. *Stroke.* 2005;36(6):1166-1171. doi:10.1161/01.STR.0000162715.43417.91.
222. Young J, Forster A. The Bradford community stroke trial: eight week results. *Clin Rehabil.* 1991;5:283-292.
223. Young J, Forster A. The Bradford community stroke trial: results at six months. *Br Med J.* 1992;304:1085-1089.