HOOK MONOMIAL IMMANANTS ARE KAZHDAN–LUSZTIG POSITIVE

KEITH MCNULTY

Abstract. Stembridge conjectured that every monomial immanant is non-negative on totally non-negative matrices, and Haiman proposed a stronger statement which at $q = 1$ says that every monomial immanant is a non-negative integer combination of the Kazhdan–Lusztig immanants of Rhoades and Skandera. We prove this for every hook shape $(a, 1^{n-a})$ and every n. More precisely, each of the per-vertex functions obtained by splitting Stembridge's formula for hook characters is Kazhdan–Lusztig positive.

The main tool is a positivity statement for products: a product of Kazhdan–Lusztig positive immanants of complementary submatrices is Kazhdan–Lusztig positive, which we deduce at $q = 1$ from Braden's hyperbolic localization, following Billey and Braden. With Haiman's theorem for irreducible characters, it shows that every class function whose Frobenius characteristic lies in the semiring generated by Schur functions and hook monomial symmetric functions is Kazhdan–Lusztig positive; for example, Skandera's chain of normalised hook immanants is Kazhdan–Lusztig positive for every n.

We place these results in a comparison of three cones of class functions, and show that the remaining obstructions to Haiman's conjecture lie at singular Schubert varieties that are not reached by an iterated doubling identity of Abreu and Nigro, for which we give an explicit criterion. Finally, on the full span of permutation monomials, an element that is non-negative on totally non-negative matrices can have a negative coefficient at the Kazhdan–Lusztig immanant Imm_w if and only if the Schubert variety X_w is singular; Skandera proved the smooth half.

1. Introduction

A real matrix is *totally non-negative* (TNN) if all its minors are non-negative; TNN_n denotes the set of such $n \times n$ matrices. For a function $f\colon S_n \to \mathbb{R}$ and an $n \times n$ matrix x put

$$d_f(x) = \sum_{\sigma \in S_n} f(\sigma)\, x_{1,\sigma(1)} \cdots x_{n,\sigma(n)}.$$

For an irreducible character χ^λ this is the immanant $\mathrm{Imm}_{\chi^\lambda}$, and the permanent and the determinant are the cases $\lambda = (n)$ and $\lambda = (1^n)$. Stembridge [36] proved that every irreducible-character immanant is non-negative on TNN_n. The *monomial immanants* $\mathrm{Imm}_\lambda = d_{\phi^\lambda}$, where ϕ^λ is the monomial virtual character (Section 2), refine them: each χ^μ is a non-negative integer combination of the ϕ^λ.

Conjecture 1.1 (Stembridge [37, Conj. 2.1])**.** *For every $\lambda \vdash n$ and every $A \in \mathrm{TNN}_n$, $\mathrm{Imm}_\lambda(A) \geq 0$.*

The conjecture is known for hooks (Dong–Wu–Gao [13]), for shapes with at most two columns (Clearman–Shelton–Skandera [11]), and for rectangles and $(2, 1^{n-2})$ (Stembridge [37]).

Haiman [16] proposed a stronger statement in the Hecke algebra. Let $P_{v,w}(q)$ be the Kazhdan–Lusztig polynomials of S_n, and let $C'_w(1) = \sum_{v \leq w} P_{v,w}(1)\, v$ be the Kazhdan–Lusztig basis of $\mathbb{Z}[S_n]$

Date: 28 October 2026.

2020 *Mathematics Subject Classification.* Primary 20C08; Secondary 05E05, 05E10, 14M15, 15A15, 15B48.

Key words and phrases. Kazhdan–Lusztig immanant, monomial immanant, hook partition, Haiman conjecture, Hecke algebra character, totally non-negative matrix, Billey–Braden pattern map.

at $q = 1$. Rhoades and Skandera [30] introduced Kazhdan–Lusztig immanants Imm_w, $w \in S_n$, which are non-negative on TNN_n and satisfy $d_f = \sum_w f(C'_w(1))\,\mathrm{Imm}_w$ for every f. We call d_f (or f) *KL-positive* if $f(C'_w(1)) \geq 0$ for every w; then $d_f \geq 0$ on TNN_n.

Conjecture 1.2 (Haiman, Stembridge; $q = 1$)**.** *For every $\lambda \vdash n$ and $w \in S_n$, $\phi^\lambda(C'_w(1)) \geq 0$. Equivalently, every monomial immanant is KL-positive.*

Haiman's conjecture [16, Conj. 2.1] is stronger: for the monomial characters ϕ^λ_q of the Hecke algebra, $\phi^\lambda_q(q^{\ell(w)/2}C'_w)$ has non-negative integer coefficients and is symmetric and unimodal, and Conjecture 1.2 is its value at $q = 1$. Haiman verified it by computer for $n \leq 7$ [16, §4]. Conjecture 1.2 implies Conjecture 1.1. It holds for shapes with at most two columns, by the Temperley–Lieb expansions of Clearman, Shelton and Skandera [11] together with [30, Prop. 3.2]. Our main theorem adds every hook.

Theorem 1.3. *For every n and $1 \leq a \leq n$, the monomial immanant $\mathrm{Imm}_{(a,1^{n-a})}$ is KL-positive. More precisely, for every $i \in [n]$ the per-vertex function*

$$f_i^{(a)}(\sigma) = (-1)^{a-1}\,\mathrm{sgn}(\sigma)\,[\text{the cycle of } \sigma \text{ through } i \text{ has length} \geq a]$$

is KL-positive, and $\phi^{(a,1^{n-a})} = \sum_i f_i^{(a)}$ for $a \geq 2$.

The decomposition into the $f_i^{(a)}$ follows from Stembridge's formula [37, Prop. 1.1], which for hooks counts the vertices in cycles of length $\geq a$ (see [13, Lemma 2.1]); the $f_i^{(a)}$ are the single-permutation versions of the marked cycle sums of [13, Def. 2.2]. Dong, Wu and Gao [13, proof of Thm. 1.2, §2.5] proved by Schur complements that each $d_{f_i^{(a)}}$ is non-negative on every matrix whose minors of order at most $\ell(\lambda) = n-a+1$ are non-negative, answering Stembridge's question [37, Question 2.9] for hooks; in particular $d_{f_i^{(a)}} \geq 0$ on TNN_n. On TNN_n, Theorem 1.3 explains this positivity: each $d_{f_i^{(a)}}$ is a non-negative integer combination of Kazhdan–Lusztig immanants. It does not give their result under the weaker hypothesis, since a Kazhdan–Lusztig immanant need not be non-negative there: for $n = 4$ and $\lambda = (3,1)$, Imm_{1342} and Imm_{1423} occur in the expansion of Imm_λ with coefficient 3, and the matrix A with rows $(3,1,1,0)$, $(3,1,1,1)$, $(2,1,1,1)$, $(2,1,1,2)$ has all 1×1 and 2×2 minors non-negative but $\mathrm{Imm}_{1342}(A) = \mathrm{Imm}_{1423}(A^T) = -1$. For $a = 2$ the expansion is explicit, $\mathrm{Imm}_{(2,1^{n-2})} = 2\sum_k \mathrm{Imm}_{s_k} + \sum_k(\mathrm{Imm}_{s_k s_{k+1}} + \mathrm{Imm}_{s_{k+1}s_k})$ (Corollary 4.2). For $a \geq 3$ the coefficients are not explicit, and the proof is an induction over per-vertex functions composed with rotations of consecutive values (Section 4). Its main tool is the following theorem, which we prove at $q = 1$ from Braden's hyperbolic localization [8], following Billey and Braden [4].

Theorem 1.4. *Let $I, T \subseteq [n]$ with $|I| = |T|$. If F is KL-positive on the submatrix $x[I,T]$ and G is KL-positive on the complementary submatrix $x[I^c,T^c]$, then FG is KL-positive. In particular, the induction product of two KL-positive class functions is KL-positive.*

We prove Theorem 1.4 as Theorem 3.1. By Haiman [16, Lemma 1.1], every irreducible character is KL-positive. Theorems 1.3 and 1.4 therefore give a semiring of KL-positive class functions.

Corollary 1.5. *Let $\mathcal{R}$ be the semiring generated by the Schur functions and the hook monomial symmetric functions $m_{(a,1^b)}$. Every class function whose Frobenius characteristic lies in $\mathcal{R}$ is KL-positive, so its immanant is non-negative on TNN_n.*

For example, Skandera [35, Thm. 27] proved that the normalised hook immanants form a chain per $\geq N_{(n-1,1)} \geq \cdots \geq N_{(1^n)} = \det$ on TNN_n. We show that each difference has Frobenius characteristic $\sum_j h_j m_{(k-j,1^{n-k})}$ up to a positive factor, so the chain holds at the level of Kazhdan–Lusztig immanants (Corollary 5.1).

Three cones. These results are best seen inside a comparison of three cones in the space CF_n of real class functions on S_n:

$$C_{\mathrm{KL}} = \{\phi : \ \phi(C'_w(1)) \geq 0 \text{ for all } w \in S_n\},$$
$$C_{\mathrm{TNN}} = \{\phi : \ d_\phi \geq 0 \text{ on } \mathrm{TNN}_n\},$$
$$C_J = \{\phi : \ d_\phi(J_\mu) \geq 0 \text{ for all } \mu \vdash n\},$$

where J_μ is the block-diagonal matrix with all-ones blocks of sizes $\mu_1, \mu_2, \ldots$. For every n, $C_{\mathrm{KL}} \subseteq C_{\mathrm{TNN}} \subseteq C_J$, and C_J is the simplicial cone spanned by the ϕ^λ (Section 6). So Conjecture 1.1 is the equality $C_{\mathrm{TNN}} = C_J$, which says that the matrices J_μ test total non-negativity of class-function immanants, and Conjecture 1.2 is the equality $C_{\mathrm{KL}} = C_J$. Since the cones are nested, Conjecture 1.2 is equivalent to $C_{\mathrm{KL}} = C_{\mathrm{TNN}} = C_J$: all three cones coincide.

A companion preprint [26] extends Haiman's verification to $n \leq 10$ by an exact computation, which verifies his conjecture in full for these n. In particular Conjecture 1.2 holds for $n \leq 10$, and all three cones coincide. The present paper describes C_{KL} for all n from three directions. Corollary 1.5 describes elements of C_{KL}. Known results show that the coefficients $\phi^\lambda(C'_w(1))$ can only fail to be non-negative at singular Schubert varieties. Clearman, Hyatt, Shelton and Skandera [10], and Abreu and Nigro [1, Thm. 1.6], reduce a smooth (3412- and 4231-avoiding) w to a codominant one. There the coefficients are values of chromatic symmetric functions, which are non-negative by Hikita's proof of the Stanley–Stembridge conjecture [18], as Skandera observed [35, Prop. 25] (Proposition 7.2). The next theorem reduces the problem further.

Theorem 1.6. *If y is joined to a smooth permutation by k* clean steps *(Section 7.2), then $\phi^\lambda(C'_y(1)) = 2^k \phi^\lambda(C'_{w'}(1)) \geq 0$ for a codominant w' and every λ. So Conjecture 1.2 holds for n iff $\phi^\lambda(C'_w(1)) \geq 0$ for every singular w that is not the top of a clean step. If* $\deg P_{e,w} \leq 1$*, a step is clean iff w has no* bad cover *there, an explicit local condition.*

Theorem 1.6 iterates a doubling identity of Abreu and Nigro [1]. Finally, the restriction to class functions matters. For class functions $C_{\mathrm{KL}} = C_{\mathrm{TNN}}$ when $n \leq 10$, and for every n if Conjecture 1.2 holds. On the span Π_n of all permutation monomials, the analogous cones differ already for $n = 4$ [30, §7]: for example, Skandera and Zelevinsky observed that $\mathrm{Imm}_{3214} + \mathrm{Imm}_{1432} - \mathrm{Imm}_{3412}$ is non-negative on TNN_4 but not KL-positive [34, (5.1)], and Theorem 9.2 gives further examples for $n = 5$, such as $\mathrm{Imm}_{21543} + \mathrm{Imm}_{54321} - \mathrm{Imm}_{52431}$. Smoothness appears here too: by a theorem of Skandera [34, Thm. 5.5], a negative coefficient at a smooth w never occurs in a TNN element of Π_n. We prove the converse.

Theorem 1.7. *For every n and every singular $w \in S_n$ there is an element of Π_n that is non-negative on TNN_n and has a negative coefficient at Imm_w. Together with Skandera's theorem, this characterises the smooth w by total non-negativity (Theorem 9.6).*

The proof lifts two elements of Π_4, given by the inequalities $\mathrm{Imm}^2_{3412} \leq \mathrm{Imm}_{1432}\,\mathrm{Imm}_{3214}$ and $\mathrm{Imm}^2_{4231} \leq 3\,\mathrm{Imm}_{2143}\,\mathrm{Imm}_{4321}$ on TNN_4, along an occurrence of 3412 or 4231. Theorem 1.4 controls the lift, and the tangent spaces of Lakshmibai and Seshadri handle the 4231-avoiding case.

The paper is organised linearly. Section 2 recalls Kazhdan–Lusztig immanants and the Temperley–Lieb colouring rule, Section 3 proves Theorem 1.4, and Section 4 proves Theorem 1.3. Section 5 derives Corollary 1.5 and the hook chain, and Section 6 sets up the three cones. Sections 7 and 8 treat smooth and singular permutations and two further families of shapes, Section 9 the full span Π_n and Theorem 1.7, and Section 10 collects conjectures and open problems.

Computational materials. The proof of Theorem 9.2 uses an exact computation in explicit charts. The other computations reported in this paper are cross-checks of the proofs or evidence

for the conjectures: checks of Theorem 1.4 and of the lemmas of Section 4, of Corollary 5.1, of the clean-step criterion of Section 7.2, of Theorem 9.6 and Lemma 9.5, and the computations behind Conjectures 10.1 and 10.2. The code for all of them, together with a script that checks every number in the worked examples and figures, is archived with the code of the companion paper [26] as [27]; the companion paper describes the computational framework and its verification.

2. Preliminaries

2.1. **Class functions.** For a composition α of n let $S_\alpha = S_{\alpha_1} \times S_{\alpha_2} \times \cdots$ be the Young subgroup preserving the consecutive blocks of sizes $\alpha_1, \alpha_2, \ldots$, and let $\alpha! = |S_\alpha|$. For $\mu \vdash n$ put $\eta^\mu = \mathrm{Ind}_{S_\mu}^{S_n} 1$. With $\langle \phi, \psi \rangle = \frac{1}{n!} \sum_\sigma \phi(\sigma)\psi(\sigma)$, the *monomial characters* ϕ^λ are the dual basis: $\langle \phi^\lambda, \eta^\mu \rangle = \delta_{\lambda\mu}$. Under the Frobenius characteristic ch, $\eta^\mu \mapsto h_\mu$, $\chi^\lambda \mapsto s_\lambda$ and $\phi^\lambda \mapsto m_\lambda$. By Young's rule $\langle \chi^a, \eta^\mu \rangle = K_{a,\mu}$, the Kostka number. The induction product $\phi \cdot \psi = \mathrm{Ind}_{S_k \times S_{n-k}}^{S_n}(\phi \times \psi)$ satisfies $\mathrm{ch}(\phi \cdot \psi) = \mathrm{ch}\,\phi\ \mathrm{ch}\,\psi$, and

$$d_{\phi\cdot\psi}(x) = \sum_{|T|=k} d_\phi(x[T,T])\, d_\psi(x[T^c, T^c]). \tag{1}$$

2.2. **Kazhdan–Lusztig immanants.** Rhoades and Skandera [30] define

$$\mathrm{Imm}_w(x) = \sum_{v \geq w} (-1)^{\ell(v)-\ell(w)} P_{w_0 v, w_0 w}(1)\, x_{1,v(1)} \cdots x_{n,v(n)}.$$

They prove that $\mathrm{Imm}_w \geq 0$ on TNN_n and that, for every $f\colon S_n \to \mathbb{R}$, extended linearly to $\mathbb{R}[S_n]$,

$$d_f = \sum_{w \in S_n} f(C'_w(1))\ \mathrm{Imm}_w\,. \tag{2}$$

So the Imm_w are the basis of the span Π_n of permutation monomials dual to $\{C'_w(1)\}$. Comparing coefficients of the monomials shows that (2) is equivalent to the Kazhdan–Lusztig inversion formula $\sum_{u \leq w \leq v} (-1)^{\ell(v)-\ell(w)} P_{u,w}(1) P_{w_0 v, w_0 w}(1) = \delta_{u,v}$ [21, Thm. 3.1]. We write $c_{\lambda,w} = \phi^\lambda(C'_w(1))$, so that $\mathrm{Imm}_\lambda = \sum_w c_{\lambda,w}\, \mathrm{Imm}_w$. For a square submatrix $x[R,C]$ we index its Kazhdan–Lusztig immanants by $S_{|R|}$, through the order-preserving bijections of R and C with $[|R|]$, and call a polynomial *KL-positive on* $x[R,C]$ if it is a non-negative real combination of these. For integer-valued f the coefficients $f(C'_w(1))$ are integers, so for characters, and for the per-vertex functions of Theorem 1.3, KL-positivity means non-negative integer coefficients.

Two special cases are used repeatedly. Since $P_{u,w_0} = 1$ for every u, $\mathrm{Imm}_e = \det$, and (2) with $f = \mathrm{sgn}$ gives $\mathrm{sgn}(C'_w(1)) = \delta_{w,e}$. Since $P_{v,w} \geq 0$ coefficientwise, every monomial $x^v = x_{1,v(1)} \cdots x_{n,v(n)} = d_{\delta_v}$ is KL-positive, with $\delta_v(C'_w(1)) = P_{v,w}(1)$.

2.3. **The Temperley–Lieb colouring rule.** Let τ be a Temperley–Lieb diagram on $n + n$ vertices: left vertices $1, \ldots, n$ and right vertices $1', \ldots, n'$, joined in pairs by non-crossing edges. Edges between two left vertices are *cups*, edges between two right vertices are *caps*, and the others are *through-lines*. Rhoades and Skandera [29] attach to τ a Temperley–Lieb immanant Imm_τ, and [30, Prop. 3.2] shows that $\mathrm{Imm}_\tau = \mathrm{Imm}_w$ for the 321-avoiding w with the same reduced word as τ. This is a bijection between diagrams and 321-avoiding permutations [14]. Their colouring rule [29, Prop. 4.3] expands a product of complementary minors: for $|I| = |I'|$,

$$\det x[I, I']\ \det x[I^c, I'^c] = \sum_\tau \mathrm{Imm}_\tau(x),$$

the sum over the diagrams τ in which every edge joins a black vertex to a white one, where the left vertex i is black iff $i \in I$ and the right vertex j' is black iff $j \notin I'$. We need the case $|I| = 1$ (Figure 1).

Proposition 2.1. *Let $v, c \in [n]$. Then $x_{vc} \det x[\hat{v}, \hat{c}] = \sum_\tau \mathrm{Imm}_\tau$, where τ runs over*

- *the identity diagram, if $c = v$;*
- *the diagrams $\tau_{\epsilon,\delta}$ with one cup $\{v, v+\epsilon\}$, one cap $\{c', (c+\delta)'\}$ and order-preserving through-lines, for $\epsilon, \delta \in \{\pm 1\}$ with $v+\epsilon, c+\delta \in [n]$.*

In particular the non-identity diagrams are determined by their cup and cap, and the diagram with cap $\{k', (k+1)'\}$ occurs for $c = k$ and $c = k+1$ and for no other c.

Proof. Here $I = \{v\}$ and $I' = \{c\}$, so the black vertices are v on the left and every $j' \neq c'$ on the right. Every cup contains the only black left vertex v, so there is at most one cup, and there are as many caps as cups. If there is no cup, τ is the identity diagram, and its edge from v must end at the white vertex c', so $c = v$. If there is one cup, it contains v, and the cap contains c', the only white right vertex. A cup or cap that encloses another vertex would cross the through-line from that vertex, so the cup is $\{v, v \pm 1\}$ and the cap is $\{c', (c \pm 1)'\}$. The remaining $n-2$ left vertices are joined to the remaining $n-2$ right vertices by non-crossing through-lines, which forces the order-preserving matching. Each of these joins a white left vertex to a black right vertex, so all such diagrams are allowed. □




FIGURE 1. Proposition 2.1 for $n = 5$, $v = c = 3$: the five diagrams in $x_{33} \det x[\hat{3}, \hat{3}]$. Black vertices are filled: the left vertex 3 and every right vertex except $3'$. Each edge joins a black vertex to a white one. The identity diagram appears because $c = v$, and the other four have cup $\{3, 3 \pm 1\}$ and cap $\{3', (3 \pm 1)'\}$, with order-preserving through-lines. They are the Temperley–Lieb immanants of s_2, s_3, $s_2 s_3$ and $s_3 s_2$ (Lemma 4.1).

3. Products of immanants of complementary submatrices

Theorem 3.1. *Let $I, T \subseteq [n]$ with $|I| = |T| = k$. If F is KL-positive on $x[I, T]$ and G is KL-positive on $x[I^c, T^c]$, then FG is KL-positive. In particular:*

1. *if G is KL-positive on $x[\hat{\imath}, \hat{\jmath}]$, then $x_{ij} G$ is KL-positive;*
2. *for every bijection γ of a set $T \subseteq [n]$, $x_\gamma \det x[T^c, T^c]$ is KL-positive, where $x_\gamma = \prod_{t \in T} x_{t,\gamma(t)}$;*
3. *if $\phi \in \mathrm{CF}_k$ and $\psi \in \mathrm{CF}_{n-k}$ are KL-positive, then so is $\phi \cdot \psi$.*

Proof. By linearity we may take $F = \mathrm{Imm}_u(x[I, T])$ and $G = \mathrm{Imm}_v(x[I^c, T^c])$ with $u \in S_k$, $v \in S_{n-k}$. For $\sigma \in S_n$ with $\sigma(I) = T$ let $\varphi(\sigma) = (\sigma_I, \sigma_{I^c}) \in S_k \times S_{n-k}$, where σ_I is the restriction $I \to T$ and σ_{I^c} the restriction $I^c \to T^c$, both standardised. Then $FG = d_f$ with $f(\sigma) = [\sigma(I) = T]\, f_u(\sigma_I) f_v(\sigma_{I^c})$, where $\mathrm{Imm}_u = d_{f_u}$ and $f_u(C'_{u'}(1)) = \delta_{uu'}$ on S_k, and similarly for v. By (2) the coefficient of Imm_w in FG is

$$f(C'_w(1)) = \sum_{\sigma(I)=T} P_{\sigma,w}(1)\, f_u(\sigma_I) f_v(\sigma_{I^c}).$$

Let $W' \subseteq S_n$ be the stabiliser of the set of values T. It is generated by the reflections whose roots are orthogonal to the indicator vector $\mathbf{1}_T$, so it is a parabolic reflection subgroup in the sense of Billey and Braden, and $W' \cong S_k \times S_{n-k}$. Its cosets $W'z$ are the sets $\{\sigma : \sigma(I) = T\}$, $I = z^{-1}(T)$, and the pattern map restricts to the bijection φ of $W'z$ onto W'. Braden's theorem on hyperbolic localization [8], applied to the fixed component of a one-parameter subgroup indexed by $W'z$, gives (see the proof of [4, Thm. 4], equation (2))

$$P_{\sigma,w}(1) = \sum_{y \in W'z} a_y \, P_{\varphi(\sigma),\varphi(y)}(1) \qquad (\sigma \in W'z) \tag{3}$$

for integers $a_y = a_y(w) \geq 0$, the multiplicities of the simple summands of the hyperbolic localization of the intersection cohomology sheaf of the Schubert variety X_w. They depend on w and the coset, but not on σ, so the same a_y serve for every $\sigma \in W'z$. The shifts in this decomposition are even, so passing to Euler characteristics introduces no signs. The Kazhdan–Lusztig polynomials of W' are products, so $\sum_{\sigma(I)=T} P_{\sigma,w}(1)\,\varphi(\sigma) = \sum_y a_y \, C'_{\varphi(y)_1}(1) \otimes C'_{\varphi(y)_2}(1)$ in $\mathbb{Z}[S_k] \otimes \mathbb{Z}[S_{n-k}]$. Applying $f_u \otimes f_v$ shows that the coefficient of Imm_w is $a_y \geq 0$ for the y with $\varphi(y) = (u, v)$.

Part (1) is the case $I = \{i\}$, $T = \{j\}$, $F = x_{ij} = \mathrm{Imm}_e(x[\{i\},\{j\}])$. Part (2) follows from (1) by adding one entry at a time, starting from $\det x[T^c, T^c] = \mathrm{Imm}_e$. Part (3) follows from (1): each term $d_\phi(x[T,T])\, d_\psi(x[T^c,T^c])$ is KL-positive by the main statement with $I = T$. □

Example 3.A. Take $n = 3$, $I = \{1\}$, $T = \{2\}$, $F = x_{12}$ and $G = \det x[\hat{1}, \hat{2}]$. The coset $\{\sigma : \sigma(1) = 2\}$ is $\{213, 231\}$, and $\varphi(213) = e$, $\varphi(231) = s_1$ in S_2. Write $a = a_{213}(w)$ and $b = a_{231}(w)$. Since $P_{e,e} = P_{e,s_1} = P_{s_1,s_1} = 1$ and $P_{s_1,e} = 0$, the identity (3) reads $P_{213,w}(1) = a + b$ and $P_{231,w}(1) = b$. Every Kazhdan–Lusztig polynomial of S_3 is 1 on its Bruhat interval, so:

w	123	132	213	231	312	321
$(P_{213,w}, P_{231,w})$	$(0,0)$	$(0,0)$	$(1,0)$	$(1,1)$	$(1,0)$	$(1,1)$
(a,b)	$(0,0)$	$(0,0)$	$(1,0)$	$(0,1)$	$(1,0)$	$(0,1)$

The coefficient of Imm_w in FG is a_y for the y with $\varphi(y) = (e,e)$, that is $a = a_{213}(w)$. So $x_{12} \det x[\hat{1}, \hat{2}] = \mathrm{Imm}_{213} + \mathrm{Imm}_{312}$, which is also what Proposition 2.1 gives with $v = 1$, $c = 2$: the cup $\{1,2\}$ with the caps $\{1', 2'\}$ and $\{2', 3'\}$.

For $k = 1$, the coset is $\{\sigma : \sigma(i) = j\}$ and $\varphi(\sigma)$ is σ with position i and value j deleted, standardised. This is the form used in Section 4. For products of two complementary minors, Proposition 2.1 and its general form [29, Prop. 4.3] give the expansion explicitly. Theorem 3.1 proves at $q = 1$ the positivity of products of Kazhdan–Lusztig immanants of complementary submatrices, which Rhoades and Skandera [31, (3.4)] state, citing Lusztig, as a property of the dual canonical basis.

4. Hooks

4.1. **Per-vertex functions.** For $\sigma \in S_n$ and $i \in [n]$ let $\ell_i(\sigma)$ be the length of the cycle of σ through i, and put

$$f_i^{(a)}(\sigma) = (-1)^{a-1} \operatorname{sgn}(\sigma)\, [\ell_i(\sigma) \geq a].$$

Stembridge's formula [37, Prop. 1.1] (see [13, Lemma 2.1]) gives, with $\kappa_{a,k} = 1$ for $a > 1$ and $\kappa_{1,k} = k + 1$,

$$\kappa_{a,k}\, \phi^{(a,1^k)} = \sum_{i=1}^{n} f_i^{(a)}. \tag{4}$$

Fix i. For $L \geq 1$ let G_L be the sum, over L-subsets $T \ni i$ and L-cycles γ on T, of $x_\gamma \det x[T^c, T^c]$. Sorting permutations by the cycle through i gives

$$\det x = G_1 - G_2 + G_3 - \cdots, \qquad d_{f_i^{(a)}} = \sum_{L \geq a} (-1)^{L-a} G_L. \tag{5}$$

Each G_L is KL-positive by Theorem 3.1(2), but $d_{f_i^{(a)}}$ is an alternating sum of them. For instance,

$$d_{f_i^{(2)}} = x_{ii} \det x[\hat{\imath}, \hat{\imath}] - \det x. \tag{6}$$

4.2. **The case $a = 2$.**

Lemma 4.1. *$f_i^{(2)}(C'_w(1)) = 1$ if $w \in \{s_{i-1}, s_i, s_{i-1}s_i, s_i s_{i-1}\}$ (those that exist), and $f_i^{(2)}(C'_w(1)) = 0$ otherwise.*

Proof. By Proposition 2.1 with $v = c = i$, $x_{ii} \det x[\hat{\imath}, \hat{\imath}]$ is the sum of Imm_τ over the identity diagram and the diagrams with cup $\{i, i \pm 1\}$ and cap $\{i', (i \pm 1)'\}$. These four are the Temperley–Lieb generators t_{i-1}, t_i and the products $t_{i-1}t_i$, $t_i t_{i-1}$, and their 321-avoiding permutations are s_{i-1}, s_i, $s_{i-1}s_i$, $s_i s_{i-1}$. The identity diagram gives $\mathrm{Imm}_e = \det$, which cancels in (6). □

Corollary 4.2. *For every n, $\mathrm{Imm}_{(2,1^{n-2})} = 2\sum_{k=1}^{n-1} \mathrm{Imm}_{s_k} + \sum_{k=1}^{n-2} \big(\mathrm{Imm}_{s_k s_{k+1}} + \mathrm{Imm}_{s_{k+1} s_k}\big)$.*

Proof. Sum Lemma 4.1 over i, using (4): s_k occurs for $i = k$ and $i = k+1$, and $s_k s_{k+1}$, $s_{k+1} s_k$ occur only for $i = k+1$. □

For $n = 3$ this reads $\mathrm{Imm}_{(2,1)} = 2\,\mathrm{Imm}_{213} + 2\,\mathrm{Imm}_{132} + \mathrm{Imm}_{231} + \mathrm{Imm}_{312}$. As a check, Lemma 6.1(3) gives $\mathrm{Imm}_w(J_{(2,1)}) = \delta_{w,213}$, so the right side is 2 at $J_{(2,1)}$, in agreement with $\mathrm{Imm}_{(2,1)}(J_{(2,1)}) = 2!\,1!$ from Lemma 6.1(2).

4.3. **Rotations.** For $a \geq 3$ the alternating sum (5) does not collapse in the same way. Instead we prove a stronger statement, in which the per-vertex function is composed with a rotation, by induction on a. For $b \geq 1$, $m \geq 1$ and $v \in [m]$ write $F_v^{(b)} = f_v^{(b)}$ for the per-vertex function on S_m. Permutations are maps from positions to values, and $(\rho\pi)(x) = \rho(\pi(x))$.

Definition 4.3. A permutation $\rho \in S_m$ is a *good rotation for v* if $\rho = e$, or ρ rotates a block $[l, h]$ of consecutive values with v at its leading end: either $v = h$, $\rho(l) = h$ and $\rho(x) = x - 1$ for $l < x \leq h$ (a *down-rotation*), or $v = l$, $\rho(h) = l$ and $\rho(x) = x + 1$ for $l \leq x < h$ (an *up-rotation*). The other end of the block is the *anchor*; for $\rho = e$ the anchor is v. A block of length 1 gives $\rho = e$, and then either description has anchor v.

Theorem 4.4. *For every $b \geq 1$, m, $v \in [m]$ and good rotation ρ for v, the function $\pi \mapsto F_v^{(b)}(\rho\pi)$ satisfies $F_v^{(b)}(\rho\, C'_u(1)) \geq 0$ for every $u \in S_m$ with $u \neq e$.*

Proof of Theorem 1.3, given Theorem 4.4. Take $\rho = e$: $f_i^{(a)}(C'_u(1)) \geq 0$ for $u \neq e$. At $u = e$, $f_i^{(a)}(C'_e(1)) = f_i^{(a)}(e) = 0$ for $a \geq 2$, while $f_i^{(1)} = \mathrm{sgn}$ is KL-positive. Sum over i with (4). □

The proof of Theorem 4.4 is an induction on b. Fix b, m, v, ρ and put $h(\pi) = F_v^{(b)}(\rho\pi)$. For $c \in [m]$, the permutations with $\pi(v) = c$ form a coset of the stabiliser of the value c. As in Section 3, let $\pi' \in S_{m-1}$ be π with position v and value c deleted, standardised, and define $H_c(\pi') = h(\pi)$.

Lemma 4.5 (closure)**.** *Let $t = \rho(c)$. If $t = v$, that is, if c is the anchor, then $H_c = 0$ (for $b \geq 2$). Otherwise $H_c(\pi') = F_{v'}^{(b-1)}(\rho'\pi')$, where v' is the position of t in $[m] \setminus \{v\}$ and ρ' is a good rotation for v' whose block has length $|c - \text{anchor}|$.*

Proof. Put $\sigma = \rho\pi$, so $\sigma(v) = t$. If $t = v$, the cycle of σ through v has length $1 < b$. Otherwise remove v from its cycle: let τ be the permutation of $[m]\setminus\{v\}$ with $\tau(x) = \sigma(x)$, except $\tau(\sigma^{-1}(v)) = t$. The cycle through t becomes one shorter, the other cycles are unchanged, and the number of cycles is the same, so $\operatorname{sgn}\sigma = -\operatorname{sgn}\tau$ and $F_v^{(b)}(\sigma) = F_t^{(b-1)}(\tau)$. For $x \neq v$, $\tau(x) = \psi(\rho(\pi(x)))$ with $\psi(v) = t$ and ψ the identity elsewhere. Here $\psi\colon [m]\setminus\{t\} \to [m]\setminus\{v\}$ is a bijection because $t \neq v$. Let $\operatorname{std}_v\colon [m]\setminus\{v\} \to [m-1]$ and $\operatorname{std}_c\colon [m]\setminus\{c\} \to [m-1]$ be the order-preserving bijections, so that $\pi' = \operatorname{std}_c \circ \pi \circ \operatorname{std}_v^{-1}$. Conjugating τ by std_v gives $\operatorname{std}_v \circ \tau \circ \operatorname{std}_v^{-1} = \rho'\pi'$ with $\rho' = \operatorname{std}_v \circ \psi \circ \rho \circ \operatorname{std}_c^{-1} \in S_{m-1}$, and conjugation preserves cycle lengths and signs, so $F_t^{(b-1)}(\tau) = F_{v'}^{(b-1)}(\rho'\pi')$ with $v' = \operatorname{std}_v(t)$. By the symmetry $w \mapsto w_0 w w_0$, which preserves Kazhdan–Lusztig polynomials, we may assume that $\rho = e$ or that ρ is the down-rotation of $[l, v]$. A direct check gives three cases: if $c > v$, ρ' is the down-rotation of $[l, c-1]$ and $v' = c-1$; if $l < c \le v$, ρ' is the down-rotation of $[l, c-1]$ and $v' = c-1$; if $c < l$, ρ' is the up-rotation of $[c, l-1]$ and $v' = c$. In each case ρ' is good for v' and its block has length $|c-l|$. □

Example 4.A. Let $m = 6$ and let $\rho = 125346$ be the down-rotation of $[3, 5]$, so $v = 5$ and the anchor is $l = 3$. The three cases of the proof, and the anchor, give:

c	$t = \rho(c)$	case	$\rho' \in S_5$	v'	$\lvert c-l\rvert$
1	1	$c < l$	21345, up-rotation of $[1, 2]$	1	2
2	2	$c < l$	e, up-rotation of $[2, 2]$	2	1
3	5	anchor	$H_3 = 0$		
4	3	$l < c \le v$	e, down-rotation of $[3, 3]$	3	1
5	4	$l < c \le v$	12435, down-rotation of $[3, 4]$	4	2
6	6	$c > v$	12534, down-rotation of $[3, 5]$	5	3

For instance, at $c = 6$ the value $t = 6$ is fixed by ρ, and removing $v = 5$ from the cycle of $\sigma = \rho\pi$ through it gives a permutation τ of $\{1, 2, 3, 4, 6\}$; after standardising, $\tau = \rho'\pi'$ with ρ' again the down-rotation of $[3, 5]$, now in S_5, and $v' = 5$ is the position of $t = 6$. In each row $H_c = F_{v'}^{(b-1)} \circ \rho'$ has been checked on all of S_5 for $b = 2, 3, 4, 5$.

Lemma 4.6 (bottom terms)**.** *For $u \neq e$, $\sum_c H_c(e)\, D_{vc}(u) \ge 0$, where $D_{vc}(u)$ is the coefficient of* Imm_u *in $x_{vc} \det x[\hat{v}, \hat{c}]$.*

Proof. Let l be the anchor and let $c \in [m]$. If $c = l$, then $H_c = 0$ by Lemma 4.5. Otherwise ρ' rotates a block of length $L = |c-l| \ge 1$, so it is a single L-cycle through v' (a fixed point if $L = 1$) and $\operatorname{sgn}\rho' = (-1)^{L-1}$. Hence, in both cases,

$$H_c(e) = F_{v'}^{(b-1)}(\rho') = (-1)^{b+|c-l|+1}\,[\,|c-l| \ge b-1\,].$$


So $H_c(e) = 0$ exactly for c in $Z \cap [m]$, where $Z = [l-b+2,\, l+b-2]$ is an interval containing l (as $b \ge 2$), and

$$H_c(e) = \kappa\,(-1)^{v+c} \quad (c \in [m]\setminus Z), \qquad \kappa = (-1)^{b+1+v+l}.$$

By Proposition 2.1, the coefficient of the diagram with cup $\{v, v \pm 1\}$ and cap $\{k', (k+1)'\}$ in $\sum_c H_c(e)\, x_{vc} \det x[\hat{v}, \hat{c}]$ is $H_k(e) + H_{k+1}(e)$ (Figure 2). If k and $k+1$ both lie outside Z, this is $\kappa(-1)^{v+k} + \kappa(-1)^{v+k+1} = 0$, and if both lie in Z it is 0. Since Z is an interval, the remaining pairs straddle its boundary. For $k = l-b+1$, with $k+1 \in Z$, the coefficient is $H_k(e) = \kappa(-1)^{v+l-b+1} = +1$. For $k+1 = l+b-1$, with $k \in Z$, it is $H_{k+1}(e) = \kappa(-1)^{v+l+b-1} = +1$. A boundary pair that does not lie inside $[m]$ gives no diagram. So

$$\sum_c H_c(e)\, x_{vc} \det x[\hat{v}, \hat{c}] \;=\; H_v(e)\, \operatorname{Imm}_e \;+ \sum_{\substack{k \in \{l-b+1,\, l+b-2\} \\ 1 \le k < m}} \sum_{\pm} \operatorname{Imm}_{\tau_{\pm,k}},$$

where $\tau_{\pm,k}$ has cup $\{v, v \pm 1\}$ and cap $\{k', (k+1)'\}$, and the inner sum is over the cups that exist. The two boundary values $k = l - b + 1$ and $k = l + b - 2$ differ, since $2b \neq 3$, so the diagrams in the double sum have distinct caps or distinct cups and are pairwise distinct by Proposition 2.1. Distinct diagrams correspond to distinct 321-avoiding permutations [14], hence to distinct Kazhdan–Lusztig immanants, so for $u \neq e$ the sum $\sum_c H_c(e) D_{vc}(u)$ is 0 or 1. □



$x_{2c} \det x[\hat{2}, \hat{c}]$, $c = k = 4$ $\qquad$ $x_{2c} \det x[\hat{2}, \hat{c}]$, $c = k + 1 = 5$

$Z = [1, 3]$

c	1	2	3	4	5	6
$H_c(e)$	0	0	0	+1	−1	+1
$H_k(e) + H_{k+1}(e)$		0	0	+1	0	0

FIGURE 2. Lemma 4.6 for $m = 6$, $v = 2$, $k = 4$. The diagram with cup $\{2, 3\}$ and cap $\{4', 5'\}$ is compatible with the colouring of $x_{24} \det x[\hat{2}, \hat{4}]$ (left) and with that of $x_{25} \det x[\hat{2}, \hat{5}]$ (right), and with no other summand; black vertices are filled. Its two coefficients $H_4(e)$ and $H_5(e)$ have opposite signs unless one of $4, 5$ lies in Z. Below, the values $H_c(e)$ for $\rho = e$ and $b = 3$, where $l = v = 2$ and $Z = [1, 3]$. The last row, placed between k and $k+1$, is the coefficient of the diagram with cap $\{k', (k+1)'\}$: the pairs $(4, 5)$ and $(5, 6)$ cancel, the only surviving pair is the boundary pair $(3, 4)$, with coefficient $H_4(e) = +1$, and the other boundary $k = l - b + 1 = 0$ lies outside $[m]$.

Proof of Theorem 4.4. Induction on b. For $b = 1$, $F_v^{(1)}(\rho\pi) = \operatorname{sgn}(\rho) \operatorname{sgn}(\pi)$, and $\operatorname{sgn}(C'_u(1)) = \delta_{u,e}$. Let $b \geq 2$. Apply (3) with $k = 1$ to the coset $\{\pi(v) = c\}$: for each c there are integers $a_y^{(c)}(u) \geq 0$, indexed by the coset, with

$$h(C'_u(1)) = \sum_c \sum_y a_y^{(c)}(u)\, H_c\big(C'_{\varphi(y)}(1)\big).$$

The terms with $\varphi(y) \neq e$ are ≥ 0 by Lemma 4.5 and induction. Each coset contains exactly one y with $\varphi(y) = e$, since φ is a bijection of the coset onto S_{m-1}, and its term is $a_y^{(c)}(u) H_c(e)$. The same multiplicity occurs in Theorem 3.1 with $I = \{v\}$, $T = \{c\}$, $F = x_{vc}$ and $G = \det x[\hat{v}, \hat{c}] = \operatorname{Imm}_e(x[\hat{v}, \hat{c}])$: there $x_{vc} \det x[\hat{v}, \hat{c}] = d_g$ with $g(\sigma) = [\sigma(v) = c] \operatorname{sgn}(\varphi(\sigma))$, and since $\operatorname{sgn}(C'_{u'}(1)) = \delta_{u',e}$, the proof of Theorem 3.1 shows that the coefficient of Imm_u is $a_y^{(c)}(u)$ for this y. So $a_y^{(c)}(u) = D_{vc}(u)$, and these terms sum to $\sum_c H_c(e) D_{vc}(u) \geq 0$ by Lemma 4.6. □

For $\rho = e$ the bottom terms are $(-1)^{a-1}$ times a tail $\sum_{|j-i| \geq a-1} (-1)^{i+j} x_{ij} \det x[\hat{\imath}, \hat{\jmath}]$ of the Laplace expansion of $\det x$ along row i, and Lemma 4.6 identifies them with at most four staircase Kazhdan–Lusztig immanants. For $a = 2$ this is Lemma 4.1.

Example 4.B. Let $n = 4$, $a = 3$, $i = 1$. The non-zero Kazhdan–Lusztig coefficients of $f_1^{(3)}$ are 1 at 2314, 3124, 3241, 4213 and 2 at 3214, 4231. In the proof, $\rho = e$, $v = l = 1$ and $b = 3$, so $H_c(e) = 0$ for $c \in \{1, 2\}$, $H_3(e) = +1$ and $H_4(e) = -1$. Only the boundary cap $\{2', 3'\}$ survives, with the cup $\{1, 2\}$, and this diagram is the staircase 3124. So the bottom terms contribute exactly Imm_{3124}. The rest of $f_1^{(3)}$, namely 1 at 2314, 3241, 4213 and 2 at 3214, 4231, comes from the pieces $H_c = F^{(2)}_{v'} \circ \rho'$, which are non-negative off e by the case $b = 2$. In detail, $H_1 = 0$, and for $c = 2, 3, 4$ the closure lemma gives $(\rho', v') = (e, 1)$, $(213, 2)$ and $(312, 3)$. Writing $D^{(c)}_{u'}(u)$ for the coefficient of Imm_u in $x_{1c}\,\mathrm{Imm}_{u'}(x[\hat{1}, \hat{c}])$, the proof of Theorem 4.4 splits $h(C'_u(1)) = \sum_c \sum_{u'} D^{(c)}_{u'}(u)\, H_c(C'_{u'}(1))$ as follows:

c	u' with $H_c(C'_{u'}(1)) \neq 0$	$x_{1c}\,\mathrm{Imm}_{u'}(x[\hat{1}, \hat{c}])$
2	213: 1	$\mathrm{Imm}_{2314} + \mathrm{Imm}_{3214}$
3	e: 1	$\mathrm{Imm}_{3124} + \mathrm{Imm}_{4123}$
	213: 1	$\mathrm{Imm}_{3214} + \mathrm{Imm}_{4213}$
	231: 1	$\mathrm{Imm}_{3241} + \mathrm{Imm}_{4231}$
4	e: -1	Imm_{4123}
	231: 1	Imm_{4231}

So the contributions of $c = 2, 3, 4$ are

$$\mathrm{Imm}_{2314} + \mathrm{Imm}_{3214}, \qquad \mathrm{Imm}_{3124} + \mathrm{Imm}_{4123} + \mathrm{Imm}_{3214} + \mathrm{Imm}_{4213} + \mathrm{Imm}_{3241} + \mathrm{Imm}_{4231},$$
$$- \mathrm{Imm}_{4123} + \mathrm{Imm}_{4231}\,.$$

The two bottom terms at Imm_{4123} cancel, as in Lemma 4.6, and the three contributions add up to the expansion of $f_1^{(3)}$ above.

The induction works at $q = 1$ only, because (3) is a statement about Euler characteristics. We do not know whether $\phi_q^{(a,1^{n-a})}(q^{\ell(w)/2} C'_w(q)) \in \mathbb{N}[q]$ for every n (Problem 3); for $n \leq 10$ it holds by [26].

5. Consequences

Proof of Corollary 1.5. The KL-positive class functions are closed under sums, and under induction products by Theorem 3.1(3). Every irreducible character is KL-positive by Haiman [16, Lemma 1.1], and every hook monomial character by Theorem 1.3. Since ch is additive and multiplicative, every class function with characteristic in $\mathcal{R}$ is KL-positive. The immanant is then non-negative on TNN_n by (2) and the positivity of the Imm_w [30]. □

Corollary 1.5 contains the $h_\mu = s_{(\mu_1)} s_{(\mu_2)} \cdots$, so every η^μ is KL-positive, as is directly clear from $d_{\eta^\mu}(x) = \sum \prod_t \mathrm{per}\, x[T_t, T_t]$. A product such as $h_j m_{(a,1^b)}$ involves monomial symmetric functions m_λ with λ not a hook, so Corollary 1.5 reaches class functions that are not combinations of hook monomial characters, and it does not follow from Haiman's theorem alone.

Write $f^a = \chi^a(e)$ and $N_a = \mathrm{Imm}_{\chi^a}/f^a$ for the normalised immanants.

Corollary 5.1. *For every n and $2 \leq k \leq n$ put $D_k = (k-1)\chi^{(k,1^{n-k})} - (n-k+1)\chi^{(k-1,1^{n-k+1})}$, so that $N_{(k,1^{n-k})} - N_{(k-1,1^{n-k+1})} = \frac{(k-2)!\,(n-k)!}{(n-1)!}\, d_{D_k}$. Then*

$$\mathrm{ch}\, D_k = \sum_{j=0}^{k-2} h_j\, m_{(k-j,1^{n-k})}, \qquad d_{D_k}(x) = \sum_{j=0}^{k-2} \sum_{|T|=j} \mathrm{per}\, x[T, T]\ \mathrm{Imm}_{(k-j,1^{n-k})}\big(x[T^c, T^c]\big),$$

and D_k is KL-positive. In particular per $\geq N_{(n-1,1)} \geq \cdots \geq$ det *on* TNN_n *for every n, with every difference a non-negative combination of Kazhdan–Lusztig immanants.*

Proof. Let $H(x) = \sum_j h_j x^j$ and $E(y) = \sum_j e_j y^j$ in variables $x_1, x_2, \dots$, and $F = \sum_{a,b\geq 0} s_{(a+1,1^b)} x^a y^b$. The hook Schur functions satisfy $(x+y)F = H(x)E(y) - 1$. The coefficient of $x^{k-1}y^{n-k}$ in $x(\partial_x - \partial_y)F$ is $\operatorname{ch} D_k$, and since $(\partial_x - \partial_y)(x+y) = 0$,

$$x(\partial_x - \partial_y)F = \frac{x\,(H'E - HE')}{x+y} = xHE \sum_i \frac{x_i^2}{(1-xx_i)(1+yx_i)},$$

using $H'/H = \sum_i x_i/(1-xx_i)$ and $E'/E = \sum_i x_i/(1+yx_i)$. On the other hand $\sum_b m_{(c,1^b)} y^b = \sum_i x_i^c \prod_{j\neq i}(1+yx_j)$ for $c \geq 2$, so $H(x)\sum_{c\geq 2,\, b\geq 0} m_{(c,1^b)} x^{c-1} y^b$ is the same series. Comparing coefficients of $x^{k-1}y^{n-k}$ gives the first identity. The second follows from (1). KL-positivity is Corollary 1.5. □

Skandera's proofs [35] give total non-negativity: one counts tableaux on planar networks, and the other expands D_k in the immanants of $\theta_\ell = \sum_{\ell(\mu)=\ell} \phi^\mu$, whose KL-positivity is open (Problem 6). Corollary 5.1 avoids the θ_ℓ. For total non-negativity alone, the theorem of Dong–Wu–Gao [13] can replace Theorem 1.3 in the proof.

6. Three cones

In this section we set up the cones $C_{\mathrm{KL}} \subseteq C_{\mathrm{TNN}} \subseteq C_J$ of the introduction (Figure 3), and show how they organise the known results. Let $w_0(\alpha)$ be the longest element of S_α.

Lemma 6.1. (1) *For every composition α of n, $J_\alpha \in \mathrm{TNN}_n$.*
(2) *For every class function ϕ, $d_\phi(J_\mu) = \mu!\,\langle \phi, \eta^\mu \rangle$. In particular $\mathrm{Imm}_\lambda(J_\mu) = \mu!\,\delta_{\lambda\mu}$.*
(3) *For every $w \in S_n$, $\mathrm{Imm}_w(J_\alpha) = \delta_{w,w_0(\alpha)}$. Hence $d_\phi(J_\mu) = \phi(C'_{w_0(\mu)}(1))$.*

Proof. (1) Order the rows and columns of a square submatrix by the blocks they lie in. If some block contributes more rows than columns, those rows are supported on too few columns and the minor vanishes. Otherwise every block contributes equally many rows and columns, since the totals agree, so the submatrix is block diagonal with square all-ones blocks, and the minor is a product of determinants of all-ones matrices, each equal to 0 or 1. (2) $\prod_i (J_\mu)_{i,\sigma(i)} = [\sigma \in S_\mu]$, so $d_\phi(J_\mu) = \sum_{\sigma\in S_\mu} \phi(\sigma) = \mu!\,\langle \phi, \eta^\mu \rangle$ by Frobenius reciprocity. (3) The lower interval of $w_0(\alpha)$ is S_α, and $P_{v,w_0(\alpha)} = 1$ for $v \in S_\alpha$, so $C'_{w_0(\alpha)}(1) = \sum_{v\in S_\alpha} v$. For every f, (2) at J_α gives $\sum_w f(C'_w(1))\,\mathrm{Imm}_w(J_\alpha) = \sum_{v\in S_\alpha} f(v) = f(C'_{w_0(\alpha)}(1))$. Since $\{C'_w(1)\}$ is a basis and f is arbitrary, $\mathrm{Imm}_w(J_\alpha) = \delta_{w,w_0(\alpha)}$. □

Theorem 6.2. *For every n, $C_{\mathrm{KL}} \subseteq C_{\mathrm{TNN}} \subseteq C_J$, and $C_J = \mathrm{cone}\{\phi^\lambda : \lambda \vdash n\}$ is simplicial. Consequently:*

(1) *Conjecture 1.1 holds for n iff $C_{\mathrm{TNN}} = C_J$, iff every class function ϕ with $d_\phi(J_\mu) \geq 0$ for all μ satisfies $d_\phi \geq 0$ on TNN_n;*
(2) *Conjecture 1.2 holds for n iff $C_{\mathrm{KL}} = C_J$, iff KL-positivity of a class function is detected by its $p(n)$ coefficients at the elements $w_0(\mu)$;*
(3) *Conjecture 1.2 holds for n iff $C_{\mathrm{KL}} = C_{\mathrm{TNN}} = C_J$; in particular it implies Conjecture 1.1 and $C_{\mathrm{KL}} = C_{\mathrm{TNN}}$.*

Proof. The first inclusion is (2) together with $\mathrm{Imm}_w \geq 0$ on TNN_n, and the second is Lemma 6.1(1). By Lemma 6.1(2), $\phi \in C_J$ iff $\langle \phi, \eta^\mu \rangle \geq 0$ for all μ, iff ϕ has non-negative coordinates in the basis $\{\phi^\lambda\}$ dual to $\{\eta^\mu\}$. So $C_{\mathrm{TNN}} = C_J$ iff every generator ϕ^λ lies in C_{TNN}, which is Conjecture 1.1, and similarly for C_{KL}. The reformulation in (2) is Lemma 6.1(3). Part (3) follows from the chain of inclusions. □




FIGURE 3. A schematic slice of the three cones of Theorem 6.2. The outer cone C_J is simplicial, with extreme rays ϕ^λ, $\lambda \vdash n$. Conjecture 1.1 says that the middle cone fills it, and Conjecture 1.2 that the inner one does. For $n \leq 10$ all three coincide [26], so the picture shows only the logical possibilities, not an actual n.

6.1. **What is known.** In this language, the known general results read as follows.

- Rhoades–Skandera [30]: $\mathrm{Imm}_w \geq 0$ on TNN_n, which gives $C_{\mathrm{KL}} \subseteq C_{\mathrm{TNN}}$.
- Haiman [16, Lemma 1.1]: every irreducible character lies in C_{KL}. This refines Stembridge's theorem [36] that $\chi^a \in C_{\mathrm{TNN}}$.
- Clearman–Shelton–Skandera [11]: $\phi^\lambda \in C_{\mathrm{KL}}$ when $\lambda_1 \leq 2$. Stembridge [37, Thms. 2.7, 2.8]: $\phi^\lambda \in C_{\mathrm{TNN}}$ for rectangles and $(2, 1^{n-2})$. Dong–Wu–Gao [13]: $\phi^\lambda \in C_{\mathrm{TNN}}$ for hooks (indeed under the weaker hypothesis of Stembridge's [37, Question 2.9]).
- This paper: $\phi^\lambda \in C_{\mathrm{KL}}$ for hooks (Theorem 1.3); C_{KL} is closed under induction products (Theorem 3.1), and so are C_{TNN}, by (1), and C_J, since products of monomial symmetric functions are monomial-positive. This is consistent with the three cones being equal.
- The companion paper [26]: Haiman's conjecture, at generic q, for $n \leq 10$; hence $C_{\mathrm{KL}} = C_{\mathrm{TNN}} = C_J$ for $n \leq 10$.

Since each ϕ^λ spans an extreme ray of C_J, induction products never produce a new λ with $\phi^\lambda \in C_{\mathrm{KL}}$. They do produce new elements of C_{KL}, as in Corollary 5.1.

6.2. **Inequalities between normalised immanants.** For $a, b \vdash n$ write $a \succeq b$ if $N_a \geq N_b$ on TNN_n. The cones turn this order into a question about Kostka numbers.

Proposition 6.3. *Let $a, b \vdash n$. If $a \succeq b$ then $K_{a,\mu}/f^a \geq K_{b,\mu}/f^b$ for all $\mu \vdash n$, and if this criterion fails at μ then $N_a(J_\mu) < N_b(J_\mu)$. Conversely, if Conjecture 1.1 holds for n and the criterion holds, then $a \succeq b$. In particular, $a \succeq b$ implies that a dominates b.*

Proof. $f^a f^b (N_a - N_b) = d_\phi$ with $\phi = f^b \chi^a - f^a \chi^b$, and $d_\phi(J_\mu) = \mu!\,(f^b K_{a,\mu} - f^a K_{b,\mu})$ by Lemma 6.1. The first statement holds because $J_\mu \in \mathrm{TNN}_n$, and the converse is Theorem 6.2(1). At $\mu = b$ the criterion gives $K_{a,b} \geq f^a K_{b,b}/f^b > 0$, so a dominates b. □

The first statement is Stembridge's [37, Prop. 3.1], and he notes that Conjecture 1.1 gives the converse [37, §3].

Proposition 6.4. *For every n and $a \vdash n$, $\mathrm{per} \geq N_a \geq \det$ on TNN_n, and both differences are KL-positive.*

Proof. The coefficient of x^v in $\mathrm{per} - N_a$ is $1 - \chi^a(v)/f^a \geq 0$, and every monomial is KL-positive. For the bottom, $N_a - \det = \sum_{w \neq e} \frac{\chi^a(C'_w(1))}{f^a} \mathrm{Imm}_w$, because $\mathrm{Imm}_e = \det$ and the coefficient of Imm_e is $\chi^a(e)/f^a - 1 = 0$, and $\chi^a(C'_w(1)) \geq 0$ by Haiman [16, Lemma 1.1]. □

Proposition 6.5. *If $d_\phi \geq 0$ on all Hermitian positive semidefinite $n \times n$ matrices, then $\phi \in C_J$. Hence $d_\phi \geq 0$ on TNN_n if Conjecture 1.1 holds for n, and d_ϕ is KL-positive if Conjecture 1.2 holds for n.*

Proof. Each J_μ is a direct sum of matrices $\mathbf{1}\mathbf{1}^T$, so it is positive semidefinite. Apply Theorem 6.2. □

For normalised characters, $\phi = f^b\chi^a - f^a\chi^b$, this is Stembridge's observation that, since each J_μ is positive semidefinite, an inequality $N_a \geq N_b$ on positive semidefinite matrices implies the criterion of Proposition 6.3 [37, §3]. So every linear inequality between immanants known on positive semidefinite matrices transfers to TNN_n under Conjecture 1.1, and to the Kazhdan–Lusztig level under Conjecture 1.2. Examples are Schur's inequality $N_a \geq \det$ [32], Heyfron's hook chain [17], and Pate's inequalities [28], which compare N_λ with N_μ when μ is obtained from λ by removing its last column and adding its boxes to the bottom of the first column. Skandera asked whether Pate's inequalities hold on TNN_n [35, Problem 29]; by Proposition 6.5 they follow from Conjecture 1.1, and hence hold for $n \leq 10$ [26]. For Schur's inequality and Heyfron's chain, Proposition 6.4 and Corollary 5.1 give the Kazhdan–Lusztig statement for every n unconditionally.

7. Smooth and singular permutations

7.1. **Smooth permutations.** A permutation is *codominant* if it avoids 312. For codominant w put $m(i) = \max(w(1), \ldots, w(i))$, a Hessenberg function, and let A_w be the 0/1 matrix with $(A_w)_{ij} = [j \leq m(i)]$.

Proposition 7.1. *For codominant w, $A_w \in \mathrm{TNN}_n$ and $\sum_v \prod_i (A_w)_{i,v(i)}\, v = C'_w(1)$. Consequently $f(C'_w(1)) = d_f(A_w)$ for every $f\colon S_n \to \mathbb{R}$, and $\mathrm{Imm}_v(A_w) = \delta_{v,w}$ for every v.*

Proof. In a square submatrix of A_w, row s is the indicator of an initial segment $[1, \mu_s]$ with μ non-decreasing. Subtracting consecutive rows leaves rows supported on disjoint intervals, so every minor is 0 or 1. Next, $\prod_i (A_w)_{i,v(i)} = [v(i) \leq m(i)\ \forall i]$, which equals $[v \leq w]$ for codominant w: a 312-avoiding w avoids the four patterns of [33, Thm. 3.1], so $[e, w]$ is the set of rook placements on its right hull, which is the Ferrers board $\{(i,j) : j \leq m(i)\}$ [33, Rem. 3.4]. Codominant w are smooth, so $P_{v,w} = 1$ for all $v \leq w$, and the sum is $C'_w(1)$. Comparing $d_f(A_w) = f(C'_w(1))$ with (2) for all f gives $\mathrm{Imm}_v(A_w) = \delta_{v,w}$. □

Proposition 7.1 is essentially known: see the proof of [35, Prop. 25] and the references there. For $w = w_0(\alpha)$ the matrix A_w is not J_α, since it has ones below the diagonal blocks, but both have $\prod_i M_{i,v(i)} = [v \in S_\alpha]$, so Proposition 7.1 extends Lemma 6.1(3).

Example 7.A. For the codominant $w = 2341$, $m = (2,3,4,4)$, and for $w = w_0(2,2) = 2143$, $m = (2,2,4,4)$:

$$A_{2341} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \qquad A_{2143} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \qquad J_{(2,2)} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Each row of A_w is the indicator of an initial segment, and the segments grow down the rows. One checks directly that $\mathrm{Imm}_v(A_{2341}) = \delta_{v,2341}$ and $\mathrm{Imm}_v(A_{2143}) = \mathrm{Imm}_v(J_{(2,2)}) = \delta_{v,2143}$ for all $v \in S_4$.

Proposition 7.2 (Clearman–Hyatt–Shelton–Skandera, Abreu–Nigro, Hikita, Skandera)**.** *For every smooth $w \in S_n$ and every $\lambda \vdash n$, $\phi^\lambda(C'_w(1)) \geq 0$.*

This is a direct combination of known results, which we record for later use. For codominant w it is [35, Prop. 25], where Skandera deduces it from Hikita's theorem.

Proof. For smooth w, $C'_w(1)$ and $C'_{w'}(1)$ have the same character values for some codominant w' (Abreu–Nigro [1, Thm. 1.6]; see also Clearman–Hyatt–Shelton–Skandera [10, Thm. 4.6]). By [10, Cor. 7.2] there is a unit interval order P with $X_P = \sum_\lambda \phi^\lambda(C'_w(1))\, e_\lambda$, where X_P is Stanley's chromatic symmetric function of the incomparability graph of P. These coefficients are non-negative by Hikita's proof of the Stanley–Stembridge conjecture [18]. □

Proposition 7.2, and through it Theorem 7.4, the reduction for the shapes $(3, 2^b, 1^c)$ in Section 8 and the smooth case of Problem 6, depend on Hikita's theorem [18]. Theorem 1.3 does not use it. For hooks, Hikita's theorem is not needed: $f_i^{(a)}(C'_w(1)) \geq 0$ for every w, smooth or not, by Theorem 1.3. More generally, Skandera [34, Thm. 5.5] showed that every immanant d_f that is non-negative on TNN_n has non-negative coefficients at all smooth w. So Conjecture 1.1 for a shape λ implies $c_{\lambda,w} \geq 0$ at every smooth w; Proposition 7.2 proves this unconditionally.

7.2. **Singular permutations: clean steps.** Here C'_w is the Kazhdan–Lusztig basis of the Hecke algebra $H_n(q)$, and $\mathrm{ch}(h) = \sum_\lambda \chi_q^\lambda(h)\, s_\lambda$ for $h \in H_n(q)$, where χ_q^λ are the irreducible characters of $H_n(q)$. At $q = 1$, $\mathrm{ch}(C'_w)$ determines the vector $c_{\cdot,w} = (\phi^\lambda(C'_w(1)))_\lambda$ and conversely. Let $s = s_l = (l\ l{+}1)$, so that ws exchanges the entries in positions $l, l+1$. Suppose $sw < w < ws$. The Kazhdan–Lusztig multiplication rule gives $C'_w C'_s = C'_{ws} + \sum_{z \in M(w,s)} \mu(z,w) C'_z$ and $C'_s C'_w = (q^{1/2} + q^{-1/2}) C'_w$, where

$$M(w,s) = \{z < w :\ zs < z,\ \mu(z,w) \neq 0\}.$$

Since characters are traces, Abreu and Nigro [1, (3a)] obtain

$$(q^{1/2} + q^{-1/2})\ \mathrm{ch}(C'_w) = \mathrm{ch}(C'_{ws}) + \sum_{z \in M(w,s)} \mu(z,w)\ \mathrm{ch}(C'_z). \tag{7}$$

Definition 7.3. The step $w \to ws$ is *clean* if $M(w,s) = \emptyset$. Left steps $w \to sw$, with $ws < w < sw$, are defined in the same way.

Each $\mathrm{ch}(C'_z)$ is a non-zero Schur-positive symmetric function [16, Lemma 1.1], so at $q = 1$ the step is clean iff $c_{\cdot,ws} = 2c_{\cdot,w}$.

Theorem 7.4. *Let $y = y_0 > y_1 > \cdots > y_k$ with each y_{i-1} obtained from y_i by a clean step, and y_k smooth. Then $\mathrm{ch}(q^{\ell(y)/2} C'_y) = (1+q)^k\, \mathrm{ch}(q^{\ell(w')/2} C'_{w'})$ for a codominant w', and $\phi^\lambda(C'_y(1)) = 2^k \phi^\lambda(C'_{w'}(1)) \geq 0$ for every $\lambda \vdash n$.*

Proof. For a clean step, (7) reads $\mathrm{ch}(q^{\ell(ws)/2} C'_{ws}) = (1+q)\, \mathrm{ch}(q^{\ell(w)/2} C'_w)$. Induct on k; the case $k = 0$ is [1, Thm. 1.6] and Proposition 7.2. □

The case $k = 1$ is [1, Thm. 1.7]. Since a clean step gives $c_{\cdot,ws} = 2c_{\cdot,w}$ also when w is singular, Conjecture 1.2 holds for n iff $c_{\lambda,w} \geq 0$ for every singular w that is not the top of a clean step. This proves the second statement of Theorem 1.6. The rest of this section gives a local criterion for clean steps.

Definition 7.5. Let $w(l) < w(l+1)$. A *bad cover* of w at l is a permutation z covered by w in Bruhat order with $z(l) > z(l+1)$.

Write $a = w(l) < b = w(l{+}1)$. A cover $z = w\,(i\ j)$ can be bad only if $i = l{+}1 < j$, $w(j) < a$ and no $k \in (l{+}1, j)$ has $w(j) < w(k) < b$; or $i < l = j$, $w(i) > b$ and no $k \in (i, l)$ has $a < w(k) < w(i)$. Covers have $\mu = 1$, so a clean step has no bad cover.

Example 7.B. List $c_{\cdot,w}$ in the order $\lambda = (4), (3,1), (2,2), (2,1,1), (1^4)$. For the smooth $w = 3142$ and $l = 2$, $sw = 2143 < w < ws = 3412$, and $M(w,s) = \emptyset$: the step is clean, and indeed $c_{\cdot,3142} = (4,2,2,0,0)$ and $c_{\cdot,3412} = (8,4,4,0,0)$. For the smooth $w = 2341$ and $l = 1$, $sw =$

$1342 < w < ws = 3241$, and $z = 2143$ is a bad cover: it is covered by w and $z(1) > z(2)$. Here $M(w, s) = \{2143\}$, $c_{\cdot,2341} = (4, 2, 2, 0, 0)$, $c_{\cdot,3241} = (8, 4, 0, 0, 0)$ and $c_{\cdot,2143} = (0, 0, 4, 0, 0)$, so (7) at $q = 1$ reads $2(4, 2, 2, 0, 0) = (8, 4, 0, 0, 0) + (0, 0, 4, 0, 0)$ and the step is not clean. In line with [1, Prop. 3.1], the top 3412 of the clean step is singular, and the top 3241 of the other is smooth.

Proposition 7.6. *Let $z < w$ with $\ell(w) - \ell(z) = 3$ and $\mu(z, w) \neq 0$. Then w and z differ in exactly four positions $p_1 < p_2 < p_3 < p_4$, and either w restricted to them has pattern 3412 and z has pattern 1324, or w has pattern 4231 and z has pattern 2143. The coatoms of $[z, w]$ are $w\,(p_1\ p_3)$, $w\,(p_2\ p_3)$, $w\,(p_2\ p_4)$, $w\,(p_1\ p_4)$ in the first case, and $w\,(p_3\ p_4)$, $w\,(p_2\ p_4)$, $w\,(p_1\ p_3)$, $w\,(p_1\ p_2)$ in the second. All of them are covered by w.*

Proof. Since $P_{z,w} \neq 1$, X_w is singular at the point z. Every irreducible component of the singular locus of X_w has codimension at least 3, and the components of codimension exactly 3 are the X_z given by the two configurations above, with the regions they span empty [5, 25, 20, 12]. Because $\ell(w) - \ell(z) = 3$, X_z is such a component. The interval $[z, w]$ consists of the permutations that agree with w outside the four positions and whose restriction lies in $[1324, 3412]$, respectively $[2143, 4231]$, of S_4, which gives the coatoms. □




FIGURE 4. The two configurations of Proposition 7.6. A point $(p, w(p))$ is drawn in column p and row $w(p)$; black points are w at the positions $p_1 < \cdots < p_4$, and hollow points are z. In every pair $z < w$ with $\ell(w) - \ell(z) = 3$ and $\mu(z, w) \neq 0$ for $n \leq 6$ (1, 18 and 243 pairs of each type for $n = 4, 5, 6$), no other point of w lies in a shaded cell; the white cells may contain points. The square marks a value c at position $l + 1 = p_3 + 1$ in a white cell, the configuration of the first row of each block of the table in Lemma 7.7. On the left $v_1 < c < v_2$, and $y = w\,(p_2\ p_3)$ is a bad cover with $y(l) = v_4 > c$. On the right $v_3 < c < v_4$, and $y = w\,(p_1\ p_3)$ is a bad cover with $y(l) = v_4 > c$.

Lemma 7.7. *Let $w(l) < w(l + 1)$, and let $z < w$ with $\ell(w) - \ell(z) = 3$, $\mu(z, w) \neq 0$ and $z(l) > z(l + 1)$. Then w has a bad cover at l.*

Proof. Use the notation of Proposition 7.6 and Figure 4, let $v_1 < v_2 < v_3 < v_4$ be the values in positions $D = \{p_1, \ldots, p_4\}$, and let c be the common value of w and z at the position of $\{l, l + 1\}$ outside D, if there is one. Since the order of the entries in positions $l, l + 1$ is reversed between w and z, $\{l, l + 1\} \cap D \neq \emptyset$. If $l, l + 1 \in D$, they are consecutive in D, but in neither configuration does a consecutive pair increase in w and decrease in z. In the remaining cases, y is a coatom from Proposition 7.6, hence covered by w, with $y(l) > y(l + 1)$:

configuration	position	bad cover
$3412 \to 1324$	$l = p_3,\ v_1 < c < v_2$	$w\,(p_2\ p_3),\ \ y(l) = v_4$
	$l = p_4,\ v_2 < c < v_4$	$w\,(p_2\ p_4),\ \ y(l) = v_4$
	$l+1 = p_1,\ v_1 < c < v_3$	$w\,(p_1\ p_3),\ \ y(l+1) = v_1$
	$l+1 = p_2,\ v_3 < c < v_4$	$w\,(p_2\ p_3),\ \ y(l+1) = v_1$
$4231 \to 2143$	$l = p_3,\ v_3 < c < v_4$	$w\,(p_1\ p_3),\ \ y(l) = v_4$
	$l = p_4,\ v_1 < c < v_3$	$w\,(p_3\ p_4),\ \ y(l) = v_3$
	$l+1 = p_1,\ v_2 < c < v_4$	$w\,(p_1\ p_2),\ \ y(l+1) = v_2$
	$l+1 = p_2,\ v_1 < c < v_2$	$w\,(p_2\ p_4),\ \ y(l+1) = v_1$

The other positions of l or $l+1$ in D are excluded because there $w(l) < w(l+1)$ and $z(l) > z(l+1)$ cannot both hold. □

Proposition 7.8. *Let $sw < w < ws$ with $s = s_l$, and $\deg P_{e,w} \le 1$. Then the step $w \to ws$ is clean iff w has no bad cover at l.*

Proof. A clean step has no bad cover. Conversely, by monotonicity of Kazhdan–Lusztig polynomials [9, Cor. 3.7], $P_{e,w} - P_{z,w}$ has non-negative coefficients for $z \le w$, so $\deg P_{z,w} \le 1$. Hence $\mu(z,w) \neq 0$ forces $\ell(w) - \ell(z) \in \{1,3\}$. A $z \in M(w,s)$ at distance 1 is a bad cover, and one at distance 3 gives a bad cover by Lemma 7.7. □

For smooth w, Proposition 7.8 is contained in [1, Prop. 3.1]. The degree condition cannot simply be dropped: for $w = 256413$ and $l = 1$ there is no bad cover, but $z = 215436$ has $\ell(w) - \ell(z) = 5$ and $\mu(z,w) = 1$, so the step is not clean. Starting from smooth permutations, the steps certified by Proposition 7.8 reach 84%, 68%, 49%, 32% and 19% of the singular permutations for $n = 5, \dots, 9$ [26]. So the criterion handles a large but shrinking share of the singular permutations, and the singular w that are not the top of a clean step are where Conjecture 1.2 remains open.

8. Two further families of shapes

Lemma 8.1. *If $\lambda_1 \le 2$ and u contains 321, then $\phi^\lambda(C'_u(1)) = 0$.*

Proof. The map $\theta\colon \mathbb{C}[S_n] \to TL_n(2)$, $1 + s_i \mapsto t_i$, is a surjective algebra homomorphism. It kills $C'_u(1)$ exactly when u contains 321, and the $\theta(C'_w(1))$ for 321-avoiding w form a basis (Fan–Green [14]). So $\ker\theta = \operatorname{span}\{C'_u(1) : u \ni 321\}$. A two-sided ideal of $\mathbb{C}[S_n]$ is a sum of isotypic components. The ideal $\ker\theta$ contains $C'_{s_1s_2s_1}(1) = \sum_{v\in S_3} v$, which acts non-trivially on V^μ iff the restriction of χ^μ to $S_3 = \langle s_1, s_2\rangle$ contains the trivial character. By Young's rule this multiplicity is $\langle \chi^\mu, \eta^{(3,1^{n-3})}\rangle = K_{\mu,(3,1^{n-3})}$, which is non-zero iff $\mu_1 \ge 3$. So $\ker\theta$ contains every component with $\mu_1 \ge 3$. Since $\dim TL_n(2) = \mathrm{Cat}_n = \sum_{\mu_1\le 2}(f^\mu)^2$, comparing dimensions shows that $\ker\theta$ is exactly their sum. Finally, ϕ^λ involves only χ^μ with $\mu \trianglelefteq \lambda$, hence with $\mu_1 \le 2$, so it vanishes on $\ker\theta$. □

In the ring of symmetric functions, $m_{(2^{b+1},1^c)}\,p_1 = m_{(3,2^b,1^c)} + (b+2)\,m_{(2^{b+2},1^{c-1})} + (c+1)\,m_{(2^{b+1},1^{c+1})}$, where the middle term is absent when $c = 0$. Hence

$$\mathrm{Imm}_{(3,2^b,1^c)} = \sum_j x_{jj}\,\mathrm{Imm}_{(2^{b+1},1^c)}(x[\hat\jmath,\hat\jmath]) - (b+2)\,\mathrm{Imm}_{(2^{b+2},1^{c-1})} - (c+1)\,\mathrm{Imm}_{(2^{b+1},1^{c+1})}\,. \tag{8}$$

Proposition 8.2. *For all $b, c \ge 0$ and every $u \in S_{3+2b+c}$ that contains 321, $\phi^{(3,2^b,1^c)}(C'_u(1)) \ge 0$.*

Proof. At $u \ni 321$, the two-column terms of (8) vanish by Lemma 8.1. The first term is KL-positive: $\mathrm{Imm}_{(2^{b+1},1^c)}$ is a non-negative combination of Temperley–Lieb immanants [11], which are Kazhdan–Lusztig immanants [30], so each $x_{jj}\,\mathrm{Imm}_{(2^{b+1},1^c)}(x[\hat\jmath,\hat\jmath])$ is KL-positive by Theorem 3.1(1). □

So Conjecture 1.2 for the shapes $(3, 2^b, 1^c)$ reduces to 321-avoiding permutations. Those avoiding 312 or 231 are smooth and covered by Proposition 7.2. For 321-avoiding, hexagon-avoiding w with reduced word $s_{i_1} \cdots s_{i_l}$, Billey and Warrington [6, Thm. 1] give $C'_w(1) = (1+s_{i_1}) \cdots (1+s_{i_l})$, so $c_{\lambda,w} = \sum_{S \subseteq [l]} \phi^\lambda(\prod_{k \in S} s_{i_k})$.

9. Beyond class functions

The cones of Section 6 consist of class functions, and for these $C_{\mathrm{KL}} = C_{\mathrm{TNN}}$ when $n \leq 10$. Without the restriction to class functions this fails already for $n = 4$ (Rhoades–Skandera [30, §7]; Skandera and Zelevinsky give an explicit element [34, (5.1)]). We recall the setting, give examples for $n = 5$, and prove that the TNN-isolated permutations are exactly the smooth ones (Theorem 9.6). Let Π_n be the span of the permutation monomials x^v. Every $F \in \Pi_n$ is uniquely $F = \sum_w a_w \operatorname{Imm}_w$, and we put

$$C_{\mathrm{KL}}(\Pi_n) = \{F : \ a_w \geq 0 \ \forall w\}, \qquad C_{\mathrm{TNN}}(\Pi_n) = \{F : \ F \geq 0 \text{ on } \mathrm{TNN}_n\}.$$

Then $C_{\mathrm{KL}}(\Pi_n) \subseteq C_{\mathrm{TNN}}(\Pi_n)$, and the cones of Section 6 are their intersections with the immanants d_ϕ, $\phi \in \mathrm{CF}_n$. Call w *TNN-isolated* if every $F \in C_{\mathrm{TNN}}(\Pi_n)$ has $a_w \geq 0$. So $C_{\mathrm{KL}}(\Pi_n) = C_{\mathrm{TNN}}(\Pi_n)$ iff every w is TNN-isolated. Equivalently, since the evaluation vectors $(\operatorname{Imm}_v(A))_v$, $A \in \mathrm{TNN}_n$, generate the cone dual to $C_{\mathrm{TNN}}(\Pi_n)$, w is TNN-isolated iff the coordinate vector e_w is a limit of positive combinations of evaluation vectors.

These cones go back to Stembridge [37, §5], who studied the cone of total positivity in $\mathbb{R}S_n$, whose dual is $C_{\mathrm{TNN}}(\Pi_n)$. Rhoades and Skandera [30, §7] noted $C_{\mathrm{KL}}(\Pi_n) \subsetneq C_{\mathrm{TNN}}(\Pi_n)$ for $n \geq 4$, observed that many Imm_w seem to span extreme rays of $C_{\mathrm{TNN}}(\Pi_n)$, and asked whether $a_w \geq 0$ for every $F \in C_{\mathrm{TNN}}(\Pi_n)$ whenever X_w is smooth [30, Question 7.1]. Skandera answered this question [34] and asked for the precise relation between the cones [34, Problem 5.8].

Theorem 9.1 (Skandera [34, Thms. 5.4, 5.5]; asked in [30, Question 7.1])**.** *For every n, every smooth $w \in S_n$ is TNN-isolated: there is $A_w \in \mathrm{TNN}_n$ with $\operatorname{Imm}_v(A_w) = \delta_{v,w}$ for all v.*

Skandera's matrix A_w is the path matrix of a planar network built from a factorisation of w. For codominant w, Proposition 7.1 gives such a matrix directly, the antiadjacency matrix of a unit interval order.

A *chart* is a factorisation into elementary factors $I + tE_{i+1,i}$ and $I + tE_{i,i+1}$ whose lower letters form a reduced word of w_0, as do its upper letters. By Fomin–Zelevinsky [15], positive parameters give every totally positive matrix up to a positive diagonal factor, which multiplies every element of Π_n by its determinant. Since totally positive matrices are dense in TNN_n, an $F \in \Pi_n$ whose expansion in the parameters of some chart has only non-negative coefficients is non-negative on TNN_n.

Theorem 9.2. *The elements*

$$\operatorname{Imm}_{15432} + \operatorname{Imm}_{43215} - \operatorname{Imm}_{45312}, \qquad \operatorname{Imm}_{21543} + \operatorname{Imm}_{54321} - \operatorname{Imm}_{52431},$$
$$\operatorname{Imm}_{32154} + \operatorname{Imm}_{54321} - \operatorname{Imm}_{53241}$$

of Π_5 are non-negative on TNN_5 and are not KL-positive. So $C_{\mathrm{KL}}(\Pi_5) \subsetneq C_{\mathrm{TNN}}(\Pi_5)$, and 45312, 52431, 53241 are not TNN-isolated.

Proof. For each element there is a chart in which its expansion has only non-negative coefficients: the standard chart (lower word $s_4\, s_3 s_4\, s_2 s_3 s_4\, s_1 s_2 s_3 s_4$, followed by its reverse in the upper factors) for the first two, and for the third the chart

$$L_1 L_2 U_1 L_3 U_3 L_1 U_2 L_2 U_4 U_3 L_1 L_4 L_3 U_1 L_2 L_1 U_2 U_4 U_1 U_3,$$

with $L_i(t) = I + tE_{i+1,i}$, $U_i(t) = I + tE_{i,i+1}$ and one parameter per factor; its lower letters $1, 2, 3, 1, 2, 1, 4, 3, 2, 1$ and upper letters $1, 3, 2, 4, 3, 1, 2, 4, 1, 3$ are reduced words of w_0. This is

verified in exact integer arithmetic: the three expansions have 870, 916 and 10,611 monomials, all with coefficients between 1 and 16 (`paper/hooks_kl_examples.py` in [27]). They are not KL-positive because the coefficients -1 are coefficients in the basis $\{\mathrm{Imm}_w\}$. □

All three non-isolated permutations are singular, as they must be by Theorem 9.1. We now show that Skandera's theorem is sharp for every n: no singular w is TNN-isolated. The proof lifts two elements of $C_{\mathrm{TNN}}(\Pi_4)$ along an occurrence of 3412 or 4231 in w, and controls the lift with Theorem 3.1.

Proposition 9.3. *On* TNN_4*,*

$$\mathrm{Imm}_{3412}^2 \le \mathrm{Imm}_{1432}\,\mathrm{Imm}_{3214}, \qquad \mathrm{Imm}_{4231}^2 \le 3\,\mathrm{Imm}_{2143}\,\mathrm{Imm}_{4321},$$

where $\mathrm{Imm}_{4321} = x_{14}x_{23}x_{32}x_{41}$*. Hence* $F = \frac{1}{2}(\mathrm{Imm}_{1432} + \mathrm{Imm}_{3214}) - \mathrm{Imm}_{3412}$ *and* $G_\varepsilon = \varepsilon\,\mathrm{Imm}_{2143} + \frac{3}{4\varepsilon}\,\mathrm{Imm}_{4321} - \mathrm{Imm}_{4231}$*, for every* $\varepsilon > 0$*, lie in* $C_{\mathrm{TNN}}(\Pi_4)$*.*

Proof. Use the chart (Figure 5)

$$L_3(t_1)L_2(t_2)L_1(t_3)L_3(t_4)L_2(t_5)L_3(t_6)\,U_3(t_7)U_2(t_8)U_1(t_9)U_3(t_{10})U_2(t_{11})U_3(t_{12}),$$

where $L_i(t) = I + tE_{i+1,i}$ and $U_i(t) = I + tE_{i,i+1}$; the word $s_3s_2s_1s_3s_2s_3$ is a reduced word of w_0. In this chart Imm_{1432}, Imm_{3214}, Imm_{2143} and Imm_{4321} have only positive coefficients, $\mathrm{Imm}_{3412} = m$ is a single monomial, and $\mathrm{Imm}_{4231} = m_1 + m_2 + m_3$, where

$$m = t_2t_3t_4t_5t_8t_9t_{10}t_{11}, \qquad m_1 = t_1t_2t_3^2t_5t_8t_9^2t_{11}t_{12},$$
$$m_2 = t_1t_2t_3^2t_9^2t_{11}t_{12}, \qquad m_3 = t_1t_2t_3t_9t_{11}t_{12}.$$

The monomial $a = t_2t_4t_5t_8t_{10}t_{11}$ of Imm_{1432} and the monomial $b = t_2t_3^2t_4t_5t_8t_9^2t_{10}t_{11}$ of Imm_{3214} satisfy $m^2 = ab$, so $\mathrm{Imm}_{3412}^2 \le \mathrm{Imm}_{1432}\,\mathrm{Imm}_{3214}$ for positive parameters. The monomials $a_1 = t_1t_3t_5^2t_8^2t_9t_{12}$ and $a_2 = t_1t_3t_9t_{12}$ of Imm_{2143} and $b_1 = t_1t_2^2t_3^3t_9^3t_{11}^2t_{12}$ and $b_2 = t_1t_2^2t_3t_9t_{11}^2t_{12}$ of Imm_{4321} satisfy $m_1^2 = a_1b_1$, $m_2^2 = a_2b_1$ and $m_3^2 = a_2b_2$. By the Cauchy–Schwarz inequality,

$$\mathrm{Imm}_{4231}^2 \le 3(m_1^2 + m_2^2 + m_3^2) \le 3(a_1 + a_2)(b_1 + b_2) \le 3\,\mathrm{Imm}_{2143}\,\mathrm{Imm}_{4321}\,.$$

A positive diagonal factor multiplies both sides of each inequality by $\det(D)^2$, so both hold on the totally positive matrices, and by density on TNN_4. The last statement follows from $\sqrt{XY} \le \frac{X+Y}{2}$ and $\sqrt{3XY} \le \varepsilon X + \frac{3}{4\varepsilon}Y$ for $X, Y \ge 0$, since Kazhdan–Lusztig immanants are non-negative on TNN_4. □




FIGURE 5. The chart of Proposition 9.3 as a planar network. Paths run from the sources $1, \dots, 4$ on the left to the sinks $1', \dots, 4'$ on the right; an edge labelled t_k from line a to line b is the factor $I + t_kE_{a,b}$, and horizontal edges have weight 1. The (i, j) entry of the product of the twelve factors is the sum of the weights of the paths from i to j' (Lindström). The first six edges are the lower factors $L_i(t) = I + tE_{i+1,i}$, and the last six the upper factors $U_i(t) = I + tE_{i,i+1}$.

The Skandera–Zelevinsky element is $\mathrm{Imm}_{3214} + \mathrm{Imm}_{1432} - \mathrm{Imm}_{3412} = 2F + \mathrm{Imm}_{3412}$, so F is the stronger statement.

Now fix $w \in S_n$ and $I \subseteq [n]$ with $|I| = k$, and put $T = w(I)$. As in the proof of Theorem 3.1, $\varphi(\sigma) = (\sigma_I, \sigma_{I^c})$ for $\sigma(I) = T$. Let $p = w_I$ and $w_c = w_{I^c}$, so that w contains the pattern p in the positions I. For $s \in S_k$ let σ_s be the permutation with $\sigma_s(I) = T$ and $\varphi(\sigma_s) = (s, w_c)$: it agrees with w outside I and has pattern s in the positions I. For $u \in S_k$ let $A(u)$ be the coefficient of Imm_w in $\mathrm{Imm}_u(x[I,T])\,\mathrm{Imm}_{w_c}(x[I^c,T^c])$. By Theorem 3.1, $A(u) \in \mathbb{N}$.

Lemma 9.4. (1) *$A(p) = 1$, and $A(u) = 0$ for every $u > p$.*
(2) *For every $u \in S_k$, $A(u) + \sum_{u'>u} A(u')\,P_{u,u'}(1) \le P_{\sigma_u,w}(1)$.*

Proof. By the proof of Theorem 3.1, $A(u) = a_y$ for the y with $\varphi(y) = (u, w_c)$. Since the Kazhdan–Lusztig polynomials of W' are products, (3) at $\sigma = \sigma_u$ reads

$$P_{\sigma_u,w}(1) = \sum_y a_y\,P_{u,\varphi(y)_1}(1)\,P_{w_c,\varphi(y)_2}(1),$$

a sum of non-negative terms. Keeping only the y with $\varphi(y)_2 = w_c$ gives (2).

For (1), the map φ^{-1} is order-preserving from $S_k \times S_{n-k}$ to the coset $\{\sigma : \sigma(I) = T\}$. Indeed, if $s(a) < s(b)$ for $a < b$, then multiplying s on the right by the transposition $(a\ b)$ corresponds to multiplying σ on the right by the transposition of the a-th and b-th elements of I, whose entries in σ also increase; the same holds in I^c. By the definition of Kazhdan–Lusztig immanants, the product $\mathrm{Imm}_p(x[I,T])\,\mathrm{Imm}_{w_c}(x[I^c,T^c]) = \sum_\sigma f(\sigma)x^\sigma$ has $f(\sigma) \neq 0$ only if $\sigma_I \ge p$ and $\sigma_{I^c} \ge w_c$, hence only if $\sigma \ge w$, and $f(w) = 1$. Write the product as $\sum_y c_y\,\mathrm{Imm}_y$. Since Imm_y involves only monomials x^v with $v \ge y$, a Bruhat-minimal y with $c_y \neq 0$ has $c_y = f(y)$, so $y \ge w$. Hence $c_y \neq 0$ implies $y \ge w$, and comparing the coefficients of x^w gives $A(p) = c_w = f(w) = 1$. Finally, (2) for $u = p$, where $\sigma_p = w$, reads $1 + \sum_{u'>p} A(u')P_{p,u'}(1) \le 1$. Since $P_{p,u'}(1) \ge 1$, $A(u') = 0$ for $u' > p$. □

Lemma 9.5. *Let $i < j$ with $w(i) > w(j)$, and let R be the set of $k \in (i,j)$ with $w(j) < w(k) < w(i)$. Then the Schubert variety X_w is smooth at $w\,(i\ j)$ iff $w(k) > w(l)$ for all $k < l$ in R, that is, iff w has no occurrence of 4231 in positions $i < k < l < j$. In that case $P_{w(i\,j),w} = 1$.*

Proof. Put $x = w\,(i\ j)$ and $r_y(p,v) = \#\{p' \le p : y(p') \ge v\}$. Then $y \le w$ iff $r_y \le r_w$ pointwise [7, Thm. 2.1.5]. If $p < q$ and $y(p) < y(q)$, then $r_{y(p\,q)} = r_y + \mathbf{1}_{[p,q-1]\times[y(p)+1,y(q)]}$. Hence $r_x = r_w - \mathbf{1}_B$ with $B = [i, j-1] \times [w(j)+1, w(i)]$, and for $p < q$ with $x(p) < x(q)$,

$$x\,(p\ q) \le w \iff [p, q-1] \times [x(p)+1, x(q)] \subseteq B,$$

that is, iff the points $(p, x(p))$ and $(q, x(q))$ both lie in the box $[i,j] \times [w(j), w(i)]$. The points of x in this box are $(i, w(j))$, $(j, w(i))$ and $(k, w(k))$, $k \in R$. So the number of such pairs is $(|R|+1)+|R|+\mathrm{asc}(R)$, where $\mathrm{asc}(R)$ is the number of pairs $k < l$ in R with $w(k) < w(l)$. All $\ell(x)$ transpositions $(p\,q)$ with $x(p) > x(q)$ give elements below $x \le w$. By Lakshmibai and Seshadri [24], the tangent space of X_w at x has dimension $\#\{(p,q) : p < q,\ x\,(p\,q) \le w\} = \ell(x)+2|R|+1+\mathrm{asc}(R)$, while $\dim X_w = \ell(w) = \ell(x) + 2|R| + 1$. So X_w is smooth at x iff $\mathrm{asc}(R) = 0$, and then $P_{x,w} = 1$ [22]. A pair $k < l$ in R with $w(k) < w(l)$ is exactly an occurrence (i,k,l,j) of 4231. □

We include a proof of Lemma 9.5 for completeness. It is a direct consequence of the tangent space formula; Figure 6 shows both cases.

Theorem 9.6. *For every n, a permutation $w \in S_n$ is TNN-isolated iff the Schubert variety X_w is smooth, that is, iff w avoids 3412 and 4231 [23]. Equivalently, for every singular w there is an element of $C_{\mathrm{TNN}}(\Pi_n)$ whose coefficient at Imm_w is negative.*




FIGURE 6. Lemma 9.5 for $i = 2$, $j = 5$. The point $(p, w(p))$ is drawn in column p and row $w(p)$; black points are w, and the hollow points are the two entries of $x = w\,(2\ 5)$ that differ from w. The shaded box is $[i, j] \times [w(j), w(i)]$, and $R = \{3, 4\}$. Left: R is decreasing, so $\mathrm{asc}(R) = 0$, and the tangent space at $x = 12435$ has dimension $\ell(x) + 2|R| + 1 = 6 = \ell(w)$: X_w is smooth at x. Right: the ascent $(3, 4)$ in R is the occurrence $(2, 3, 4, 5)$ of 4231, the tangent space at $x = e$ has dimension $0 + 5 + 1 = 6 > \ell(w) = 5$, and $P_{e,w} = 1 + q$.

Proof. Smooth permutations are TNN-isolated by Theorem 9.1. Let w be singular. For $W = \sum_u c_u \operatorname{Imm}_u \in C_{\mathrm{TNN}}(\Pi_4)$ and an occurrence of a pattern of length 4 in positions I, put $H = W(x[I, T]) \operatorname{Imm}_{w_c}(x[I^c, T^c])$. Submatrices of totally non-negative matrices are totally non-negative, and $\operatorname{Imm}_{w_c} \geq 0$ on TNN_{n-4}, so $H \in C_{\mathrm{TNN}}(\Pi_n)$. Its coefficient at Imm_w is $\sum_u c_u A(u)$.

If w contains 4231, take an occurrence of 4231 and $W = G_\varepsilon$. By Lemma 9.4(1), with $4321 > 4231$, the coefficient is $\varepsilon A(2143) + \frac{3}{4\varepsilon} A(4321) - A(4231) = \varepsilon A(2143) - 1$, which is negative for small $\varepsilon > 0$.


Otherwise w contains 3412 and avoids 4231. Take any occurrence $i_1 < i_2 < i_3 < i_4$ of 3412 and $W = F$. Here $\sigma_{1432} = w\,(i_1\ i_3)$ and $\sigma_{3214} = w\,(i_2\ i_4)$. Since w avoids 4231, Lemma 9.5 gives $P_{\sigma_{1432},w} = P_{\sigma_{3214},w} = 1$. Lemma 9.4(2) for $u = 1432 < 3412$, together with $A(3412) = 1$ and $P_{1432,3412} = 1$, gives $A(1432) + 1 \leq 1$, so $A(1432) = 0$, and likewise $A(3214) = 0$. The coefficient is $\frac{1}{2}A(1432) + \frac{1}{2}A(3214) - A(3412) = -1$. □

Example 9.A. Let $w = 45312$, one of the permutations of Theorem 9.2. It avoids 4231 and contains 3412 in the positions $I = \{1, 2, 4, 5\}$, with $T = w(I) = \{1, 2, 4, 5\}$, $p = 3412$ and $w_c = e \in S_1$ (Figure 7). Then $\sigma_{1432} = w\,(1\ 4) = 15342$ and $\sigma_{3214} = w\,(2\ 5) = 42315$, and $P_{\sigma_{1432},w} = P_{\sigma_{3214},w} = 1$, as Lemma 9.5 predicts. Computing the products directly, the coefficient $A(u)$ of Imm_w in $\operatorname{Imm}_u(x[I, T])\, x_{33}$ is 1 for $u = 3412$ and 0 for every other $u \in S_4$. So

$$H = \big(\tfrac{1}{2} \operatorname{Imm}_{1432} + \tfrac{1}{2} \operatorname{Imm}_{3214} - \operatorname{Imm}_{3412}\big)(x[I, T])\ x_{33}$$

lies in $C_{\mathrm{TNN}}(\Pi_5)$, and its coefficient at $\operatorname{Imm}_{45312}$ is -1; it is the only negative coefficient of H. For the 4231 case, $w = 52431$ contains 4231 in the positions $\{1, 2, 4, 5\}$, and there $A(4231) = 1$ and $A(u) = 0$ for $u \neq 4231$, so the lift of G_ε has coefficient -1 at $\operatorname{Imm}_{52431}$ for every $\varepsilon > 0$.

So the full-span cones differ in the coordinate of every singular w, while for class functions $C_{\mathrm{KL}} = C_{\mathrm{TNN}}$ when $n \leq 10$ [26]. The companion paper [26] proves the cases $n = 4, 5$ of Theorem 9.6 by a different method, a vertex criterion on Newton polytopes. It also checks the proof by exact computation. For every singular w with $n \leq 10$, the lifted element ($2F$ for 3412, and $\operatorname{Imm}_{2143} + \operatorname{Imm}_{4321} - \operatorname{Imm}_{4231} = G_1 + \frac{1}{4} \operatorname{Imm}_{4321}$ for 4231) has a negative coefficient at Imm_w already at the first occurrence tried. Lemma 9.5 is checked for every inversion of every w with $n \leq 10$. For 4231 the proof works at every occurrence. For 3412 the elements $\varepsilon \operatorname{Imm}_{1432} + \frac{1}{4\varepsilon} \operatorname{Imm}_{3214} - \operatorname{Imm}_{3412}$, which lie in $C_{\mathrm{TNN}}(\Pi_4)$ by Proposition 9.3, show that it suffices that one of $w\,(i_1\ i_3)$ and $w\,(i_2\ i_4)$ is a smooth point of X_w.



$w = 45312$

FIGURE 7. The occurrence of 3412 in $w = 45312$ used in Example 9.A: positions $I = \{1, 2, 4, 5\}$ (solid lines) and values $T = w(I) = \{1, 2, 4, 5\}$ (dashed lines). The remaining point $(3, 3)$ gives $w_c = e \in S_1$.

10. Conjectures and open problems

10.1. **Explicit expansions for hooks.** Theorem 1.3 does not make the Kazhdan–Lusztig coefficients of $f_i^{(a)}$ explicit, except for $a = 2$. Computation suggests that for $a \geq 2$, $f_i^{(a)}(C'_w(1)) = 0$ unless w moves a single block of consecutive letters containing i (observed for $n \leq 8$). For $a = 3$, (5) gives $f_i^{(3)}(C'_w(1)) = \sum_{j \neq i} B_{ij}(w) - f_i^{(2)}(C'_w(1))$, where $B_{ij}(w) \geq 0$ is the coefficient of Imm_w in $x_{ij}x_{ji} \det x[\{i,j\}^c, \{i,j\}^c]$.

Conjecture 10.1. *Let $i < j$.*

(1) *If $j - i \geq 2$, let $Q = \{i-1, i, j, j+1\} \cap [n]$. Then $B_{ij}(w) = 1$ if w fixes $[n] \setminus Q$ and its pattern on Q lies in the Bruhat interval $[(i\ j), (\min Q\ \max Q)]$ of $S_{|Q|}$, and $B_{ij}(w) = 0$ otherwise. For interior pairs this interval is $[1324, 4231]$, with 16 elements.*

(2) *If $j = i+1$, B_{ij} equals 1 on the patterns $\geq (i\ i{+}1)$ on $\{i-1, \ldots, i+2\}$, except the longest element when $|Q| = 4$, and 2 on the pattern 3412. It is also 1 on the 5-cycles*

$$(i{-}2, i, i{+}2, i{-}1, i{+}1),\ (i{-}2, i{+}1, i{-}1, i{+}2, i) \qquad (i \geq 3),$$
$$(i{-}1, i{+}1, i{+}3, i, i{+}2),\ (i{-}1, i{+}2, i, i{+}3, i{+}1) \qquad (i + 3 \leq n),$$

and 0 elsewhere.

Table 1 lists the support of $B_{i,i+1}$ in the interior case $3 \leq i \leq n-3$, where all three kinds of term occur.

w	$B_{i,i+1}(w)$	number
fixes $[n] \setminus Q$, pattern on Q in $[1324, 4321)$, not 3412 1324, 1342, 1423, 1432, 2314, 2341, 2413, 2431, 3124, 3142, 3214, 3241, 3421, 4123, 4132, 4213, 4231, 4312	1	18
fixes $[n] \setminus Q$, pattern 3412 on Q	2	1
one of the four 5-cycles of Conjecture 10.1(2)	1	4
any other w	0	

TABLE 1. The conjectured support of $B_{i,i+1}$ for $3 \leq i \leq n-3$, with $Q = \{i-1, i, i+1, i+2\}$. For $n = 6$, $i = 3$ it has 23 elements, and the four 5-cycles are 345126, 451236, 145623, 156234.

This matches the computation for every pair (i, j) with $n \leq 8$. Two parts are proved: $B_{ij}(w) = 0$ if w has a left or right descent s_k with $\{k, k{+}1\} \cap \{i, j\} = \emptyset$; and B_{ij} is supported on minimal double coset representatives for the Young subgroup with blocks $[1, i-1], \{i\}, [i+1, j-1], \{j\}, [j+1, n]$.

With Lemma 4.1, Conjecture 10.1 gives a closed form for $\mathrm{Imm}_{(3,1^{n-3})}$ in the Kazhdan–Lusztig basis. It would follow from an inertness property: for a cycle γ on $T \subset [n]$, a point $k \notin T$ at distance at least 2 from T appears to be a fixed point of every w in the support of $x_\gamma \det x[T^c, T^c]$, with deletion of k giving the corresponding coefficients in S_{n-1}, if $\min T < k < \max T$, or no two elements of T are adjacent, or γ is an adjacent transposition and k is at distance at least 3 (checked for 2- and 3-cycles with $n \le 8$).

10.2. **A rooted refinement.** Cut the cycles of $\sigma \in S_n$ into *arcs*, runs $v, \sigma(v), \dots, \sigma^{l-1}(v)$ of consecutive elements of a cycle. Expanding each p_k in the h-basis by brick tabloids gives $\phi^\lambda(\sigma) = (-1)^{n-\ell(\lambda)} \operatorname{sgn}(\sigma) N_\lambda(\sigma)$, where $N_\lambda(\sigma)$ is the number of cuttings into arcs of lengths λ. For a vertex v, an integer $a \ge 1$ and $\mu \vdash n-a$, let $G_v^{(a;\mu)}(\sigma) = (-1)^{n-1-\ell(\mu)} \operatorname{sgn}(\sigma) M$, where M is the number of cuttings with an arc of length a starting at v and the other arcs of lengths μ, the arcs of equal length ≥ 2 being ordered. Sorting the cuttings by the arc through v gives, for every v,

$$\phi^\lambda = \sum_a \frac{a}{\operatorname{aut}(\lambda \setminus a)} \, G_v^{(a;\lambda\setminus a)},$$

the sum over the distinct parts a of λ, with $\operatorname{aut}(\mu)$ the product of the factorials of the multiplicities of the parts ≥ 2 of μ. For hooks and $a = \lambda_1$ these are the per-vertex functions of Theorem 1.3.

Conjecture 10.2. *For $v \in \{1, n\}$, every $a \ge 1$, every $\mu \vdash n-a$ and every $w \in S_n$, $G_v^{(a;\mu)}(C'_w(1)) \ge 0$.*

Conjecture 10.2 implies Conjecture 1.2. It is a combinatorial form, at $q = 1$, of a conjecture of Abreu and Nigro. For every w, the decomposition theorem for the map $Y_w(X) \to \mathbb{P}^{n-1}$, $V_\bullet \mapsto V_1$, from a regular semisimple Lusztig variety splits $\operatorname{ch} IH^*(Y_w(X)) = \operatorname{ch}(q^{\ell(w)/2} C'_w)$ as $\sum_i [n-i]_q h_{n-i} \operatorname{ch}(\rho_{i,w})$ for representations $\rho_{i,w}$ of S_i [2, Cor. 1.9], and they conjecture that the local systems come from permutation representations [2, Conj. 1.4]. Our decomposition has the same shape, $\operatorname{ch}(C'_w(1)) = \sum_a a\, h_a \sum_\mu G_1^{(a;\mu)}(C'_w(1)) \operatorname{aut}(\mu)^{-1} h_\mu$. For codominant w the pieces agree with Abreu–Nigro's g-functions [3] (checked for $n \le 6$, and in graded form for $n \le 5$), which are e-positive at $q = 1$ by Huh, Hwang, Kim, Kim and Oh [19]. So Conjecture 10.2 holds for codominant w. We checked it for every w with $n \le 8$. At the interior vertices $1 < v < n$ it fails, already for $n = 3$. The pieces $G_1^{(a;\mu)}(C'_w(1))/\operatorname{aut}(\mu)$ are integers for $n \le 6$, and for $n \le 6$ they lift to palindromic graded pieces in $\mathbb{N}[q]$ compatible with $\operatorname{ch}(q^{\ell(w)/2} C'_w)$, as the decomposition theorem predicts. Two structural facts hold for every n. If $sw < w < ws$ and the step is clean, then $G_v^{(a;\mu)}(C'_{ws}(1)) = G_v^{(a;\mu)}(C'_w(1)) + G_{s(v)}^{(a;\mu)}(C'_w(1))$, because $C'_w(1)\, s = s\, C'_w(1)\, s$ and conjugation by s relabels the vertices. For $s \ne s_1$, which fixes vertex 1, the functions $G_1^{(a;\mu)}$ satisfy the analogue at $q = 1$ of the relation (7). For $n \le 6$ these relations determine them from their values at codominant w.

10.3. **Problems.** Conjectures 10.1 and 10.2 are concrete proposals supported by computation. Of the problems below, the first four are the main directions; the others are narrower or more exploratory targets.

(1) Find a structural reason why, for class functions, KL-positivity and TNN-positivity should agree ($C_{\mathrm{KL}} = C_{\mathrm{TNN}}$), independently of Conjecture 1.2. This would make Conjectures 1.1 and 1.2 equivalent.
(2) Prove Conjecture 10.2, for example by a restricted modular law for the vertex-1 functions in the spirit of [19], or by identifying the permutation representations of [2, Conj. 1.4]. The hook-style induction of Section 4 does not extend: for non-hook λ the coset pieces of $G_v^{(a;\mu)}$ are not KL-positive.

(3) Prove Haiman's q-version for hooks, $\phi_q^{(a,1^{n-a})}(q^{\ell(w)/2}C'_w(q)) \in \mathbb{N}[q]$. The proof of Theorem 1.3 works at $q = 1$ only. These polynomials are automatically symmetric about $q^{\ell(w)/2}$, because C'_w is invariant under the bar involution; the full conjecture [16, Conj. 2.1] also asks that they be unimodal. Haiman verified both for $n \leq 7$ [16], and the companion paper for $n \leq 10$ [26].

(4) Clean steps.
   (a) Describe the singular w that are not the top of a clean step, and prove $c_{\lambda,w} \geq 0$ for them; by Theorem 7.4 this is equivalent to Conjecture 1.2.
   (b) Replace the condition $\deg P_{e,w} \leq 1$ in Proposition 7.8 by a pattern condition.

(5) Prove $\phi^{(3,2^b,1^c)}(C'_w(1)) \geq 0$ for 321-avoiding w containing both 312 and 231; by Proposition 8.2 this is Conjecture 1.2 for these shapes.

(6) Prove that $\theta_\ell = \sum_{\ell(\mu)=\ell} \phi^\mu$ is KL-positive for every n (Skandera's Problem 30 [35], at $q = 1$). It holds at smooth w by Proposition 7.2, and for $n \leq 10$, since θ_ℓ is a sum of monomial characters, by [26]. Since $\sum_\ell t^\ell \theta_\ell(\sigma) = \prod_C (1 - (1-t)^{|C|})$ over the cycles C of σ, $\theta_\ell = \sum_{|T|=n-\ell} g_T$ with $g_T(\sigma) = (-1)^{|T|}\operatorname{sgn}(\sigma)[\text{no cycle of } \sigma \text{ lies in } T]$, and every g_T is KL-positive for $n \leq 8$.

(7) Describe the cone $C_{\mathrm{TNN}}(\Pi_n)$, for example its extreme rays (Stembridge [37, Question 5.4]; Skandera [34, Problem 5.8]). By Theorem 9.6, $C_{\mathrm{KL}}(\Pi_n)$ and $C_{\mathrm{TNN}}(\Pi_n)$ differ in the coordinate of every singular w. A first step is to find the extreme rays for $n = 4, 5$ and look for families that persist for all n.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this work the author used Claude (Anthropic), through Claude Code, to write and test the software, to run and cross-check the computations, to search and summarise the literature, to check arguments against computed examples, and to draft and edit the text. After using this tool, the author reviewed and edited the content as needed and takes full responsibility for the content of the published article.

References

[1] A. Abreu and A. Nigro, An update on Haiman's conjectures, Forum Math. Sigma 12 (2024), e86; arXiv:2206.00073v2.

[2] A. Abreu and A. Nigro, Parabolic Lusztig varieties and chromatic symmetric functions, J. Math. Pures Appl. 203 (2025), 103771, doi:10.1016/j.matpur.2025.103771; arXiv:2212.13497v1 (numbering cited from the arXiv version).

[3] A. Abreu and A. Nigro, Splitting the cohomology of Hessenberg varieties and e-positivity of chromatic symmetric functions, arXiv:2304.10644v1.

[4] S. Billey and T. Braden, Lower bounds for Kazhdan–Lusztig polynomials from patterns, Transform. Groups 8 (2003), 321–332.

[5] S. Billey and G. Warrington, Maximal singular loci of Schubert varieties in $SL(n)/B$, Trans. Amer. Math. Soc. 355 (2003), 3915–3945.

[6] S. Billey and G. Warrington, Kazhdan–Lusztig polynomials for 321-hexagon-avoiding permutations, J. Algebraic Combin. 13 (2001), 111–136.

[7] A. Björner and F. Brenti, Combinatorics of Coxeter Groups, Grad. Texts in Math. 231, Springer, 2005.

[8] T. Braden, Hyperbolic localization of intersection cohomology, Transform. Groups 8 (2003), 209–216.

[9] T. Braden and R. MacPherson, From moment graphs to intersection cohomology, Math. Ann. 321 (2001), 533–551.

[10] S. Clearman, M. Hyatt, B. Shelton and M. Skandera, Evaluations of Hecke algebra traces at Kazhdan–Lusztig basis elements, Electron. J. Combin. 23(2) (2016), P2.7.

[11] S. Clearman, B. Shelton and M. Skandera, Path tableaux and combinatorial interpretations of immanants for class functions on S_n, Discrete Math. Theor. Comput. Sci. Proc. AO (2011), 233–244.

[12] A. Cortez, Singularités génériques et quasi-résolutions des variétés de Schubert pour le groupe linéaire, Adv. Math. 178 (2003), 396–445.
[13] X. Dong, T. Wu and X. Gao, On the nonnegativity of monomial immanants for hook partitions, arXiv:2609.36665v1.
[14] C. K. Fan and R. M. Green, Monomials and Temperley–Lieb algebras, J. Algebra 190 (1997), 498–517.
[15] S. Fomin and A. Zelevinsky, Double Bruhat cells and total positivity, J. Amer. Math. Soc. 12 (1999), 335–380.
[16] M. Haiman, Hecke algebra characters and immanant conjectures, J. Amer. Math. Soc. 6 (1993), 569–595.
[17] P. Heyfron, Immanant dominance orderings for hook partitions, Linear Multilinear Algebra 24 (1988), 65–78.
[18] T. Hikita, A proof of the Stanley–Stembridge conjecture, J. Amer. Math. Soc., published electronically 28 August 2026, doi:10.1090/jams/1090; arXiv:2410.12758.
[19] J. Huh, B.-H. Hwang, D. Kim, J. S. Kim and J. Oh, Refinement of Hikita's e-positivity theorem via Abreu–Nigro's g-functions and restricted modular law, arXiv:2504.09123v1.
[20] C. Kassel, A. Lascoux and C. Reutenauer, The singular locus of a Schubert variety, J. Algebra 269 (2003), 74–108.
[21] D. Kazhdan and G. Lusztig, Representations of Coxeter groups and Hecke algebras, Invent. Math. 53 (1979), 165–184.
[22] D. Kazhdan and G. Lusztig, Schubert varieties and Poincaré duality, in: Geometry of the Laplace Operator, Proc. Sympos. Pure Math. 36, Amer. Math. Soc., 1980, 185–203.
[23] V. Lakshmibai and B. Sandhya, Criterion for smoothness of Schubert varieties in $SL(n)/B$, Proc. Indian Acad. Sci. Math. Sci. 100 (1990), 45–52.
[24] V. Lakshmibai and C. S. Seshadri, Singular locus of a Schubert variety, Bull. Amer. Math. Soc. (N.S.) 11 (1984), 363–366.
[25] L. Manivel, Le lieu singulier des variétés de Schubert, Internat. Math. Res. Notices 2001, no. 16, 849–871.
[26] K. McNulty, Haiman's conjecture for $n \leq 10$: Kazhdan–Lusztig positivity of monomial immanants, preprint, 2026. Available at `https://github.com/keithmcnulty/haiman-immanants-papers-n10-hooks`.
[27] K. McNulty, `keithmcnulty/haiman-immanants-papers-n10-hooks`: code and data for the papers, version v1.4.0, software, Zenodo, 2026, doi:10.5281/zenodo.23730750.
[28] T. H. Pate, Partitions, irreducible characters, and inequalities for generalized matrix functions, Trans. Amer. Math. Soc. 325 (1991), 875–894.
[29] B. Rhoades and M. Skandera, Temperley–Lieb immanants, Ann. Comb. 9 (2005), 451–494.
[30] B. Rhoades and M. Skandera, Kazhdan–Lusztig immanants and products of matrix minors, J. Algebra 304 (2006), 793–811.
[31] B. Rhoades and M. Skandera, Bitableaux and zero sets of dual canonical basis elements, Ann. Comb. 15 (2011), 499–528.
[32] I. Schur, Über endliche Gruppen und Hermitesche Formen, Math. Z. 1 (1918), 184–207.
[33] J. Sjöstrand, Bruhat intervals as rooks on skew Ferrers boards, J. Combin. Theory Ser. A 114 (2007), 1182–1198.
[34] M. Skandera, On the dual canonical and Kazhdan–Lusztig bases and 3412, 4231-avoiding permutations, J. Pure Appl. Algebra 212 (2008), 1086–1104.
[35] M. Skandera, Hook immanantal inequalities for totally nonnegative matrices, Electron. J. Combin. 33(4) (2026), P4.23, doi:10.37236/14684; arXiv:2510.00327v2.
[36] J. Stembridge, Immanants of totally positive matrices are nonnegative, Bull. London Math. Soc. 23 (1991), 422–428.
[37] J. Stembridge, Some conjectures for immanants, Canad. J. Math. 44 (1992), 1079–1099.

Email address: `keith.mcnulty@gmail.com`