WORKED SOLUTION

STEP I 1989, Question 15

Probability and Statistics • Comparing Distributions

STEP I 1989, Question 15

Read the question before you go on. It is in the STEP Support Programme's question database:

step.maths.org/questions/89-s1-q15

The key idea

At $12\,\mathrm{km\,h^{-1}}$ every kilometre takes exactly 5 minutes, so each journey time is a fixed number of minutes plus a simple multiple of one random variable: uniform, Poisson or binomial. Every question about time then becomes a question about that one variable. The middle part is Bayes' theorem with three equally likely routes.

Solution

Journey times

Let T_A, T_B, T_D be the journey times in minutes by the three routes. Since $12\,\mathrm{km\,h^{-1}}$ is 1 km every 5 minutes:

- Angle Avenue: $T_A = 25 + A$, with $A \sim \mathrm{U}(0, 10)$.
- Bend Boulevard: $T_B = 20 + 3B$, with $B \sim \mathrm{Po}(4)$.
- Detour Drive: let X be the number of detours. Assuming the five road works act independently, each with probability $\frac{4}{5}$, we have $X \sim \mathrm{B}(5, \frac{4}{5})$, the distance is $2 + X$ km and $T_D = 10 + 5X$.

At least 27 minutes

Angle Avenue. $T_A \geq 27 \iff A \geq 2$, so

$$\mathrm{P}(T_A \geq 27) = \frac{10 - 2}{10} = \frac{4}{5}.$$

Bend Boulevard. $T_B \geq 27 \iff 3B \geq 7 \iff B \geq 3$, since B is a whole number. So

$$\mathrm{P}(T_B \geq 27) = 1 - \mathrm{e}^{-4}\left(1 + 4 + \frac{4^2}{2!}\right) = 1 - 13\mathrm{e}^{-4} \approx 0.7619.$$

Detour Drive. $T_D \geq 27 \iff 5X \geq 17 \iff X \geq 4$. So

$$\mathrm{P}(T_D \geq 27) = 5\left(\frac{4}{5}\right)^4 \frac{1}{5} + \left(\frac{4}{5}\right)^5 = \left(\frac{4}{5}\right)^4 \left(1 + \frac{4}{5}\right) = \frac{256}{625} \cdot \frac{9}{5} = \frac{2304}{3125} = 0.73728.$$

Which route, given a long journey?

Let R be the route chosen, with $\mathrm{P}(R = A) = \mathrm{P}(R = B) = \mathrm{P}(R = D) = \frac{1}{3}$, and let L be the event that the journey lasts at least 27 minutes. By Bayes' theorem the factors of $\frac{1}{3}$ cancel:

$$\begin{aligned}\mathrm{P}(R = B \mid L) &= \frac{\mathrm{P}(L \mid B)}{\mathrm{P}(L \mid A) + \mathrm{P}(L \mid B) + \mathrm{P}(L \mid D)} = \frac{1 - 13\mathrm{e}^{-4}}{\frac{4}{5} + 1 - 13\mathrm{e}^{-4} + \frac{2304}{3125}} \\ &= \frac{1 - 13\mathrm{e}^{-4}}{\frac{7929}{3125} - 13\mathrm{e}^{-4}} \approx \frac{0.7619}{2.2992} \approx 0.331.\end{aligned}$$

This is barely above $\frac{1}{3}$, which makes sense: the three routes are almost equally likely to give a journey of 27 minutes or more.

Part (i)

Using $\mathrm{E}(A) = 5$, $\mathrm{E}(B) = 4$ and $\mathrm{E}(X) = 5 \times \frac{4}{5} = 4$, and linearity of expectation,

$$\mathrm{E}(T_A) = 25 + 5 = 30, \qquad \mathrm{E}(T_B) = 20 + 3 \times 4 = 32, \qquad \mathrm{E}(T_D) = 10 + 5 \times 4 = 30.$$

Ask yourself Two routes tie. Is that a mistake?

No. Angle Avenue and Detour Drive both average exactly 30 minutes, and Bend Boulevard averages 32. So to make the average as small as possible I should use *either* Angle Avenue or Detour Drive: they are equally good, and both beat Bend Boulevard.

Part (ii)

Now compare $\mathrm{P}(T < 32)$ for each route.

- $T_A < 32 \iff A < 7$, so $\mathrm{P}(T_A < 32) = \frac{7}{10} = 0.7$.
- $T_B < 32 \iff 3B < 12 \iff B \leq 3$, so

$$\mathrm{P}(T_B < 32) = \mathrm{e}^{-4}\left(1 + 4 + 8 + \frac{32}{3}\right) = \frac{71}{3}\mathrm{e}^{-4} \approx 0.4335.$$

- $T_D < 32 \iff 5X < 22 \iff X \leq 4$, so

$$\mathrm{P}(T_D < 32) = 1 - \left(\frac{4}{5}\right)^5 = 1 - \frac{1024}{3125} = \frac{2101}{3125} \approx 0.6723.$$

Since $0.7 > 0.6723 > 0.4335$, I should always use **Angle Avenue**, which gets me there in under 32 minutes on 70% of days.

Checks

- A simulation of 400 000 journeys gave $\mathrm{P}(T \geq 27) \approx 0.801,\ 0.761,\ 0.740$, mean times $\approx 30.0,\ 32.0,\ 30.0$ minutes, $\mathrm{P}(T < 32) \approx 0.700,\ 0.435,\ 0.670$, and $\mathrm{P}(R = B \mid L) \approx 0.331$. All agree with the exact values.
- Boundary sense-check: on Detour Drive with 3 detours the time is 25 minutes and with 4 it is 30, so "at least 27" really does mean "at least 4 detours".

Common slips

- Treating the integer conditions loosely: $3B \geq 7$ means $B \geq 3$, not $B \geq 2$, and $3B < 12$ means $B \leq 3$, not $B \leq 4$.
- Forgetting the 2 km base distance on Detour Drive, or the five-minute-per-km riding time on the other routes.
- Claiming a single winner in (i). The two means are exactly equal.

Beyond STEP

Parts (i) and (ii) show that "best" depends on what we are optimising: the mean is just one summary of a distribution, and two routes with the same mean can have very different spreads and tail probabilities. In decision theory this is the choice of a *loss function*. Grimmett and Welsh, *Probability: An Introduction*, covers the distributions here and Bayes' theorem. Blitzstein and Hwang, *Introduction to Probability*, has many problems of this kind.

Beyond STEP worked solutions: https://beyondstep.keithmcnulty.org/. Solution text CC0 1.0; the question itself is © Cambridge University Press & Assessment and is not reproduced here.