4.5 The Sophomore's Dream

WORKSHEET

The question

Let m be a positive integer and let n be a non-negative integer.

(i) Use the result $\lim_{t\to\infty} e^{-mt}t^n = 0$ to show that

$$\lim_{x\to 0} x^m(\ln x)^n = 0.$$

By writing x^x as $e^{x\ln x}$ show that

$$\lim_{x\to 0} x^x = 1.$$

(ii) Let $I_n = \int_0^1 x^m(\ln x)^n \,\mathrm{d}x$. Show that

$$I_{n+1} = -\frac{n+1}{m+1} I_n$$

and hence evaluate I_n.

(iii) Show that

$$\int_0^1 x^x \,\mathrm{d}x = 1 - \left(\tfrac{1}{2}\right)^2 + \left(\tfrac{1}{3}\right)^3 - \left(\tfrac{1}{4}\right)^4 + \cdots.$$

Hints

Try the question on your own first. If you get stuck, read the hints *one at a time*, and go back to the question after each one. Hints marked *Pause and try* are the key steps.

Ask yourself The function x^x is not defined at $x = 0$ by any formula you know (0^0 is ambiguous). How does it behave as x approaches 0 from above? Try $x = 0.1, 0.01, 0.001$ on a calculator.

STAGE 1 Powers beat logarithms

Ask yourself As $x \to 0^+$, $\ln x \to -\infty$. What substitution turns $x \to 0^+$ into $t \to \infty$, and turns $x^m(\ln x)^n$ into something like $e^{-mt}t^n$?

STAGE 2 A reduction formula

Ask yourself Integrate $I_{n+1} = \int_0^1 x^m(\ln x)^{n+1}\,dx$ by parts. Which factor should you integrate? What happens to the boundary term at $x = 0$?

STAGE 3 The series

Ask yourself Expand $x^x = e^{x\ln x}$ using the series for e^y. What does each term look like? Which value of m makes the integrals in part (ii) appear?

Pause and try Integrate term by term from 0 to 1, using part (ii) with $m = n$ (and treating $n = 0$ separately). What do you get?

Try this next

1 **The companion.** Show that $\displaystyle\int_0^1 x^{-x}\,dx = \sum_{n=1}^{\infty}\frac{1}{n^n} \approx 1.2913$.

2 **The minimum.** Show that x^x has its minimum on $(0,\infty)$ at $x = \frac{1}{e}$, with value $e^{-1/e}$.

3 **How many terms?** For an alternating series with decreasing terms, the error after stopping is less than the first omitted term. How many terms of the series for $\int_0^1 x^x\,dx$ are needed for six decimal places?

4 **A cousin.** Use the same method to show that $\displaystyle\int_0^1 x^2\ln x\,dx = -\frac{1}{9}$, and that

$\displaystyle\int_0^1 \frac{\ln x}{1-x}\,\mathrm{d}x = -\sum_{n=1}^{\infty}\frac{1}{n^2}$. (By Section 2.4, this is $-\frac{\pi^2}{6}$.)

Further reading

- **Jonathan Borwein, David Bailey & Roland Girgensohn, *Experimentation in Mathematics*** (A. K. Peters). The source of the name, and many more identities discovered by computer.
- **William Dunham, *The Calculus Gallery*** (Princeton University Press). Masterpieces of analysis from Newton to Lebesgue, including Bernoulli's work.
- **Mary Hart, *Guide to Analysis*** (Palgrave). Clear treatment of uniform convergence and when sums and integrals can be swapped.

The full guided solution, and what the question leads to at university, is in Section 4.5:
https://keithmcnulty.github.io/step_book_mcnulty/ch04/sophomores-dream.html