CHAPTER

9

Counting and Chance

Counting is the oldest mathematics there is. Doing it cleverly is still one of the hardest.

IN THIS CHAPTER

9.1	Lines Cutting the Plane	STEP II 1990
9.2	Ramsey's Theorem in Miniature	STEP I 1989
9.3	Colouring Fence Posts	STEP II 1995
9.4	Sequences as Functions	STEP I 2016
9.5	Making Change	STEP I 1997
9.6	Strange Dice	STEP I 1993
9.7	Equal Sums of Powers	STEP I 1992
9.8	Nobody Gets Their Own Shirt	STEP II 2016
9.9	The Birthday Problem	STEP II 2007
9.10	Always Able to Give Change	STEP I 2011
9.11	Collecting the Set	STEP II 1994
9.12	Random Subsets	STEP III 2019
9.13	Loaded Dice	STEP I 1999
A	Answers to the exercises	p. 89

The Art of Counting

Combinatorics is the slums of topology.

— J. H. C. Whitehead, in the 1950s. It is not a view anyone holds today.

Everyone can count. But ask how many ways there are to make £1 from coins, or to seat ten people so that nobody sits in their own place, or to colour a map with four colours, and simple counting quickly breaks down. There are too many cases to list, and the cases overlap. *Combinatorics*, the mathematics of counting and arrangement, is the art of organising such problems so that they can be solved without listing.

For a long time, combinatorics was regarded as a collection of clever tricks rather than a serious subject, as Whitehead's remark suggests. That changed in the second half of the twentieth century, driven partly by computer science, where counting the steps an algorithm takes is a combinatorial problem, and partly by the discovery of deep connections with algebra, geometry and probability. Today it is one of the most active areas of mathematics. In 2022 June Huh was awarded a Fields Medal, the highest honour in mathematics, partly for proving a 1968 conjecture about the chromatic polynomials of Section 9.3, using ideas from algebraic geometry.

Why counting matters

Probability, in its simplest form, is counting: when all outcomes are equally likely, the probability of an event is the number of favourable outcomes divided by the total. So every question about cards, dice and lotteries is a counting problem, and the second half of this chapter is drawn from the probability papers for exactly this reason. But counting also matters well beyond games. Computer scientists count the operations an algorithm needs, and the ways it might fail. Chemists count the possible structures of molecules. Biologists count the ways a strand of DNA can fold. Cryptographers count the keys an attacker must try, and, as Section 9.9 shows, how many are needed before two of something coincide.

What you will meet at university

Most universities offer a first- or second-year course in combinatorics or discrete mathematics, and its themes are the themes of this chapter. You will learn the basic counting principles (the product rule, bijections, inclusion–exclusion, the pigeonhole principle), and how to set up and solve recurrence relations. You will meet *generating functions*, which turn sequences into functions and counting into algebra. You will study graphs, from colourings to Ramsey theory, and partially ordered sets such as the lattice of subsets. And probability courses will use the same tools, together with linearity of expectation, to analyse random structures. Further on, combinatorics

connects with algebra (symmetric functions, representation theory), with number theory (partitions, additive combinatorics) and with computer science (the analysis of algorithms, complexity theory).

The questions that run through this chapter

Three big questions

1. **How do you count without listing?** What structures (recurrences, bijections, generating functions) let you count huge collections exactly?
2. **When must something happen?** When does size alone force a pattern, a coincidence or a match?
3. **How do counting and probability help each other?** When is a probability best found by counting, and when is a count best found by probability?

The chapter has two halves. The first seven sections come from the pure papers. They begin with recurrences (lines cutting the plane, colouring fence posts) and the pigeonhole principle (Ramsey's theorem), then develop generating functions and use them to count coins, design strange dice and split numbers into teams with equal power sums. The last six sections are classic counting problems from the probability papers: derangements and the appearance of $1/e$, the birthday problem, the ballot problem and its reflection principle, the coupon collector, random subsets, and the impossibility of loading dice so that every total is equally likely.

Ideas in this chapter

§	Topic	Idea
9.1	Regions of the plane	What changes when you add one more
9.2	Ramsey theory	The pigeonhole principle
9.3	Graph colouring	Coupled recurrences; chromatic polynomials
9.4	Generating functions	A sequence becomes a function
9.5	Partitions	Counting via coefficients
9.6	Sicherman dice	Factorising polynomials
9.7	Thue–Morse	Repeated roots give balanced sums
9.8	Derangements	Inclusion–exclusion; $1/e$
9.9	Coincidences	Complements; the square-root law
9.10	The ballot problem	The reflection principle
9.11	Coupon collector	Linearity of expectation; $n \ln n$
9.12	Random subsets	One element at a time
9.13	Loaded dice	Impossibility by AM–GM

How to read this chapter

Always start with small cases. Draw three lines, list the six orders of three cards, count the ways to make 5p. Small cases give you data to guess from, and they check your formulas at the end. But do not trust a pattern until you have a reason for it: Section 9.1 includes a famous sequence, $1, 2, 4, 8, 16, \ldots$, whose next term is 31.

9.1 Lines Cutting the Plane

The question

STEP II 1990, Question 4

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/90-s2-q4`

The hook

How many pieces can you cut a pizza into with n straight cuts, if the pieces need not be the same size? One cut gives 2, two cuts give 4, three cuts give 7 (not 8: try it), four give 11. The sequence $1, 2, 4, 7, 11, 16, \ldots$ is known as the *lazy caterer's sequence*, and this question asks for its formula.

The question is a first lesson in a technique that runs through all of combinatorics: do not try to count everything at once. Instead, ask what changes when you add one more object. And the answer turns out to be a special case of Euler's formula $V - E + F = 2$, which you met with the Platonic solids in Section 7.9.

Before you start

Pause and try Draw the cases $n = 1, 2, 3, 4$ carefully, keeping the lines in general position (no two parallel, no three through a point). Count the segments and the regions in each case. Can you guess the patterns?

You should find:

n	0	1	2	3	4
segments	0	1	4	9	16
regions	1	2	4	7	11

The segments look like n^2. The regions go up by $1, 2, 3, 4$: the differences are the natural numbers.

Working it through

STAGE 1 Counting segments

Ask yourself Concentrate on one line. How many of the other lines cross it, and into how many pieces do they cut it?

Each line is crossed by each of the other $n-1$ lines (none is parallel to it), and at $n-1$ different points (no three lines meet at a point). Those $n-1$ points cut the line into n segments: two infinite ones at the ends, and $n-2$ finite ones between. With n lines, there are

$$n \times n = n^2 \text{ segments.} \qquad \blacksquare$$

This is a direct count, with no induction needed: we counted segments line by line.

STAGE 2 What happens when we add a line?

Ask yourself Suppose $n-1$ lines are drawn, and you add the nth. It is cut into n segments by the others. What does each segment do to the region it passes through?

As the new line is drawn, each of its n segments runs through one of the existing regions and splits it into two (Figure 9.1.1). Different segments lie in different regions, because to pass from one region to another the line must cross an existing line. So adding the nth line increases the number of regions by exactly n:

$$R(n) = R(n-1) + n.$$

STAGE 3 Solving the recurrence

Starting from $R(0) = 1$ (no lines, one region),

$$R(n) = 1 + (1 + 2 + \cdots + n) = 1 + \frac{n(n+1)}{2} = \frac{n^2 + n + 2}{2}.$$

To present this as an induction: the formula holds for $n = 1$, since $\frac{1}{2}(1+1+2) = 2$; and if $R(n-1) = \frac{1}{2}\big((n-1)^2 + (n-1) + 2\big) = \frac{1}{2}(n^2 - n + 2)$, then $R(n) = \frac{1}{2}(n^2 - n + 2) + n = \frac{1}{2}(n^2 + n + 2)$. $\blacksquare$




Figure 9.1.1 Three lines (teal) divide the plane into 7 regions. The fourth line (plum) is crossed by the other three at the amber points, which cut it into four segments, numbered 1 to 4. Each segment splits one region into two, so the fourth line adds 4 regions, giving 11.

Looking back

- **Count one piece at a time.** Segments were counted line by line; regions were counted by what one new line adds.
- **Turn a geometric step into a recurrence.** The observation "each new segment splits one region" became $R(n) = R(n-1) + n$.
- **General position matters.** Parallel lines, or three lines through a point, would give fewer regions. Find where each assumption was used.

What's really going on

Euler's formula in disguise

Here is a second proof, which explains the answer rather than just deriving it. Think of the arrangement of lines as a map: the crossing points are *vertices*, the segments are *edges* and the regions are *faces*. There are $V = \binom{n}{2}$ vertices (one for each pair of lines) and $E = n^2$ edges. The infinite segments have only one end, so add a single extra vertex "at infinity" where they all meet; now the picture is a map drawn on a sphere (imagine the plane wrapped round a ball, with infinity at the top). Euler's formula $V - E + F = 2$ from Section 7.9 gives

$$\left(\binom{n}{2} + 1\right) - n^2 + F = 2, \qquad \text{so} \qquad F = n^2 - \frac{n(n-1)}{2} + 1 = \frac{n^2 + n + 2}{2}.$$

The same number, from a completely different direction. The regions also split neatly into types: $2n$ of them are unbounded (one between each pair of neighbouring infinite rays), and the remaining $\frac{1}{2}(n^2 + n + 2) - 2n = \binom{n-1}{2}$ are bounded.

Binomial coefficients and higher dimensions

The formula can be written as

$$R(n) = \binom{n}{0} + \binom{n}{1} + \binom{n}{2},$$

and this is not a coincidence. Turn the picture so that no line is horizontal. A region that is bounded below has a lowest point, which must be a crossing, and each crossing is the lowest point of exactly one region (the one directly above it): that gives $\binom{n}{2}$ regions, one for each pair of lines. Every other region stretches down for ever, and so meets a horizontal line drawn far below all the crossings; the n lines cut that horizontal line into $n+1$ intervals, one in each such region. In total, $\binom{n}{2} + n + 1$. In three dimensions, n planes in general position cut space into

$$\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \binom{n}{3} = \frac{n^3 + 5n + 6}{6}$$

pieces (the *cake numbers* $1, 2, 4, 8, 15, 26, \ldots$), and the pattern continues in every dimension. The proof is the same induction: a new plane is cut by the others into regions arranged like lines in a plane, and each of those regions splits a piece of space in two.

Steiner and the arrangement of hyperplanes

The formula for regions cut by lines and planes was found by Jakob Steiner in 1826. Today it is the starting point of the theory of *hyperplane arrangements*, which counts the regions formed by arbitrary collections of lines, planes and their higher-dimensional analogues, including non-general ones. A remarkable 1975 theorem of Thomas Zaslavsky counts the regions of any arrangement using a polynomial that encodes which hyperplanes meet. The same ideas appear in machine learning: a neural network built from "ReLU" units cuts its input space into regions by hyperplanes, and the number of regions measures how complicated a function it can represent.

The thinking skill

Ask what changes when you add one more

To count a configuration of n objects, compare it with the configuration of $n-1$. The change is often much easier to count than the whole.

This is induction used as a tool for discovery, not just for proof. It is one of the most common moves in combinatorics, and it turns a geometric question into a recurrence relation that can be solved.

Try this next

1. **Circles.** Show that n circles, each pair meeting in two points and no three through a point, divide the plane into $n^2 - n + 2$ regions.

2. **Parallel lines.** If the n lines include k that are all parallel to each other (and otherwise general position), how many regions are there?

3. **Cake numbers.** Prove by induction that n planes in general position divide space into $\frac{1}{6}(n^3 + 5n + 6)$ regions.

4. **A pattern that breaks.** Put n points on a circle and join every pair with a chord, with no three chords through one point inside the circle. For $n = 1, 2, 3, 4, 5$ you get $1, 2, 4, 8, 16$ regions. Count the regions for $n = 6$. (*Hint:* use Euler's formula; each set of four points gives one crossing.) Show that the number is $1 + \binom{n}{2} + \binom{n}{4}$.

Further reading

- **Ronald Graham, Donald Knuth & Oren Patashnik, *Concrete Mathematics*** (Addison-Wesley), Chapter 1. Starts with exactly this problem, and the "pizza" and "zig" variants.
- **Miklós Bóna, *A Walk Through Combinatorics*** (World Scientific). A friendly first course, with chapters on induction, graphs and Euler's formula.
- **Richard Guy, "The strong law of small numbers"**, *American Mathematical Monthly* 95 (1988). Many patterns that look obvious and turn out to be false, including the circle problem in the exercises.

9.2 Ramsey's Theorem in Miniature

The question

STEP I 1989, Question 4

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/89-s1-q4`

The hook

At any party of six people, there are either three who all know each other or three who are all strangers to each other. This is guaranteed, whoever the six people are. With five people it is not guaranteed. This little fact is the first case of *Ramsey's theorem*, and it captures a principle that runs through modern combinatorics: *complete disorder is impossible*. Any sufficiently large structure, however it is coloured or arranged, must contain a highly ordered piece.

The proof takes four lines. But the obvious next question (how many people guarantee four mutual friends or four mutual strangers? Five?) leads to some of the hardest open problems in mathematics.

Before you start

Ask yourself The six points matter only through which pairs are joined by which colour: the geometry is a distraction. How many segments are there, and how many leave each point?

There are $\binom{6}{2} = 15$ segments, and 5 leave each point. (The general-position condition just makes sure every three points form a genuine triangle. The problem is really about the *complete graph* on six vertices, K_6, with its edges coloured in two colours.)

Working it through

STAGE 1 **Five segments, two colours**

Ask yourself Five segments leave A, each gold or silver. What can you say for certain about their colours?

At least three of them have the same colour. If at most two were gold and at most two silver, there would be at most four segments. This is the *pigeonhole principle*: five objects in two boxes put at least three in one box. Say (swapping the names of the colours if necessary) that AB, AC and AD are all gold.

STAGE 2 Look at the far ends


Ask yourself Now look at the triangle BCD. What happens if any one of its sides is gold? What if none of them is?

If any side of triangle BCD is gold, say BC, then ABC has all three sides gold: AB, AC and BC. If no side of BCD is gold, then BC, CD and DB are all silver, and BCD is a silver triangle. Either way there is a one-colour triangle (Figure 9.2.1, right). ■


STAGE 3 Five points are not enough

Ask yourself With five points, each point has only four segments. How might you colour them so that each point has exactly two of each colour?

Place the five points as a regular pentagon (lifting them slightly off a plane if you want general position). Colour the five sides of the pentagon gold and the five diagonals silver (Figure 9.2.1, left). The gold edges form a five-cycle, which contains no triangle; the silver edges form a pentagram, which is also a five-cycle and contains no triangle. So there is no one-colour triangle. ■



five points: no one-colour triangle



six points: look at BCD

Figure 9.2.1 Left: five points with the pentagon gold (solid) and the pentagram silver (dashed) have no one-colour triangle. Right: with six points, three of the five segments from A share a colour, say gold to B, C, D. The triangle BCD (plum) then decides: one gold side makes a gold triangle with A, and no gold side makes BCD a silver triangle.

Looking back

- **Strip away the irrelevant.** The points in space became a coloured graph; only the colours of the edges mattered.
- **Pigeonhole.** Five edges and two colours forced three of one colour.
- **Case analysis on the right object.** Once three gold edges were found, the triangle at their far ends settled everything.
- **Sharpness.** The pentagon shows the number six cannot be lowered. A good theorem comes with an example showing it is best possible.

What's really going on

Ramsey numbers

The *Ramsey number* $R(s,t)$ is the smallest N such that every colouring of the edges of K_N in gold and silver contains either a gold K_s (a set of s points with all edges between them gold) or a silver K_t. The question has shown that $R(3,3) = 6$: six points always work, and the pentagon shows five do not.

Frank Ramsey proved in 1928 that these numbers always exist: for every s and t there is some N that works. The proof is an induction that generalises the argument above, and shows $R(s,t) \le R(s-1,t) + R(s,t-1)$. But exact values are extraordinarily hard to find:

$$R(3,3) = 6, \qquad R(4,4) = 18, \qquad 43 \le R(5,5) \le 46.$$

The value of $R(4,4)$ was found in 1955. Despite enormous computer searches, $R(5,5)$ is still unknown; the upper bound of 46 was proved only in 2024. A computer cannot simply check every colouring of K_{43}: there are 2^{903} of them, far more than the number of atoms in the universe.

Erdős and the aliens

Paul Erdős, who did more than anyone to develop Ramsey theory, liked to tell this story. Suppose aliens invade the Earth and threaten to destroy it unless we tell them the value of $R(5,5)$. We should marshal all our computers and mathematicians, and we might find it. But if they asked for $R(6,6)$, we should try to destroy the aliens. Erdős also proved, in 1947, that $R(k,k) > 2^{k/2}$, using a startling method: he showed that a *randomly* coloured graph on fewer than $2^{k/2}$ points has a positive probability of containing no one-colour K_k, so a good colouring must exist, even though nobody knows how to construct one. This *probabilistic method* is now one of the most powerful tools in combinatorics.

Order from chaos

Ramsey's theorem has many relatives, all expressing the same idea: large enough structures cannot avoid order. *Van der Waerden's theorem* says that if the whole numbers are

coloured with finitely many colours, there are arbitrarily long arithmetic progressions of one colour. The *Erdős–Szekeres theorem* says that any sequence of $n^2 + 1$ distinct numbers contains an increasing or a decreasing subsequence of length $n + 1$. And Ramsey himself was interested in a question of logic: he proved his theorem as a lemma in a paper on deciding the truth of logical statements.

The thinking skill

The pigeonhole principle

If more than kn objects are placed in n boxes, some box contains more than k objects.

The principle is obvious, but using it well is an art: the skill is in choosing what the objects and boxes should be. Here they were the five edges at one point and the two colours. At university, pigeonhole arguments appear in number theory (approximating irrationals by fractions), in analysis (the Bolzano–Weierstrass theorem) and throughout combinatorics.

Try this next

1 **At least two.** Show that in any two-colouring of K_6 there are in fact at least *two* one-colour triangles. (*Hint:* count the "bichromatic angles": pairs of edges of different colours at a common vertex. A point with g gold and $5 - g$ silver edges is at the apex of $g(5 - g) \leq 6$ such angles. Each triangle that is not one-colour contains exactly two.)

2 **Three colours.** If the edges of K_{17} are coloured in three colours, show that there is a one-colour triangle. (*Hint:* one point has at least six edges of one colour.)

3 **Three or four.** Show that $R(3,4) \leq 10$: in any gold/silver colouring of K_{10} there is a gold triangle or a silver K_4. (*Hint:* a point has either at least 4 gold edges or at least 6 silver edges.) In fact $R(3,4) = 9$.

4 **Monotone subsequences.** Prove the Erdős–Szekeres theorem for $n = 2$: any sequence of 5 distinct numbers contains an increasing or decreasing subsequence of length 3. Find a sequence of 4 with neither.

Further reading

- **Miklós Bóna, *A Walk Through Combinatorics*** (World Scientific), the chapters on the pigeonhole principle and Ramsey theory.
- **Ronald Graham & Joel Spencer, "Ramsey theory"**, *Scientific American* (July 1990). A popular introduction by two leading researchers.

- **Noga Alon & Joel Spencer, *The Probabilistic Method*** (Wiley). The standard book on Erdős's method; the first chapter is accessible.
- **Paul Hoffman, *The Man Who Loved Only Numbers*** (Fourth Estate). A biography of Erdős.

9.3 Colouring Fence Posts

The question

STEP II 1995, Question 2

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/95-s2-q2`

The hook

Painting posts in a line is easy: 3 choices for the first, then 2 for each of the rest (anything but the colour of the previous post), giving $3 \times 2^{n-1}$. Bend the line into a circle and it becomes surprisingly awkward, because the last post now has to differ from the first as well as from its neighbour, and whether that rules out one colour or none depends on everything that came before.

The question gets round this with a classic trick: split the count into cases according to the colour of the last post, and find how the cases feed into each other. The answer, $2^n + 2(-1)^n$, is the first example of a *chromatic polynomial*, part of the mathematics behind the four colour theorem.

Before you start

Pause and try Count the ways to paint 3 posts in a circle, and 4. Does $3 \times 2^{n-1}$ overcount, and by how much?

Three posts in a circle are all adjacent to each other, so they need three different colours: $3! = 6$ ways. Four posts in a circle: 18 ways (you can check this at the end). Compare $3 \times 2^2 = 12$ and $3 \times 2^3 = 24$ for the line: the circle loses the colourings in which the last post matches the first.

Working it through

STAGE 1 Two kinds of colouring

Ask yourself Take a colouring of $n+1$ posts with the first and last red. Remove the last post. What can you say about the colouring of the first n posts that remains?

Post n is next to post $n+1$, which is red, so post n is not red. Removing post $n+1$ leaves a colouring of n posts with the first red and the last not red: one of the s_n colourings. Conversely, any of the s_n colourings can be extended in exactly one way, by painting post $n+1$ red. So the two sets correspond exactly, and $r_{n+1} = s_n$.

Ask yourself What is $r_n + s_n$?

Every colouring with the first post red has its last post either red or not, so $r_n + s_n$ counts all colourings of n posts with the first red. After the first post, each post has 2 choices, so $r_n + s_n = 2^{n-1}$. Hence

$$r_{n+1} + r_n = s_n + r_n = 2^{n-1}. \qquad \blacksquare$$

STAGE 2 The formula by induction

For $n = 1$ there is one post, painted red, so $r_1 = 1$, and the formula gives $\frac{1}{3}(1+2) = 1$. Suppose the formula holds for r_n. Then

$$r_{n+1} = 2^{n-1} - r_n = 2^{n-1} - \frac{2^{n-1} + 2(-1)^{n-1}}{3} = \frac{2 \cdot 2^{n-1} - 2(-1)^{n-1}}{3} = \frac{2^n + 2(-1)^n}{3},$$

which is the formula for r_{n+1}. $\blacksquare$

Pause and try Where did the formula come from? The recurrence $r_{n+1} + r_n = 2^{n-1}$ is like a differential equation: a "particular solution" $c \cdot 2^n$ plus a "complementary function" $A(-1)^n$ that solves $r_{n+1} + r_n = 0$. Find c and A.

STAGE 3 The circle

Ask yourself In a circle of n posts, fix the colour of post 1, say red. Now cut the circle between post n and post 1, and straighten it into a line. Which of the line colourings correspond to valid circle colourings?

Straightened out, the circle becomes a line of n posts starting with red, in which the last post must also differ from post 1: the last post is not red. These are exactly the s_n colourings. Post 1 can be any of 3 colours, so the number of circle colourings is

$$3s_n = 3(2^{n-1} - r_n) = 3 \cdot 2^{n-1} - 2^{n-1} - 2(-1)^{n-1} = 2^n + 2(-1)^n. \qquad \blacksquare$$

Check: $n = 3$ gives $8 - 2 = 6$, and $n = 4$ gives $16 + 2 = 18$.




Figure 9.3.1 Left: in a line, each post after the first just has to differ from its predecessor. Right: in a circle, the last post must also differ from post 1. Cutting the circle at the dashed line turns a circle colouring into a line colouring whose first and last posts differ: one of the s_n colourings counted in the question.

Looking back

- **Split into cases that feed each other.** Recording whether the last post was red gave two sequences, r_n and s_n, linked by simple recurrences.
- **Bijections.** $r_{n+1} = s_n$ was proved by matching colourings one-to-one, not by calculation.
- **Cut the circle.** A cyclic problem became a linear one with a condition on the ends.

What's really going on

Chromatic polynomials

The posts and their adjacencies form a *graph*: a path for the line, a cycle for the circle. A *proper colouring* gives adjacent vertices different colours. The number of proper colourings with k colours is a polynomial in k, the *chromatic polynomial* of the graph, introduced by George Birkhoff in 1912. Repeating the question's argument with k colours gives

$$\text{path: } k(k-1)^{n-1}, \qquad \text{cycle: } (k-1)^n + (-1)^n(k-1).$$

With $k = 3$ the cycle formula is $2^n + 2(-1)^n$. Birkhoff hoped to prove the four colour theorem (every map can be coloured with four colours so that neighbouring countries differ) by showing that the chromatic polynomial of every map's graph is positive at $k = 4$. That approach never succeeded; the theorem was finally proved in 1976 by Kenneth Appel and Wolfgang Haken, with extensive computer help. Chromatic polynomials remain an active field, with surprising connections to statistical physics, where they appear as the *Potts model* of magnetism.

The same count by matrices

There is another way to see the answer, using linear algebra from Chapter 7. Let $\mathbf{M}$ be the 3×3 matrix with 0 on the diagonal and 1 elsewhere: entry (i, j) is 1 when colour j may follow colour i. Then entry (i, j) of $\mathbf{M}^{n-1}$ counts line colourings that start with colour i and end with colour j, so r_n is a diagonal entry of $\mathbf{M}^{n-1}$. A circle colouring is a sequence that returns to its start after n steps, so the number of circle colourings is the *trace* of $\mathbf{M}^n$. Since $\mathbf{M} = \mathbf{J} - \mathbf{I}$, where every entry of $\mathbf{J}$ is 1, the eigenvalues of $\mathbf{M}$ are $3 - 1 = 2$ (eigenvector $(1, 1, 1)$) and -1 (twice), and the trace of $\mathbf{M}^n$ is the sum of the nth powers of the eigenvalues:

$$\operatorname{tr} \mathbf{M}^n = 2^n + (-1)^n + (-1)^n = 2^n + 2(-1)^n.$$

This *transfer matrix* method is used throughout combinatorics and statistical physics to count arrangements built up one step at a time.

The thinking skill

Set up coupled recurrences

When one sequence will not satisfy a simple recurrence on its own, record a little extra information (here, the colour of the last post) and find recurrences linking the cases.

This is the idea of a *state*: keep track of exactly what you need to know to continue, and nothing more. It is the basis of dynamic programming in computer science, of Markov chains in probability (Section 12.16), and of the transfer matrix method.

Try this next

1. **k colours.** Repeat the question with k colours. Show that $r_{n+1} = s_n$ still holds and that $r_{n+1} + r_n = (k-1)^{n-1}$, find r_n, and deduce the chromatic polynomial of the cycle, $(k-1)^n + (-1)^n(k-1)$.

2. **No two reds together.** How many ways are there to paint n posts in a line red or white, with no two adjacent posts red? (You should find the Fibonacci numbers.)

3. **A wheel.** Add a central post adjacent to every post of the circle (a "wheel"). How many ways are there to colour it with 4 colours? (*Hint:* the centre uses one colour, and the circle must use the other three.)

4. **Necklaces.** If two colourings of the circle count as the same when one is a rotation of the other, how many different colourings are there for $n = 4$? For $n = 5$? (This is the start of *Burnside's lemma*.)

Further reading

- **Robin Wilson, *Introduction to Graph Theory*** (Pearson). The standard first course; covers colourings and the four colour problem.
- **Robin Wilson, *Four Colours Suffice*** (Penguin). The story of the four colour theorem, for a general reader.
- **Herbert Wilf, *generatingfunctionology*** (free online). Recurrences and how to solve them, including the transfer matrix method.

9.4 Sequences as Functions

The question

STEP I 2016, Question 8

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/16-s1-q8`

The hook

Work out $1/89$ on a calculator: $0.011235955\ldots$. The digits begin $0,1,1,2,3,5$, the Fibonacci numbers. Is this a coincidence? It is not, and this question explains why.

The idea is one of the most powerful in combinatorics. Take an infinite sequence of numbers and package it as a single function, by using the terms as the coefficients of a power series. Then facts about the sequence become facts about the function, and you can use algebra and calculus on them. Herbert Wilf put it memorably: "A generating function is a clothesline on which we hang up a sequence of numbers for display."

Before you start

? **Ask yourself** What sequence has generating function $\dfrac{1}{1-x}$? What about $\dfrac{1}{1-2x}$? What happens to the sequence if you multiply its generating function by x?

The geometric series $\frac{1}{1-x} = 1 + x + x^2 + \cdots$ is the generating function of $1,1,1,\ldots$, and $\frac{1}{1-2x} = 1 + 2x + 4x^2 + \cdots$ that of the powers of 2. Multiplying by x shifts every term one place along: $x \cdot \frac{1}{1-x} = x + x^2 + \cdots$ belongs to $0,1,1,1,\ldots$.

Working it through

STAGE 1 **The natural numbers and the triangle numbers**

Ask yourself Expand $(1-x)^{-2}$ by the binomial series. What is the coefficient of x^n?

The binomial series gives

$$(1-x)^{-2} = 1 + 2x + 3x^2 + 4x^3 + \cdots = \sum_{n\geq 0} (n+1)x^n,$$

so $x(1-x)^{-2} = x + 2x^2 + 3x^3 + \cdots = \sum nx^n$: the generating function of $u_n = n$.

Similarly, the coefficient of x^n in $(1-x)^{-3}$ is $\frac{(-3)(-4)\cdots(-3-n+1)}{n!}(-1)^n = \frac{(n+1)(n+2)}{2}$, so

$$x(1-x)^{-3} = \sum_{n\geq 1} \frac{n(n+1)}{2}\, x^n = x + 3x^2 + 6x^3 + 10x^4 + \cdots,$$

the generating function of the triangle numbers $u_n = \frac{1}{2}n(n+1)$.

Ask yourself How can you write n^2 in terms of n and $\frac{1}{2}n(n+1)$?

Since $n^2 = 2 \cdot \frac{1}{2}n(n+1) - n$, and generating functions add just as sequences do,

$$\sum n^2 x^n = \frac{2x}{(1-x)^3} - \frac{x}{(1-x)^2} = \frac{2x - x(1-x)}{(1-x)^3} = \frac{x(1+x)}{(1-x)^3}. \qquad \blacksquare$$

STAGE 2 A recurrence becomes an equation

Ask yourself Multiply $u_n = ku_{n-1}$ by x^n and add up over all $n \geq 1$. What is each side, in terms of f?

The left side is $\sum_{n\geq 1} u_n x^n = \mathrm{f}(x) - u_0 = \mathrm{f}(x) - a$. The right side is $kx \sum_{n\geq 1} u_{n-1}x^{n-1} = kx\,\mathrm{f}(x)$. So $\mathrm{f}(x) = a + kx\,\mathrm{f}(x)$, and

$$\mathrm{f}(x) = \frac{a}{1-kx}. \qquad \blacksquare$$

Of course, $u_n = ak^n$; expanding $\frac{a}{1-kx}$ as a geometric series confirms it.

STAGE 3 The Fibonacci numbers

Do the same with $u_n = u_{n-1} + u_{n-2}$, for $n \geq 2$:

$$\sum_{n\geq 2} u_n x^n = x \sum_{n\geq 2} u_{n-1}x^{n-1} + x^2 \sum_{n\geq 2} u_{n-2}x^{n-2},$$

that is,

$$\mathrm{f}(x) - u_0 - u_1 x = x\big(\mathrm{f}(x) - u_0\big) + x^2\,\mathrm{f}(x).$$

With $u_0 = 0$ and $u_1 = 1$, this says $\mathrm{f}(x) - x = x\,\mathrm{f}(x) + x^2\,\mathrm{f}(x)$, so

$$\mathrm{f}(x) = \frac{x}{1-x-x^2}. \qquad \blacksquare$$

STAGE 4 Explaining $1/89$

Pause and try Ignoring convergence (which is fine here: the series converges for $|x| < 0.618$), put $x = \frac{1}{10}$ in f. What do you get?

$$\frac{1}{10}F_1 + \frac{1}{100}F_2 + \frac{1}{1000}F_3 + \cdots = \mathrm{f}\Big(\frac{1}{10}\Big) = \frac{1/10}{1 - 1/10 - 1/100} = \frac{10}{89}.$$

Dividing by 10, $\frac{1}{89} = \sum F_n/10^{n+1}$. The Fibonacci numbers are placed in successive decimal places, overlapping when they get bigger than 9 (Figure 9.4.1). The early digits of $1/89$ show the Fibonacci numbers; later ones are blurred by carrying.

```
0.  1
0.    1
0.      2
0.        3
0.          5
0.            8
0.            1 3
0.              2 1
0.                3 4
0.                  5 5
 ⋮
0. 0 1 1 2 3 5 9 5 5 0 5
```

Figure 9.4.1 Adding $F_n/10^{n+1}$ for $n = 1, 2, 3, \ldots$. Each Fibonacci number ends one decimal place further to the right. Once they have two digits they overlap, and carrying changes the digits from the seventh place on: $\frac{1}{89} = 0.01123595505\ldots$.

Looking back

- **Turn a sequence into a function.** The whole infinite sequence became one rational function.
- **Recurrences become equations.** Multiplying by x^n and summing turned the rule $u_n = u_{n-1} + u_{n-2}$ into an equation for f that could be solved by algebra.
- **Build new from old.** Adding, shifting and differentiating generating functions produced new ones without starting from scratch.

What's really going on

From generating function to formula

Generating functions do not just encode a sequence: they can solve for it. The denominator factorises as $1 - x - x^2 = (1 - \varphi x)(1 - \psi x)$, where $\varphi = \frac{1+\sqrt{5}}{2}$ and $\psi = \frac{1-\sqrt{5}}{2}$ are the roots of $t^2 = t + 1$. Partial fractions give

$$\frac{x}{1 - x - x^2} = \frac{1}{\sqrt{5}}\left(\frac{1}{1 - \varphi x} - \frac{1}{1 - \psi x}\right),$$

and expanding each fraction as a geometric series,

$$F_n = \frac{\varphi^n - \psi^n}{\sqrt{5}}.$$

This is *Binet's formula*. Since $|\psi| < 1$, the second term is tiny, so F_n is the nearest integer to $\varphi^n / \sqrt{5}$, and the ratio of successive Fibonacci numbers tends to the golden ratio (the subject of Section 2.2). The same method solves any linear recurrence with constant coefficients; the exercises include an example from STEP I 2004.

Why generating functions are so useful

The real power appears when we multiply. The coefficient of x^n in a product $f(x)g(x)$ is $\sum_k u_k v_{n-k}$, which adds up over all ways of splitting n into two parts. That is exactly what we need to count things built from two pieces. The next two sections use it to count the ways of making change (Section 9.5) and to find strangely numbered dice with the same totals as ordinary ones (Section 9.6). In probability, the same idea gives the *probability generating function*, which turns sums of independent random variables into products.

De Moivre, Euler and Laplace

Abraham de Moivre used generating functions in the 1720s to solve linear recurrences, essentially by the method above. Leonhard Euler used them in 1748 to study partitions of whole numbers, and Pierre-Simon Laplace made them the foundation of his *Théorie analytique des probabilités* (1812). In the twentieth century they became central to combinatorics, and analytic methods (studying a generating function as a function of a complex variable) now give precise estimates for counting problems far too complicated to solve exactly.

The thinking skill

Change the representation

A problem about a sequence may become easy when the sequence is turned into a function, a matrix, or a picture.

Generating functions translate recurrences into algebra and counting into multiplication. At university you will meet many such translations: Fourier series turn functions into sequences, Laplace transforms turn differential equations into algebra, and matrices turn geometry into arithmetic.

Try this next

1 **Two digits at a time.** Try STEP II 2022, Question 3 (in OCR's download of the 2022 papers, `www.ocr.org.uk/Images/696587-2022-past-papers.zip`). It explains why the decimal expansion of $\frac{1}{89}$ begins with the Fibonacci numbers, and asks for a fraction whose decimals show the pattern two digits at a time.

2 **Solving a recurrence.** Try STEP I 2004, Question 7 (`step.maths.org/questions/04-s1-q7`). It solves a linear recurrence with a generating function and partial fractions, then runs the method in reverse.

3 **Counting numbers.** Show that $\frac{1}{9801} = 0.00\,01\,02\,03\,04\ldots$, with the two-digit blocks counting upwards. Where does the pattern first go wrong, and why?

4 **Sums of squares.** Explain why $\frac{1}{1-x}f(x)$ is the generating function of the partial sums $u_0 + u_1 + \cdots + u_n$. Use this and the answer to part (i) to find a formula for $1^2 + 2^2 + \cdots + n^2$.

Further reading

- **Herbert Wilf, *generatingfunctionology*** (A K Peters; free online). The classic introduction. Chapter 1 is perfect after this section.
- **Ronald Graham, Donald Knuth & Oren Patashnik, *Concrete Mathematics*** (Addison-Wesley), Chapter 7.
- **Philippe Flajolet & Robert Sedgewick, *Analytic Combinatorics*** (Cambridge University Press; free online). Far beyond first year, but the opening pages show where the subject leads.

9.5 Making Change

The question

STEP I 1997, Question 1

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/97-s1-q1`

The hook

This looks like the easiest question in the book: just list the possibilities. But listing gets out of hand very quickly. There are 41 ways to make 20p, 451 ways to make 50p, and 4563 ways to make £1 from the coins in a British pocket. The question is really asking for a *system*, a way to organise the count so that nothing is missed and nothing counted twice.

Behind the system is a beautiful piece of mathematics. Each coin corresponds to a geometric series, the number of ways to make n pence is a coefficient in their product, and the same idea, applied to all whole numbers at once, gives Euler's theory of *partitions*, one of the deepest topics in number theory.

Before you start

Ask yourself How many ways are there to make n pence using only 2p and 1p coins? Why does the answer depend only on how many 2p coins you use?

Once the number of 2p coins is chosen, the rest must be 1p coins, so there is exactly one way for each possible number of 2p coins: $0, 1, \ldots, \lfloor n/2 \rfloor$. That gives $\lfloor n/2 \rfloor + 1$ ways. Call this $a(n)$. For example, $a(10) = 6$ and $a(5) = 3$.

Working it through

STAGE 1 **Ten pence**

Ask yourself Organise the count by the largest coins. How many 10p coins can you use? Given that, how many 5p coins?

- One 10p coin: 1 way.
- No 10p, two 5p: 1 way.
- No 10p, one 5p: the other 5p is made from 2p and 1p, in $a(5) = 3$ ways.
- No 10p or 5p: 10p from 2p and 1p, in $a(10) = 6$ ways.

The total is $1 + 1 + 3 + 6 = 11$. ■

STAGE 2 Building up layer by layer

Ask yourself Let $b(n)$ be the number of ways to make n pence from 5p, 2p and 1p coins. How can you express $b(n)$ using the function a?

Choose the number k of 5p coins; the remaining $n - 5k$ pence are made from 2p and 1p coins. So

$$b(n) = a(n) + a(n-5) + a(n-10) + \cdots.$$

This gives

n	0	5	10	15	20
$a(n)$	1	3	6	8	11
$b(n) = a(n) + b(n-5)$	1	4	10	18	29

Similarly, let $c(n)$ count ways using 10p coins as well; then $c(n) = b(n) + b(n-10) + \cdots$, so $c(10) = 10 + 1 = 11$ (agreeing with Stage 1) and $c(20) = b(20) + b(10) + b(0) = 29 + 10 + 1 = 40$.

STAGE 3 Twenty pence

Finally the 20p coin: either we use it (one way) or we do not ($c(20)$ ways). So there are

$$40 + 1 = 41 \text{ ways.}$$ ■

Pause and try Notice the structure: each new coin adds a layer, and each layer is a sum of the previous layer at spacings of the new coin's value. Can you see how a computer would carry out this calculation for £1, or £100?

Looking back

- **Organise by the largest part.** Fixing the number of the largest coins first made the count systematic.

- **Build a table.** Each new coin added a row computed from the row before. This is *dynamic programming*, the standard way computers count such things.
- **Check against a known case.** The table reproduced the 11 ways for 10p.

What's really going on

Coins as geometric series

Here is the idea that turns this into algebra. The number of 5p coins used can be $0, 1, 2, \ldots$, contributing $0, 5, 10, \ldots$ pence. Record this as the series

$$1 + x^5 + x^{10} + x^{15} + \cdots = \frac{1}{1 - x^5}.$$

Do the same for each coin and multiply:

$$\frac{1}{(1-x)(1-x^2)(1-x^5)(1-x^{10})(1-x^{20})} = \sum_n w(n) x^n.$$

When the brackets are multiplied out, each term x^n arises from choosing one term from each bracket, that is, a number of each coin, with total value n. So $w(n)$, the coefficient of x^n, is exactly the number of ways to make n pence. The table in Stage 2 was simply multiplying out this product one bracket at a time. This is the generating function idea of Section 9.4, and multiplication is where it shows its power: products of generating functions count combinations of choices.




Figure 9.5.1 The number of ways to make n pence from 1p, 2p, 5p, 10p, 20p and 50p coins. The jumps at multiples of 5 and 10 show where a new coin first becomes usable. With six coins the count grows roughly like n^5: in general, like a polynomial of degree one less than the number of coins.

Partitions

What if every whole number is allowed as a coin? Then the number of ways to write n as a sum of positive integers, ignoring order, is the *partition number* $p(n)$. For example

$p(4) = 5$, since $4 = 3+1 = 2+2 = 2+1+1 = 1+1+1+1$. Euler saw in 1748 that

$$\sum_{n\geq 0} p(n)x^n = \frac{1}{(1-x)(1-x^2)(1-x^3)(1-x^4)\cdots},$$

and used it to discover remarkable identities. One of them: the number of ways to write n as a sum of *distinct* parts equals the number of ways to write it as a sum of *odd* parts. (The proof is a single line of algebra with generating functions: try it in the exercises.)

Hardy, Ramanujan and $p(200)$

The partition numbers grow fast: $p(10) = 42$, $p(100) = 190\,569\,292$. In 1918 G. H. Hardy and Srinivasa Ramanujan found an astonishing formula for $p(n)$, an infinite series whose first term is

$$p(n) \approx \frac{1}{4n\sqrt{3}}\, e^{\pi\sqrt{2n/3}}.$$

To test it, Percy MacMahon computed $p(200) = 3\,972\,999\,029\,388$ by hand, using Euler's identities. The first few terms of Hardy and Ramanujan's series gave the right answer to within a fraction of one. Ramanujan also discovered that $p(5n+4)$ is always divisible by 5, $p(7n+5)$ by 7, and $p(11n+6)$ by 11. Explaining why there are no more such patterns took until the twenty-first century.

The thinking skill

Count via coefficients

When an object is built from independent choices, each choice is a factor, and the number of objects of total size n is the coefficient of x^n in the product.

This turns counting into algebra. At university it is the basis of enumerative combinatorics, and it is how probability generating functions handle sums of independent random variables.

Try this next

1 **Two kinds of six.** Try STEP II 2012, Question 1 (`step.maths.org/questions/12-s2-q1`). It counts with products of geometric series. At the end, describe a coin-counting problem that the coefficients answer.

2 **Distinct and odd.** Show that $(1+x)(1+x^2)(1+x^3)\cdots$ is the generating function for partitions into distinct parts. Use $1+x^k = \frac{1-x^{2k}}{1-x^k}$ to show that it equals $\frac{1}{(1-x)(1-x^3)(1-x^5)\cdots}$, and deduce Euler's theorem. Check it for $n = 6$.

3 **A formula.** Show that the number of ways to make n pence using only 1p, 2p and 5p coins is the nearest integer to $\frac{(n+4)^2}{20}$. (Check it first for small n.)

4 **Order matters.** In how many ways can you make n pence by putting 1p and 2p coins into a slot one at a time, if the order matters? (You have met this sequence before.)

Further reading

- **George Andrews & Kimmo Eriksson, *Integer Partitions*** (Cambridge University Press). A short, readable introduction for undergraduates.
- **George Pólya, "On picture-writing"**, *American Mathematical Monthly* 63 (1956). Pólya's classic explanation of generating functions using the problem of changing a dollar (there are 292 ways).
- **Robert Kanigel, *The Man Who Knew Infinity*** (Abacus). The life of Ramanujan, and his partnership with Hardy.

9.6 Strange Dice

The question

STEP I 1993, Question 1

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/93-s1-q1`

The hook

Is there any other way to number two six-sided dice, with positive whole numbers, so that every total from 2 to 12 comes up exactly as often as with ordinary dice? In 1978 Martin Gardner announced in *Scientific American* that there is exactly one: a die numbered $1, 2, 2, 3, 3, 4$ and a die numbered $1, 3, 4, 5, 6, 8$. They had been discovered by Colonel George Sicherman, and are now sold as *Sicherman dice*. A game of Monopoly played with them would be indistinguishable from an ordinary one.

This question is a warm-up: deduce how two dice are numbered from the table of totals. The systematic way to do it, and the proof that Sicherman's dice are the only strange pair, both come from turning each die into a polynomial.

Before you start

Ask yourself If the first die has p_i faces showing i and the second has q_j faces showing j, how many ways are there to throw a total of t?

The number of ways is $\sum_i p_i q_{t-i}$: choose the number i on the first die, then the second must show $t - i$. For the first pair of dice, a total of 6 comes from $1 + 5$ (in 1×1 way), $2 + 4$ (2×1), $3 + 3$ (2×2), and nothing else, since the first die has no 4 or 5 and the second has no 0. That is $1 + 2 + 4 = 7$ ways. ■

Working it through

STAGE 1 The extremes

Call the dice A and B, and write a_i, b_i for the number of faces of each showing i.

Ask yourself There is no way to throw 2, nor 12. What does that say about the 1s and the 6s? There *are* two ways to throw 3. Which die has the 1s?

No total of 2 means $a_1 b_1 = 0$: at most one die has any 1s. A total of 3 can only be $1+2$ or $2+1$, and there are two ways, so some die does have a 1; call it A, so $b_1 = 0$. Then

$$\text{total } 3: \quad a_1 b_2 = 2, \qquad \text{so } (a_1, b_2) = (2,1) \text{ or } (1,2).$$

Similarly, no total of 12 means $a_6 b_6 = 0$.

STAGE 2 Working upwards

Ask yourself Write down the equation for a total of 4, remembering $b_1 = 0$. What does it say in each of the two cases?

A total of 4 is $1+3$ or $2+2$ (and $3+1$ is impossible as $b_1 = 0$), so $a_1 b_3 + a_2 b_2 = 1$.

- **Case (a)**, $a_1 = 2$, $b_2 = 1$: $2b_3 + a_2 = 1$, so $b_3 = 0$ and $a_2 = 1$.
- **Case (b)**, $a_1 = 1$, $b_2 = 2$: $b_3 + 2a_2 = 1$, so $a_2 = 0$ and $b_3 = 1$.

Total 5: $a_1 b_4 + a_2 b_3 + a_3 b_2 = 1$.

- Case (a): $2b_4 + 0 + a_3 = 1$, so $b_4 = 0$, $a_3 = 1$. So far A has $1,1,2,3$ and B has 2; B's other five faces are 5s and 6s.
- Case (b): $b_4 + 0 + 2a_3 = 1$, so $a_3 = 0$, $b_4 = 1$. So far A has 1 and B has $2,2,3,4$; A's other five faces are from $4,5,6$, and B's other two from $5,6$.

STAGE 3 Case (a) fails

Ask yourself In case (a), write down the equation for a total of 7, and use $b_5 + b_6 = 5$.

Total 7: $a_1 b_6 + a_2 b_5 + a_3 b_4 + a_4 b_3 + a_5 b_2 = 2b_6 + b_5 + 0 + 0 + a_5 = 3$. With $b_6 = 5 - b_5$ this says $10 - b_5 + a_5 = 3$, so $b_5 = 7 + a_5 \geq 7$. But B only has six faces. Case (a) is impossible.

STAGE 4 Case (b) succeeds

Ask yourself In case (b), use the totals 6 and 11, and the fact that $a_6 b_6 = 0$.

Total 6: $a_1 b_5 + a_4 b_2 = b_5 + 2a_4 = 4$ (the other products involve $a_2 = a_3 = 0$ or $b_1 = 0$). Since $b_5 \leq 2$ and $b_5 = 4 - 2a_4$ is even, either $b_5 = 0, a_4 = 2$ or $b_5 = 2, a_4 = 1$.

- If $b_5 = 0$ then B is $2,2,3,4,6,6$, so $a_6 = 0$ and A is $1,4,4,5,5,5$. But then total 7 has $a_1 b_6 + a_4 b_3 + a_5 b_2 = 2 + 2 + 6 = 10$ ways, not 3.

- If $b_5 = 2$ then B is $2,2,3,4,5,5$, with $b_6 = 0$, and A is $1,4$ and four faces from 5 and 6. Total 11: $a_6 b_5 = 2a_6 = 6$, so $a_6 = 3$ and $a_5 = 1$.

The only possibility is

$$A:\ 1,4,5,6,6,6, \qquad B:\ 2,2,3,4,5,5.$$

Finally we must check that these dice give the whole table, since we have used only some of the totals. Multiplying out (or using the polynomials below) confirms every entry. ■

Looking back

- **Start at the extremes.** The smallest and largest totals involve only one or two products, so they give the strongest information.
- **Split into cases and kill them.** Each total cut the possibilities down, until only one case survived.
- **Check sufficiency.** The deduction showed what the dice *must* be; a final check showed that they work.

What's really going on

Dice as polynomials

Write each die as a polynomial, with one term x^i for each face showing i. An ordinary die is $x + x^2 + x^3 + x^4 + x^5 + x^6$, and the dice in the question are

$$A(x) = x + x^4 + x^5 + 3x^6, \qquad B(x) = 2x^2 + x^3 + x^4 + 2x^5.$$

Multiplying, each pair of faces contributes $x^i x^j = x^{i+j}$, so the coefficient of x^t in $A(x)B(x)$ is the number of ways to throw a total of t. This is the generating function idea of Section 9.4 once more: the table of totals is just the product $A(x)B(x)$, and our case-by-case deduction was a way of factorising it.

Sicherman's dice

For two ordinary dice the product is $(x + x^2 + \cdots + x^6)^2$. Any pair of dice with the same totals must have polynomials $P(x)$, $Q(x)$ with $P(x)Q(x) = (x + \cdots + x^6)^2$. Now factorise:

$$x + x^2 + \cdots + x^6 = x(1+x)(1+x+x^2)(1-x+x^2),$$

and these factors cannot be broken down further (into polynomials with integer coefficients). So each of P and Q must be built from the factors x, $1+x$, $1+x+x^2$, $1-x+x^2$, each appearing twice in total. Three conditions narrow the choice:

- each die has 6 faces, so $P(1) = Q(1) = 6$; the factors take the values 1, 2, 3, 1 at $x = 1$, so each die gets one $1 + x$ and one $1 + x + x^2$;

- no face shows 0, so $P(0) = Q(0) = 0$: each die gets one factor x;
- the only freedom left is how the two copies of $1 - x + x^2$ are shared. One each gives ordinary dice. Both on one die gives

$$x(1+x)(1+x+x^2) = x + 2x^2 + 2x^3 + x^4,$$
$$x(1+x)(1+x+x^2)(1-x+x^2)^2 = x + x^3 + x^4 + x^5 + x^6 + x^8.$$

These are Sicherman's dice, $1,2,2,3,3,4$ and $1,3,4,5,6,8$ (Figure 9.6.1), and the argument proves they are the only alternative. The key fact is that polynomials, like integers, factorise uniquely into irreducibles: number theory and combinatorics meet.




Figure 9.6.1 Sicherman's dice. Thrown together, they give every total from 2 to 12 with exactly the same frequency as two ordinary dice (right): one way to throw 2, two ways to throw 3, and so on up to six ways to throw 7.

Probability generating functions

In probability the same idea appears as the *probability generating function*: a random variable X taking values $0,1,2,\ldots$ has $G(t) = E(t^X) = \sum P(X = k)t^k$, and for independent X and Y, $E(t^{X+Y}) = E(t^X)E(t^Y)$. Sums of independent random variables correspond to products of generating functions. STEP III 2003 Q14 (in the exercises) finds Sicherman's dice this way, and Section 9.13 uses a related argument to show that no loading of two dice can make all totals equally likely.

The thinking skill

Encode a counting problem as a polynomial

When objects combine by adding their values, represent each one as a polynomial in x. Combining becomes multiplying, and questions about the combinations become questions about factorisation.

Here a question about dice turned into the unique factorisation of polynomials. At university, such encodings are everywhere: in combinatorics, in probability, and in algebra, where polynomials are studied in their own right as a ring with arithmetic like that of the integers.

Try this next

1 **Tetrahedral dice.** Try STEP III 2003, Question 14 (`step.maths.org/questions/03-s3-q14`), which asks Sicherman's question for four-faced dice. (*Hint:* $x + x^2 + x^3 + x^4 = x(1+x)(1+x^2)$.)

2 **Checking the table.** Multiply out $(x + x^4 + x^5 + 3x^6)(2x^2 + x^3 + x^4 + 2x^5)$ and check it against the table in the question.

3 **Three dice.** Are there three six-sided dice, not all standard, labelled with positive integers, whose totals have the same distribution as three standard dice? (*Hint:* the factors of $(x + \cdots + x^6)^3$ now include three copies of $1 - x + x^2$.)

4 **A coin and an octahedron.** An eight-sided die labelled $1, \ldots, 8$ and a two-sided "die" (a coin) labelled $1, 2$: can they be relabelled differently with the same totals?

Further reading

- **Martin Gardner, *Penrose Tiles to Trapdoor Ciphers*** (Mathematical Association of America). Includes the column on Sicherman dice.
- **Joseph Gallian & David Rusin, "Cyclotomic polynomials and nonstandard dice"**, *Discrete Mathematics* 27 (1979). The general theory, for dice with any number of faces.
- **Herbert Wilf, *generatingfunctionology*** (free online), Chapter 1.

9.7 Equal Sums of Powers

The question

STEP I 1992, Question 5

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/92-s1-q5`

The hook

Split the numbers 1 to 16 into two teams of eight:

$$\{1,4,6,7,10,11,13,16\} \quad \text{and} \quad \{2,3,5,8,9,12,14,15\}.$$

Each team sums to 68. The sums of their squares are both 748. The sums of their cubes are both 9248. It is hard to believe this could be anything but a fluke, yet it is the third step of a pattern that goes on for ever: split 1 to 32 the right way and the fourth powers agree as well.

The split was found by Eugène Prouhet in 1851, and the rule behind it, the *Thue–Morse sequence*, turns up in an astonishing range of places: in number theory, in the theory of computation, in chess, and in the fairest way to take turns. This question uncovers it through a polynomial and its derivatives.

Before you start

Ask yourself If a polynomial $\mathrm{p}(x) = \sum c_k x^k$ has $\mathrm{p}(1) = 0$, what does that say about the coefficients? What about $\mathrm{p}'(1) = 0$?

$\mathrm{p}(1) = \sum c_k$, so $\mathrm{p}(1) = 0$ says the coefficients add to zero. And $\mathrm{p}'(x) = \sum k c_k x^{k-1}$, so $\mathrm{p}'(1) = \sum k c_k$. If the coefficients are all ± 1, these say that the positions of the $+1$s and the -1s are equal in number, and have equal sums.

Working it through

STAGE 1 A triple root

Ask yourself Factorise $1-x^2$ and $1-x^4$. How many factors of $1-x$ appear?

Since $1-x^2=(1-x)(1+x)$ and $1-x^4=(1-x)(1+x)(1+x^2)$,

$$p_0(x)=(1-x)^3(1+x)^2(1+x^2),$$

so $(1-x)^3$ is a factor. ■

STAGE 2 Derivatives at 1

Ask yourself If $(1-x)^3$ divides p_0, what power of $(1-x)$ divides p_0'? And then $p_1=xp_0'$ and p_1'?

Write $p_0(x)=(1-x)^3q(x)$. By the product rule, $p_0'(x)=-3(1-x)^2q(x)+(1-x)^3q'(x)$, which has $(1-x)^2$ as a factor. So $p_0'(1)=0$. Then $p_1(x)=xp_0'(x)$ also has $(1-x)^2$ as a factor, and the same argument shows p_1' has $(1-x)$ as a factor, so $p_1'(1)=0$. ■

STAGE 3 Reading off the coefficients

Multiplying out, $(1-x)(1-x^2)=1-x-x^2+x^3$, and then

$$p_0(x)=1-x-x^2+x^3-x^4+x^5+x^6-x^7.$$

Ask yourself What are $p_0'(1)$ and $p_1'(1)$ in terms of the coefficients c_k?

Since $p_1(x)=xp_0'(x)=\sum kc_kx^k$, we have $p_1'(1)=\sum k^2c_k$. The coefficient is $+1$ at $k=0,3,5,6$ and -1 at $k=1,2,4,7$. So $p_0'(1)=0$ and $p_1'(1)=0$ say

$$0+3+5+6=1+2+4+7\quad(=14),$$
$$0^2+3^2+5^2+6^2=1^2+2^2+4^2+7^2\quad(=70),$$

which, dropping the 0, are the identities required. ■

STAGE 4 One more factor

Pause and try To make the cubes balance too, we need a fourth factor of $(1-x)$. Which polynomial should replace p_0?

Take $P(x)=(1-x)(1-x^2)(1-x^4)(1-x^8)$, which has $(1-x)^4$ as a factor. Its coefficients are $p_0(x)-x^8p_0(x)$: the pattern of signs for 0 to 7, followed by the same pattern reversed in sign for 8 to 15:

k	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
c_k	+	−	−	+	−	+	+	−	−	+	+	−	+	−	−	+

The argument of Stage 2, applied one more time, shows that $\sum c_k k^r = 0$ for $r = 0, 1, 2, 3$. Take the $+$ positions $\{0, 3, 5, 6, 9, 10, 12, 15\}$ and the $-$ positions $\{1, 2, 4, 7, 8, 11, 13, 14\}$. For $r = 1, 2, 3$, the term from 0 is zero, so it can be dropped. So the order

$$a_1, \ldots, a_8 = 1, 2, 4, 7, 8, 11, 13, 14, \qquad a_9, \ldots, a_{15} = 3, 5, 6, 9, 10, 12, 15$$

works: both halves sum to 60, their squares to 620, and their cubes to 7200. ■

Looking back

- **Derivatives detect repeated roots.** A factor $(1 - x)^m$ forces the first $m - 1$ derivatives to vanish at $x = 1$.
- **Multiplying by x before differentiating** turned $\sum k c_k$ into $\sum k^2 c_k$, and so on to higher powers.
- **Doubling.** Each extra factor $(1 - x^{2^m})$ doubled the length and added one more power that balances.

What's really going on

The Thue–Morse sequence

The signs of the coefficients of $(1 - x)(1 - x^2)(1 - x^4)(1 - x^8) \cdots$ form the *Thue–Morse sequence*. Writing 0 for $+$ and 1 for $-$:

$$0\,1\,1\,0\,1\,0\,0\,1\,1\,0\,0\,1\,0\,1\,1\,0 \ldots$$

There are three equivalent descriptions.

- **Doubling:** start with 0, and repeatedly append the complement (swap 0s and 1s) of what you have: $0 \to 01 \to 0110 \to 01101001 \to \cdots$.
- **Binary digits:** the kth term is 0 if k has an even number of 1s when written in binary, and 1 if odd. (This is why the sign of x^k is $(-1)^{\text{number of 1s in } k}$: to make x^k from the product, you choose the factors x^{2^m} that make up k in binary, and each brings a minus sign.)
- **Polynomial:** the coefficients of $\prod_m (1 - x^{2^m})$, as in the question.

Splitting $0, 1, \ldots, 2^N - 1$ by the Thue–Morse sequence gives two sets whose rth powers have equal sums for every $r < N$ (Figure 9.7.1). This is the *Prouhet–Tarry–Escott problem*: find two different sets of integers with equal sums of rth powers for $r = 1, \ldots, k$. Prouhet's solution uses 2^{k+1} numbers; it is a famous open problem to find out how few numbers are really needed.



$1+4+6+7+10+11+13+16 = 68$

$2+3+5+8+9+12+14+15 = 68$

squares: $748 = 748$ cubes: $9248 = 9248$ fourth powers: $122\,692 \neq 121\,156$

Figure 9.7.1 The numbers 1 to 16 split by the Thue–Morse sequence. The two sets have equal sums, equal sums of squares and equal sums of cubes, but not equal sums of fourth powers; for that you need 1 to 32. Reading the colours left to right gives the pattern teal–plum–plum–teal, plum–teal–teal–plum, ...

The fairest way to take turns

Two children take turns choosing sweets from a pile, best first. If they alternate $ABABAB\ldots$, then A always gets the better sweet of each pair, a systematic advantage. The order $ABBA$ is fairer, and $ABBABAAB$ fairer still: this is the Thue–Morse sequence, and the balanced power sums are exactly the reason. If the sweets have values given by a smooth function of their position, expanding that function as a polynomial shows that the Thue–Morse order cancels its leading terms. The same idea has been proposed for penalty shoot-outs in football (the "$ABBA$" order trialled by football's rule-makers from 2017) and is used for the order of serving in tennis tie-breaks.

Thue, Morse and Euwe

The sequence has been discovered many times. Prouhet used it in 1851. Axel Thue rediscovered it in 1912 while studying sequences without repetitive patterns: it never contains the same block three times in a row (it is "cube-free"). Marston Morse found it again in 1921 in the study of geodesics on surfaces. And the chess world champion Max Euwe used it in 1929 to show that the rule then in force for drawing a game by repetition did not prevent infinitely long games. Today it is a central example in the theory of *automatic sequences*, sequences computed by simple machines reading the binary digits of n.

The thinking skill

Encode a combinatorial property in a polynomial

Balanced sums of powers are about where the $+$ and $-$ signs fall. A polynomial with those signs as coefficients turns the property into a statement about roots.

The question never asks you to search: a repeated root at $x = 1$ does all the work. Using algebra to produce combinatorial objects, rather than finding them by trial, is a recurring idea in university combinatorics.

Try this next

1 **Shifting.** Try STEP II 2025, Question 8 (in OCR's download of the 2025 papers, `www.ocr.org.uk/Images/750486-2025-past-papers.zip`). It shows how to turn a split that balances sums and squares into a larger one that balances cubes as well, by shifting one list and combining.

2 **Squares of squares.** (Also from STEP II, 2025.) The sets $A = \{1, 3, 4, 5, 9, 11\}$ and $B = \{2, 6, 7, 8, 10\}$ have equal sums and equal sums of squares. Split $\{n^2, (n+1)^2, \ldots, (n+11)^2\}$ into two sets with equal sums, for every positive integer n. (*Hint:* add 0 to the second set.)

3 **Translation invariance.** Show that if two sets of numbers have equal sums of rth powers for $r = 0, 1, \ldots, k$, then so do the sets obtained by adding the same number c to every element. Why does this mean that the question's sets $\{1, \ldots, 15\}$ could equally be replaced by any 16 consecutive integers?

4 **Fair sweets.** Sweets have values $1, \frac{1}{2}, \frac{1}{3}, \ldots, \frac{1}{8}$. Compare the totals A gets with the orders $ABABABAB$ and $ABBABAAB$.

Further reading

- **Jean-Paul Allouche & Jeffrey Shallit, "The ubiquitous Prouhet–Thue–Morse sequence"**, in *Sequences and Their Applications* (Springer, 1999). A survey of the sequence's many appearances; free online.
- **Matt Parker, *Things to Make and Do in the Fourth Dimension*** (Penguin). Includes equal power sums and other number curiosities.
- **Joshua Cooper & Aaron Dutle, "Greedy Galois games"**, *American Mathematical Monthly* 120 (2013). Fair turn-taking and Thue–Morse.

9.8 Nobody Gets Their Own Shirt

The question

STEP II 2016, Question 12

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/16-s2-q12`

The hook

A football team of eleven players changes in a hurry and grabs shirts at random. What is the chance that nobody ends up in their own shirt? Now suppose it is a squad of fifty, or a crowd of a million. Does the chance go up, because there are more people to get it wrong? Or down, because there are more people who might get it right by luck? Remarkably, it hardly changes at all: from five people onwards, the chance is within half a per cent of $1/e \approx 0.368$.

This is the oldest problem in combinatorial probability, posed by Pierre de Montmort in 1708 as the *problème des rencontres*. The question solves it with the *inclusion–exclusion principle*, a tool for counting things that overlap, and the answer is one of the most unexpected appearances of e in mathematics.

Before you start

Pause and try Try $n = 3$: list all six orders of cards $1, 2, 3$ and count how many have no card in its own place. Then try $n = 4$, if you are patient (there are 24 orders).

For $n = 3$ only 231 and 312 have no card in its place, so the probability is $\frac{2}{6} = \frac{1}{3}$. For $n = 4$ there are 9 such orders out of 24, giving $\frac{3}{8} = 0.375$. An arrangement with no fixed card is called a *derangement*.

Working it through

STAGE 1 **Inclusion–exclusion for three events**

Ask yourself How can you write $A \cup B \cup C$ as a union of two events, so that the two-event formula applies?

Treat $A \cup B$ as a single event and apply the formula to it and C:

$$\mathrm{P}(A \cup B \cup C) = \mathrm{P}(A \cup B) + \mathrm{P}(C) - \mathrm{P}((A \cup B) \cap C).$$

Now $(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$, and applying the formula once more, $\mathrm{P}((A \cap C) \cup (B \cap C)) = \mathrm{P}(A \cap C) + \mathrm{P}(B \cap C) - \mathrm{P}(A \cap B \cap C)$. Expanding $\mathrm{P}(A \cup B)$ too and collecting terms gives the result. ■

For four events the pattern continues: add the four single probabilities, subtract the six pairwise intersections, add the four triple intersections, and subtract $\mathrm{P}(A \cap B \cap C \cap D)$.

STAGE 2 The probabilities

Ask yourself If card i is in position i, how many ways are there to arrange the rest? And if cards i and j are both fixed?

There are $n!$ equally likely orders. If card i is fixed in position i, the other $n-1$ cards can be in any order: $(n-1)!$ ways. So

$$\mathrm{P}(E_i) = \frac{(n-1)!}{n!} = \frac{1}{n}, \qquad \mathrm{P}(E_i \cap E_j) = \frac{(n-2)!}{n!} = \frac{1}{n(n-1)},$$
$$\mathrm{P}(E_i \cap E_j \cap E_k) = \frac{(n-3)!}{n!} = \frac{1}{n(n-1)(n-2)}.$$

In general, the probability that k specified cards are all in their own positions is $\frac{(n-k)!}{n!}$.

STAGE 3 Adding it up

Ask yourself In the inclusion–exclusion formula for $\mathrm{P}(E_1 \cup \cdots \cup E_n)$, how many terms are there involving intersections of k events, and what is their total?

There are $\binom{n}{k}$ ways to choose k of the events, and each intersection has probability $\frac{(n-k)!}{n!}$. So the k-fold terms add up to

$$\binom{n}{k}\frac{(n-k)!}{n!} = \frac{n!}{k!\,(n-k)!} \cdot \frac{(n-k)!}{n!} = \frac{1}{k!}.$$

Everything except the $k!$ cancels. With alternating signs,

$$\mathrm{P}(\text{at least one card in its place}) = 1 - \frac{1}{2!} + \frac{1}{3!} - \cdots + (-1)^{n+1}\frac{1}{n!}. \qquad \blacksquare$$

So the probability of a derangement is

$$d_n = 1 - 1 + \frac{1}{2!} - \frac{1}{3!} + \cdots + (-1)^n \frac{1}{n!} = \sum_{k=0}^{n} \frac{(-1)^k}{k!}.$$

STAGE 4 **Exactly one**

> **Ask yourself** For exactly one card to be in place, which card is it, and what must the others do?

Choose which card is in its place (n ways). The probability that card i is in place is $\frac{1}{n}$, and given that, the other $n-1$ cards must form a derangement among themselves, which has probability d_{n-1}. So

$$\mathrm{P}(\text{exactly one}) = n \cdot \frac{1}{n} \cdot d_{n-1} = \sum_{k=0}^{n-1} \frac{(-1)^k}{k!}. \qquad \blacksquare$$

Looking back

- **Overcount, then correct.** Adding the $\mathrm{P}(E_i)$ counts arrangements with two fixed cards twice; subtracting the pairs corrects this but goes too far for triples; and so on.
- **Symmetry makes the terms equal.** Every k-fold intersection has the same probability, so each layer of the formula is one number times a binomial coefficient.
- **Reduce to a smaller case.** "Exactly one" became a derangement of $n-1$ cards.

What's really going on

Where e comes from

The series $\sum(-1)^k/k!$ is the exponential series $e^x = \sum x^k/k!$ at $x = -1$. So as $n \to \infty$,

$$d_n \to e^{-1} \approx 0.3679,$$

and the probability of exactly one match tends to the same limit. The convergence is extraordinarily fast (Figure 9.8.1): the error in stopping at $1/n!$ is less than $1/(n+1)!$, so already for $n = 7$ the answer agrees with $1/e$ to four decimal places. In fact the number of derangements of n objects, $n!\,d_n$, is exactly the nearest integer to $n!/e$.




Figure 9.8.1 The probability that a random shuffle of n cards leaves no card in its own place. It alternates above and below $1/e$, and from $n = 6$ onwards it is indistinguishable from $1/e$ on this scale.

The number of matches is Poisson

More is true. Let N be the number of cards in their own place. Then $\mathrm{P}(N = k) \to \frac{e^{-1}}{k!}$ for each k: the number of matches has, in the limit, the *Poisson distribution* with mean 1. You can see why the mean is 1 without any calculation: N is the sum of the indicators of $E_1, \ldots, E_n$, each with expectation $\frac{1}{n}$, and expectation adds even though the events are not independent. STEP III 2004 Q12 (in the exercises) shows that the variance is 1 too. The Poisson distribution appears whenever there are many unlikely events that are nearly independent; derangements are a classic example, and the subject of Section 12.11 in Chapter 12.

Montmort's game of *treize*

Pierre Rémond de Montmort, a French nobleman, studied games of chance in his *Essay d'analyse sur les jeux de hazard* (1708). In the game of *treize* (thirteen), a player deals thirteen cards, calling out "one, two, three, ..." and wins if the card dealt ever matches the number called. Montmort found the probability of winning, and in correspondence with Nicolaus Bernoulli the general formula was established by 1713. Euler rediscovered it in 1753. The inclusion–exclusion principle was later formalised by Abraham de Moivre and, in its general form, by James Joseph Sylvester in the nineteenth century.

The thinking skill

Inclusion–exclusion

To count a union of overlapping sets, add the sizes of the sets, subtract the pairwise overlaps, add the triple overlaps, and so on.

It is most powerful when symmetry makes all the overlaps of each size equal, as here. At university it appears in combinatorics, in probability and in number theory, where it counts integers not divisible by any of a list of primes, and leads to Euler's totient function and the Möbius function.

Try this next

1 **Shirts.** Try STEP III 2004, Question 12 (`step.maths.org/questions/04-s3-q12`). It counts the players wearing their own shirts with indicator variables, finds the mean and variance, and compares the result with a Poisson distribution.

2 **A recurrence.** Let D_n be the number of derangements of n objects. By considering where card 1 goes, and whether the card in that position goes to position 1, show that $D_n = (n-1)(D_{n-1} + D_{n-2})$. Check it against $D_3 = 2$, $D_4 = 9$.

3 **Exactly k.** Show that the probability that exactly k cards are in their own places is $\frac{1}{k!} d_{n-k}$, and that it tends to $\frac{e^{-1}}{k!}$.

4 **Coprime integers.** Use inclusion–exclusion to count the integers from 1 to 30 divisible by none of 2, 3 and 5. How does your answer relate to $30\left(1-\frac{1}{2}\right)\left(1-\frac{1}{3}\right)\left(1-\frac{1}{5}\right)$?

Further reading

- **Geoffrey Grimmett & Dominic Welsh, *Probability: An Introduction*** (Oxford University Press). A first-year text that covers inclusion–exclusion and the matching problem.
- **William Feller, *An Introduction to Probability Theory and Its Applications*, Vol. 1** (Wiley), Chapter IV. A classic, with many variations on the matching problem.
- **Miklós Bóna, *A Walk Through Combinatorics*** (World Scientific), on inclusion–exclusion.

9.9 The Birthday Problem

The question

STEP II 2007, Question 13

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/07-s2-q13`

The hook

How many people do you need in a room for it to be more likely than not that two of them share a birthday? Most people guess somewhere around 180, half of 365. The answer is 23. In a class of thirty there is a 70% chance of a shared birthday; at a party of fifty, 97%.

The surprise comes from confusing two different questions: "does someone share *my* birthday?" (which does need about 253 people) and "do *any two people* share a birthday?". With 23 people there are $\binom{23}{2} = 253$ pairs, and each pair is a chance for a match. This question makes that intuition precise, and the same mathematics decides how secure the cryptography protecting your bank account is.

Before you start

Ask yourself Why is it easier to calculate the probability that *no* two guests share a birthday? Line the guests up: what is the chance that the second avoids the first's birthday? That the third avoids both?

"At least two share" covers many overlapping cases (two share, three share, two different pairs, ...). Its complement, "all different", is a single clean event. The second guest avoids the first's birthday with probability $\frac{364}{365}$; given that, the third avoids both with probability $\frac{363}{365}$; and so on.

Working it through

STAGE 1 A useful approximation

Ask yourself What is the tangent line to $y = e^{-x}$ at $x = 0$?

Since $e^{-x} = 1 - x + \frac{x^2}{2} - \cdots$, for small x we have $e^{-x} \approx 1 - x$. With $x = r/n$,

$$\frac{n-r}{n} = 1 - \frac{r}{n} \approx e^{-r/n}. \qquad \blacksquare$$

The error is about $\frac{r^2}{2n^2}$, small when r is much smaller than n.

STAGE 2 The probability of a shared birthday

The probability that all k birthdays are different is

$$\frac{365}{365} \cdot \frac{364}{365} \cdot \frac{363}{365} \cdots \frac{365-(k-1)}{365} = \prod_{r=0}^{k-1} \frac{365-r}{365}.$$

Using the approximation on each factor, and the fact that exponentials multiply by adding their exponents,

$$\prod_{r=0}^{k-1} e^{-r/365} = \exp\Big(-\frac{0+1+\cdots+(k-1)}{365}\Big) = \exp\Big(-\frac{k(k-1)}{730}\Big).$$

So the probability of at least one shared birthday is approximately $1 - e^{-k(k-1)/730}$. $\blacksquare$

Ask yourself Look at the exponent: $\frac{k(k-1)}{730} = \binom{k}{2}\Big/365$. What does that say in words?

It is the number of pairs of guests, times the probability $\frac{1}{365}$ that a given pair shares a birthday: the expected number of shared pairs. The approximation says the chance of *no* match is $e^{-(\text{expected number of matches})}$, a hint of the Poisson distribution again, as in Section 9.8.

STAGE 3 Twenty-three

We need $1 - e^{-k(k-1)/730} \geq \frac{1}{2}$, that is $e^{-k(k-1)/730} \leq \frac{1}{2}$, or

$$\frac{k(k-1)}{730} \geq \ln 2 \approx \frac{253}{365}, \qquad \text{that is} \qquad k(k-1) \geq 506.$$

Since $22 \times 21 = 462$ and $23 \times 22 = 506$, the smallest value is $k = 23$. $\blacksquare$

(The exact probability for 23 guests is 0.507, and for 22 guests 0.476, so the approximation gives the right answer.)

STAGE 4 Sharing with the host

Ask yourself For each guest, what is the probability of *not* sharing the host's birthday? Are the guests independent of each other in this respect?

Each guest independently misses the host's birthday with probability $\frac{364}{365} \approx e^{-1/365}$. So the probability that none of k guests shares it is $\left(\frac{364}{365}\right)^k \approx e^{-k/365}$, and we need

$$e^{-k/365} \leq \tfrac{1}{2}, \qquad \text{that is} \qquad k \geq 365 \ln 2 \approx 253.$$

So there must be 253 guests. ■

The contrast is the whole point: 253 guests to match one particular person, but only 23 for some pair to match, because 23 people make 253 pairs.




Figure 9.9.1 The probability that at least two of k people share a birthday: exact values (teal) and the approximation $1 - e^{-k(k-1)/730}$ (plum, dashed). The two are very close, and both pass $\frac{1}{2}$ between $k = 22$ and $k = 23$.

Looking back

- **Use the complement.** "All different" was a single product; "at least one match" was a mess.
- **Linearise, then exponentiate.** Replacing each factor $1 - x$ by e^{-x} turned a product into a sum.
- **Count pairs.** The exponent was the number of pairs times the chance that a pair matches. The square in $k(k-1)$ is the whole explanation of the paradox.

What's really going on

The square-root law

With n equally likely days instead of 365, the same argument gives

$$\mathrm{P}(\text{some match among } k) \approx 1 - e^{-k^2/(2n)},$$

so matches become likely once k is around $\sqrt{n}$, not n. More precisely, the probability reaches $\frac{1}{2}$ at $k \approx 1.18\sqrt{n}$, and the expected number of people you need to meet before the first repeat is about $\sqrt{\pi n/2}$ (for $n = 365$, about 24.6). This *square-root law* is the reason coincidences are so common: in any large collection of events, some pair is likely to match in some way, even though any particular match is unlikely.

Birthday attacks

Computers check that a file has not been tampered with by computing a *hash*: a fingerprint of, say, $n = 2^{128}$ possible values. A forger who wants two documents with the same hash (a harmless contract and a fraudulent one) does not need to try 2^{128} documents. By the birthday argument, generating about $\sqrt{2^{128}} = 2^{64}$ variants of each will probably produce a match. This *birthday attack* is why modern hash functions use 256 bits or more: half the bits are lost to the square-root law. Similar calculations decide how long random identifiers must be so that two computers never accidentally choose the same one.

Who first asked?

The birthday problem is usually credited to Richard von Mises, who discussed it in 1939, though it circulated earlier. Harold Davenport is also said to have known it earlier, and not to have published it because it seemed too simple. As Warren Weaver told the story, he liked to demonstrate it at dinners of army officers during the Second World War: he bet that two of the twenty-two people present shared a birthday, and lost, until a waitress revealed that she shared her birthday with one of the officers.

The thinking skill

Calculate the complement; approximate carefully

When "at least one" is complicated, find the probability of "none". When a product is awkward, approximate each factor by an exponential and add the exponents.

Both moves are standard at university. The second, $1 - x \approx e^{-x}$, is the start of the theory of Poisson approximation and of the asymptotic methods used to estimate quantities too complicated to calculate exactly.

Try this next

1 **The first repeat.** Try STEP II 2014, Question 13 (`step.maths.org/questions/14-s2-q13`). It studies the first repeated value from a random number generator, which is the birthday problem in another guise, and turns the probabilities into two surprising identities.

2 **Three in common.** Roughly how many people are needed for a better-than-even chance that some *three* share a birthday? (*Hint:* the expected number of triples sharing a birthday is $\binom{k}{3}/365^2$. Set this to about $\ln 2$.) The true answer is 88.

3 **Near misses.** How many people are needed for an even chance that two have birthdays on the same day or on adjacent days?

4 **Uneven birthdays.** Real birthdays are not uniform. Show that if the days have probabilities $p_1, \ldots, p_{365}$, the chance that two people share a birthday is $\sum p_i^2$, and that this is smallest when all p_i are equal. (So non-uniformity only makes matches more likely.)

Further reading

- **William Feller, *An Introduction to Probability Theory and Its Applications*, Vol. 1** (Wiley), Chapter II. The birthday problem and its approximations.
- **Persi Diaconis & Frederick Mosteller, "Methods for studying coincidences"**, *Journal of the American Statistical Association* 84 (1989). Why coincidences are common; very readable.
- **David Hand, *The Improbability Principle*** (Bantam). The square-root law and other reasons why unlikely things happen all the time.

9.10 Always Able to Give Change

The question

STEP I 2011, Question 12

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/11-s1-q12`

The hook

The pattern is irresistible: $\frac{m}{m+1}$, $\frac{m-1}{m+1}$, $\frac{m-2}{m+1}$. Surely, with n people holding £2 coins, the answer is $\frac{m-n+1}{m+1}$. It is, and the slickest proof is one of the most elegant ideas in combinatorics: the *reflection principle*. It turns a hard count (queues that never run out of change) into an easy one by reflecting part of a picture in a line.

The same problem has many disguises: an election count in which the winner is never behind (Bertrand's ballot problem of 1887), a gambler who never goes broke, brackets that match correctly. When $m = n$ the count gives the *Catalan numbers*, perhaps the most frequently occurring sequence in combinatorics after the binomial coefficients.

Before you start

Ask yourself Each £1 customer gives you a coin; each £2 customer takes one away (as change). When do you get stuck?

Keep track of the number of £1 coins you hold. It goes up by 1 for each £1 customer and down by 1 for each £2 customer. You are stuck exactly when a £2 customer arrives and you hold no coins: the count would go to -1. So you succeed if and only if, at every point in the queue, the number of £1 customers so far is at least the number of £2 customers so far.

Working it through

STAGE 1 One £2 coin

With $n = 1$, you fail only if the £2 customer is first. The £2 customer is equally likely to

be in any of the $m+1$ positions, so

$$\mathrm{P}(\text{success}) = 1 - \frac{1}{m+1} = \frac{m}{m+1}. \qquad \blacksquare$$

STAGE 2 Two £2 coins

Ask yourself The first person must have a £1 coin. What can happen with the second person, and in each case, what situation are you left in?

The first person has a £1 coin with probability $\frac{m}{m+2}$; otherwise you fail at once. Then you hold one coin, and the remaining queue has $m-1$ £1 people and two £2 people.

- If the second person has a £1 coin (probability $\frac{m-1}{m+1}$), you hold two coins, enough to change both £2 coins whenever they come. Success is certain.
- If the second person has a £2 coin (probability $\frac{2}{m+1}$), you are back to no coins, with $m-1$ £1 people and one £2 person left. By part (i) with $m-1$ in place of m, you succeed with probability $\frac{m-1}{m}$.

So

$$\mathrm{P}(\text{success}) = \frac{m}{m+2}\left(\frac{m-1}{m+1} + \frac{2}{m+1}\cdot\frac{m-1}{m}\right) = \frac{m}{m+2}\cdot\frac{m-1}{m+1}\cdot\frac{m+2}{m} = \frac{m-1}{m+1}. \qquad \blacksquare$$

Pause and try Try the same approach for $n = 3$. It works, but the cases multiply. Is there a way to count the successful queues all at once?

STAGE 3 Queues as paths

Ask yourself Draw the number of coins you hold against the number of people served. What does a queue look like? What does a successful queue look like?

A queue is a path made of m up-steps (£1) and n down-steps (£2), starting at height 0 and ending at height $m-n$. There are $\binom{m+n}{n}$ such paths, all equally likely. A queue is successful exactly when its path never goes below 0. So we need to count the *bad* paths, those that touch height -1.

STAGE 4 The reflection

Ask yourself Take a bad path. Find the first moment it touches -1, and reflect the part of the path *before* that moment in the line $y = -1$. Where does the new path start, and where does it end?

The reflected path starts at -2 (the mirror image of 0 in the line $y = -1$) and ends where the original did, at $m-n$ (Figure 9.10.1). Conversely, every path from -2 to

$m - n$ must cross $y = -1$ somewhere (it starts below and ends above, since $m \geq n$); reflecting it before its first visit gives back a bad path from 0. So bad paths from 0 correspond exactly to *all* paths from -2 to $m - n$.

A path from -2 to $m - n$ in $m + n$ steps rises by $m - n + 2$, so it has $m + 1$ up-steps and $n - 1$ down-steps: there are $\binom{m+n}{n-1}$ of them. Hence

$$\mathrm{P}(\text{failure}) = \frac{\binom{m+n}{n-1}}{\binom{m+n}{n}} = \frac{n!\,m!}{(n-1)!\,(m+1)!} = \frac{n}{m+1},$$

and

$$\mathrm{P}(\text{success}) = 1 - \frac{n}{m+1} = \frac{m-n+1}{m+1}.$$

With $n = 3$ this is $\frac{m-2}{m+1}$. ■




Figure 9.10.1 A queue with $m = 4$ £1 coins and $n = 3$ £2 coins, drawn as a path (teal). It first hits -1 at the third person (amber), when you cannot give change. Reflecting the path up to that point in the line $y = -1$ gives a path from -2 (plum, dashed) to the same end point. This matching between bad queues and paths starting at -2 is the reflection principle.

Looking back

- **Find the right state.** The only thing that mattered was the number of coins held, which turned the queue into a walk.
- **Condition on the first steps.** Parts (i) and (ii) were done by asking what the first few people do, reducing to smaller cases.
- **Count the complement by a bijection.** The bad paths were hard to count directly, but reflection matched them with a set that was easy to count.

What's really going on

Catalan numbers

When $m = n$ (exactly enough £1 coins), the number of successful queues is

$$\binom{2n}{n} - \binom{2n}{n-1} = \frac{1}{n+1}\binom{2n}{n},$$

the nth *Catalan number* C_n: $1, 1, 2, 5, 14, 42, 132, \ldots$. Richard Stanley's book *Catalan Numbers* lists more than two hundred different things that they count. Among them: the number of ways to arrange n pairs of brackets correctly ("(()()" is allowed, ")()(()" is not, by exactly the coin argument with "(" as £1 and ")" as £2), the number of ways to cut a polygon with $n+2$ sides into triangles by non-crossing diagonals, and the number of ways to bracket a product of $n+1$ factors.

Random walks

Queues are random walks, and the reflection principle is one of the basic tools for studying them. It shows, for example, that a fair random walk starting at 0 is certain to return to 0 eventually, and that the probability that it stays strictly positive for the first $2n$ steps is $\frac{1}{2}\binom{2n}{n}2^{-2n} \approx \frac{1}{2\sqrt{\pi n}}$. In continuous form, the same reflection argument applied to *Brownian motion* gives the distribution of its maximum. This is the basis of the pricing of certain financial options, whose payoff depends on whether a share price ever crosses a barrier.

Bertrand's ballot problem

In 1887 Joseph Bertrand asked: in an election where candidate A receives a votes and B receives $b < a$, what is the probability that A is strictly ahead throughout the count? The answer, $\frac{a-b}{a+b}$, was proved in the same year by Désiré André, whose argument is usually remembered as the reflection principle (although historians have pointed out that André's actual proof used a slightly different bijection). Our raffle problem is a variant in which ties are allowed: success means the £1 count never falls *below* the £2 count. One of the exercises deduces Bertrand's answer from ours.

The thinking skill

Bijective counting

To count a set, find a one-to-one correspondence with a set you can count.

Reflection matched bad queues with unrestricted paths from a different start. Bijections are the most satisfying proofs in combinatorics, because they explain *why* two numbers are equal rather than just showing that they are. At university you will meet bijective proofs throughout combinatorics, and the reflection principle again in the study of random walks and Brownian motion.

Try this next

1 **Three by cases.** Prove part (iii) by conditioning on the first people in the queue, as in part (ii), to see how much work the reflection principle saves.

2 **Bertrand's ballot.** In an election, A gets a votes and B gets $b < a$. Show that the probability that A is strictly ahead throughout the count is $\frac{a-b}{a+b}$. (*Hint:* the first vote must be for A; after that, A must never fall behind B by more than…)

3 **Brackets.** List the 5 correct arrangements of three pairs of brackets, and the 5 ways to cut a pentagon into triangles. Can you find a bijection between them?

4 **A £5 note.** A queue contains m people with £1 coins and n with £5 notes, for a ticket costing £1. Explain why the reflection principle no longer gives a simple answer. (This generalisation is much harder.)

Further reading

- **William Feller, *An Introduction to Probability Theory and Its Applications*, Vol. 1** (Wiley), Chapter III. The classic treatment of random walks and the reflection principle.
- **Richard Stanley, *Catalan Numbers*** (Cambridge University Press). Over two hundred interpretations, with exercises.
- **Marc Renault, "Lost (and found) in translation: André's actual method and its application to the generalized ballot problem"**, *American Mathematical Monthly* 115 (2008). The history of the reflection principle.

9.11 Collecting the Set

The question

STEP II 1994, Question 13

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/94-s2-q13`

The hook

A football sticker album has 600 stickers to collect, sold one at a time at random. How many stickers should you expect to buy to fill the album? The first few hundred go quickly, since almost every sticker is new. But as the album fills, new stickers get rarer, and the very last one takes on average 600 purchases on its own. The total comes to about 4200, seven times the size of the album.

This is the *coupon collector's problem*. Its answer combines two ideas: the waiting time for a success, and the harmonic series $1 + \frac{1}{2} + \frac{1}{3} + \cdots$, which grows like $\ln n$. You met the harmonic series diverging in Section 2.3; here its slow growth is exactly what you pay for completing a collection.

Before you start

Ask yourself If each packet independently gives a new card with probability p, what is the expected number of packets you open up to and including the first new card?

The number of packets X has a *geometric distribution*: $\mathrm{P}(X = k) = (1-p)^{k-1}p$. Its expectation is $\frac{1}{p}$. (One quick way to see this: $\mathrm{E}(X) = 1 + (1-p)\mathrm{E}(X)$, since you open one packet, and with probability $1-p$ you are back where you started.) If a new card turns up a third of the time, you expect to wait three packets.

Working it through

STAGE 1 Waiting for a new card

With r different cards already, a packet gives a new card with probability $\frac{n-r}{n}$, independently of previous packets. So the expected number of packets until the next new card is

$$\frac{1}{(n-r)/n} = \frac{n}{n-r}. \qquad \blacksquare$$

STAGE 2 Comparing a sum with an integral

Ask yourself On the interval $r \le x \le r+1$, between which two values does $\frac{1}{x}$ lie? What does that say about the area under the curve?

Since $\frac{1}{x}$ is decreasing, $\frac{1}{r+1} \le \frac{1}{x} \le \frac{1}{r}$ on $[r, r+1]$. Integrating over an interval of length 1 gives

$$\frac{1}{r+1} \le \int_r^{r+1} \frac{\mathrm{d}x}{x} \le \frac{1}{r}. \qquad \blacksquare$$

In pictures (Figure 9.11.1): the area under the curve is squeezed between a rectangle of height $\frac{1}{r+1}$ and one of height $\frac{1}{r}$. Adding these from $r = 1$ to $r = n-1$, the integrals join up to $\int_1^n \frac{\mathrm{d}x}{x} = \ln n$, and so

$$\frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} \le \ln n \le 1 + \frac{1}{2} + \cdots + \frac{1}{n-1}. \qquad \blacksquare$$




Figure 9.11.1 On each interval $[r, r+1]$ the area under $y = 1/x$ lies between the teal rectangle (height $\frac{1}{r+1}$) and the larger plum one (height $\frac{1}{r}$). Adding up, $\ln n$ lies between $\frac{1}{2} + \cdots + \frac{1}{n}$ and $1 + \frac{1}{2} + \cdots + \frac{1}{n-1}$. The gaps between the curve and the plum rectangles add up to a finite total: Euler's constant $\gamma \approx 0.577$.

STAGE 3 Completing the set

Ask yourself The total number of packets is the sum of the waiting times for the first card, the second new card, and so on. What is its expectation?

Let X_r be the number of packets needed to go from r different cards to $r+1$. The total is $X_0 + X_1 + \cdots + X_{n-1}$, and since expectation adds,

$$\mathrm{E}(\text{total}) = \sum_{r=0}^{n-1} \frac{n}{n-r} = n\Big(\frac{1}{n} + \frac{1}{n-1} + \cdots + 1\Big) = nH_n,$$

where $H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}$ is the nth *harmonic number*. From Stage 2, $\ln n + \frac{1}{n} \le H_n \le \ln n + 1$, so H_n is about $\ln n$ when n is large, and I must expect to eat roughly

$$n \ln n \text{ packets.} \qquad \blacksquare$$

Looking back

- **Break a long wait into stages.** The total time split into n waiting times, each of a simple geometric kind.
- **Expectation is additive.** We never needed the distribution of the total, only the expectations of the pieces.
- **Compare sums with integrals.** The diagram turned a sum we cannot evaluate into a logarithm, with error at most 1.

What's really going on

How good is the estimate?

The comparison can be sharpened. The shaded gaps between the curve and the upper rectangles in Figure 9.11.1 have a finite total area, so $H_n - \ln n$ tends to a limit,

$$\gamma = \lim_{n\to\infty} (H_n - \ln n) = 0.5772156649\ldots,$$

Euler's constant. (Nobody knows whether γ is rational.) So the expected number of packets is $n \ln n + \gamma n + \frac{1}{2} + \cdots$. For the 600-sticker album, $600 \times (\ln 600 + 0.577) \approx 4185$, which is the figure quoted in the hook.

Where does the time go? The stages are very unequal. Collecting the first half of the set takes only about $n \ln 2 \approx 0.69n$ packets, while the last card alone takes n on average. Most of the effort goes into the last few cards, which is why collectors swap duplicates.

Concentration

The expected value is only the start: how much does the total vary? The variance of the total is $\sum nr/(n-r)^2$, which is about $\frac{\pi^2}{6}n^2$ (the sum $\sum 1/k^2 = \pi^2/6$ from Section 2.4 appears). So the standard deviation is about $1.28n$, much smaller than the mean $n \ln n$ when n is large: the total is *concentrated* around its mean. Paul Erdős and Alfréd Rényi showed in 1961 that, more precisely, the total minus $n \ln n$, divided by n, has a limiting distribution, the *Gumbel distribution*, which also describes the largest of many random quantities (Section 12.25).

Coupons everywhere

The problem goes back to Abraham de Moivre (1712) and Laplace. It now appears in many disguises. How many random tests are needed to check every component of a system? How long until a random walk on a network has visited every node? How many times must a computer hash table be probed? In each case the answer involves $n \ln n$. In biology, the same calculation estimates how many sequencing "reads" are needed to cover a whole genome.

The thinking skill

Linearity of expectation

The expectation of a sum is the sum of the expectations, whether or not the pieces are independent. Split a complicated random quantity into simple pieces.

This is one of the most useful facts in probability. Here it reduced the coupon collector's problem to n geometric waits; in Section 9.8 it gave the expected number of matches without any calculation. At university, the same idea powers the analysis of algorithms and the probabilistic method in combinatorics.

Try this next

1. **Half a set.** Show that the expected number of packets to collect $\frac{n}{2}$ different cards (for even n) is $n(H_n - H_{n/2})$, and that this is about $n \ln 2$.

2. **Distinct cards.** After opening k packets, show that the expected number of different cards you have is $n\big(1 - (1 - \frac{1}{n})^k\big)$. (*Hint:* for each card, find the probability that you have it, and add.) Roughly how many packets give you 95% of the set?

3. **Two collectors.** Two children collect the same set, sharing packets so that a duplicate for one may be new for the other. Show that the expected number of packets to complete both sets is about $n(\ln n + \ln \ln n)$. (This is harder; first convince yourself that it is much less than $2n \ln n$.)

4. **A sharper bound.** Use the trapezium rule picture instead of rectangles to show that $H_n - \ln n$ is decreasing, and that $H_n - \ln n - \frac{1}{2n}$ is increasing. Deduce $0.5 < \gamma < 0.7$ from small values of n.

Further reading

- **Geoffrey Grimmett & David Stirzaker, *One Thousand Exercises in Probability*** (Oxford University Press). Many variations on the coupon collector.

- **Julian Havil, *Gamma: Exploring Euler's Constant*** (Princeton University Press). Everything about γ and the harmonic series, at an accessible level.
- **Michael Mitzenmacher & Eli Upfal, *Probability and Computing*** (Cambridge University Press). The coupon collector and its uses in computer science.

9.12 Random Subsets

The question

STEP III 2019, Question 12

Read the question before you go on. It is in the STEP III paper in OCR's download of the 2019 STEP papers (a zip file):

`www.ocr.org.uk/Images/696590-2019-past-papers.zip`

The hook

Choose two subsets of $\{1, 2, \ldots, 100\}$ completely at random. How likely is it that they have no element in common? Your intuition may say "quite likely: most pairs of sets are different". In fact the probability is $(\frac{3}{4})^{100}$, about 3×10^{-13}. Random sets are big (about half of everything) and so they almost always overlap.

The question is short, and its one idea is powerful: do not think about the sets as a whole; think about one element at a time. A random subset is just n independent coin tosses, one for each element, and every question about random subsets becomes a question about those tosses.

Before you start

Ask yourself How can you describe a subset of $\{1, \ldots, n\}$ by a sequence of n answers to yes/no questions?

For each element i, answer the question "is i in the set?". The subset determines the answers, and the answers determine the subset. There are 2 choices for each element, so 2^n subsets in all. Choosing a subset uniformly at random is the same as answering each question by tossing a fair coin, independently.

Working it through

STAGE 1 One element

By the coin-tossing description, $\mathrm{P}(1 \in A_1) = \frac{1}{2}$. Indeed, exactly half of the 2^n subsets contain 1: pair each subset that does not contain 1 with the one obtained by adding 1.

STAGE 2 Disjoint sets

Ask yourself For $A_1 \cap A_2 = \varnothing$, what must be true of each element i? What is the probability of that, for one element? Why can you multiply over the elements?

A_1 and A_2 are disjoint exactly when no element is in both. For a single element i, there are four equally likely possibilities: in neither set, in A_1 only, in A_2 only, in both (Figure 9.12.1, left). Only the last is forbidden, so the probability that i is not in both is $\frac{3}{4}$. Different elements are decided by different, independent coin tosses, so

$$\mathrm{P}(A_1 \cap A_2 = \varnothing) = \left(\frac{3}{4}\right)^n. \qquad \blacksquare$$

For three sets, element i is in all three with probability $\frac{1}{8}$, so $\mathrm{P}(A_1 \cap A_2 \cap A_3 = \varnothing) = \left(\frac{7}{8}\right)^n$. In general,

$$\mathrm{P}(A_1 \cap \cdots \cap A_m = \varnothing) = \left(1 - \frac{1}{2^m}\right)^n. \qquad \blacksquare$$

STAGE 3 Subsets

Ask yourself For $A_1 \subseteq A_2$, which of the four possibilities for an element is forbidden?

$A_1 \subseteq A_2$ fails exactly when some element is in A_1 but not in A_2. Again one pattern out of four is forbidden, so

$$\mathrm{P}(A_1 \subseteq A_2) = \left(\frac{3}{4}\right)^n,$$


the same as for disjointness. That is not a coincidence: $A_1 \subseteq A_2$ exactly when A_1 is disjoint from the complement of A_2, and the complement of a random set is another random set.

Ask yourself For $A_1 \subseteq A_2 \subseteq A_3$, which of the 8 patterns of membership (in or out of each set) are allowed for an element?


Going along the chain A_1, A_2, A_3, once an element is in, it must stay in. So the allowed patterns are "out, out, out", "out, out, in", "out, in, in" and "in, in, in": 4 out of 8 (Figure 9.12.1, right). Hence $\mathrm{P}(A_1 \subseteq A_2 \subseteq A_3) = \left(\frac{4}{8}\right)^n = \left(\frac{1}{2}\right)^n$.

For a chain of m sets, an allowed pattern is determined by the first set the element enters (one of $A_1, \ldots, A_m$, or never), so $m+1$ of the 2^m patterns are allowed:

$$\mathrm{P}(A_1 \subseteq A_2 \subseteq \cdots \subseteq A_m) = \left(\frac{m+1}{2^m}\right)^n. \qquad \blacksquare$$



each element: 4 equally likely places



allowed for $A_1 \subseteq A_2 \subseteq A_3$: 4 of 8

Figure 9.12.1 Left: an element of S can be in one of four places relative to A_1 and A_2, each with probability $\frac{1}{4}$. Disjointness forbids "both" (amber); $A_1 \subseteq A_2$ forbids "only A_1". Right: the eight patterns of membership of A_1, A_2, A_3 (filled circle means "in"). The chain $A_1 \subseteq A_2 \subseteq A_3$ allows only the four boxed patterns, in which an element, once in, stays in.

Looking back

- **Decompose by elements.** Each set-level condition (disjoint, subset, chain) was a condition on each element separately.
- **Independence across elements.** Because the elements are decided by independent coin tosses, the probabilities multiply.
- **Count patterns.** Each answer was (allowed patterns ÷ all patterns)^n.

What's really going on

Counting without probability

Multiplying by the number of choices turns each probability into a count. The number of ordered pairs (A_1, A_2) with $A_1 \subseteq A_2$ is $4^n \times \left(\frac{3}{4}\right)^n = 3^n$. You can also count these pairs directly: choose the size k of A_2, then A_2 itself in $\binom{n}{k}$ ways, then A_1 as any of its 2^k subsets. So

$$\sum_{k=0}^{n} \binom{n}{k} 2^k = 3^n,$$

which is the binomial theorem for $(1+2)^n$. Counting the same thing in two ways, once by elements and once by sizes, proves an identity: this is the method of *double counting*, which you met with the Platonic solids in Section 7.9. Similarly, the number of chains $A_1 \subseteq \cdots \subseteq A_m$ is $(m+1)^n$.

The lattice of subsets

The subsets of $\{1, \ldots, n\}$, ordered by inclusion, form a structure called the *Boolean lattice*, which can be pictured as an n-dimensional cube: each subset is a corner, labelled by its sequence of yes/no answers, and moving along an edge adds or removes one element. Chains are paths going upwards. Deep results describe how large families of subsets

can be, given conditions on them. *Sperner's theorem* (1928) says that the largest family in which no set contains another has $\binom{n}{\lfloor n/2 \rfloor}$ sets; the *Erdős–Ko–Rado theorem* (1961) says how many k-element sets can be chosen so that every two overlap. Both are standard in a first course on combinatorics, and the coin-tossing view of subsets gives some of the slickest proofs.

Boole's algebra

George Boole's *The Laws of Thought* (1854) treated logical statements as sets and "and", "or" and "not" as intersection, union and complement. Encoding a set as a string of yes/no answers is exactly how a computer represents it: a subset of $\{1, \dots, 64\}$ is a single 64-bit number, and intersection is the "bitwise AND" instruction. The method of this question, one element (one bit) at a time, is how such operations are analysed.

The thinking skill

Think one element at a time

When objects are built from independent choices for each element, a condition that applies to every element separately has probability equal to a product.

The method turned a question about 2^n subsets into a question about 4 or 8 patterns. At university it reappears in combinatorics (generating functions for sets and multisets), in probability (independent coordinates) and in computer science (bitwise reasoning).

Try this next

1. **Covering.** Find $\mathrm{P}(A_1 \cup A_2 = S)$ and $\mathrm{P}(A_1 \cup A_2 \cup \cdots \cup A_m = S)$.
2. **Same size.** Show that $\mathrm{P}(|A_1| = |A_2|) = \binom{2n}{n}/4^n$. (*Hint:* $|A_1| = |A_2|$ exactly when $|A_1| + |S \setminus A_2| = n$.) Show that this is about $1/\sqrt{\pi n}$ for large n.
3. **Partitions into three.** Show that the number of ordered pairs of disjoint subsets (A, B) of S is 3^n, and find the number of unordered pairs $\{A, B\}$ of disjoint subsets.
4. **Sperner for small n.** For $n = 3$ and $n = 4$, find the largest family of subsets of $\{1, \dots, n\}$ in which no set contains another.

Further reading

- **Miklós Bóna, *A Walk Through Combinatorics*** (World Scientific), the chapters on subsets and partially ordered sets.

- **Béla Bollobás,** ***Combinatorics: Set Systems, Hypergraphs, Families of Vectors and Combinatorial Probability*** (Cambridge University Press). A short, elegant book on families of subsets, from Sperner's theorem onwards.
- **Peter Cameron,** ***Combinatorics: Topics, Techniques, Algorithms*** (Cambridge University Press). A first course with a strong emphasis on counting in two ways.

9.13 Loaded Dice

The question

STEP I 1999, Question 12

Read the question before you go on. It is in the STEP Support Programme's question database:

`step.maths.org/questions/99-s1-q12`

The hook

With two fair dice, a total of 7 is six times as likely as a total of 2. Board-game designers sometimes wish it were otherwise: could you load the dice (weight them so that some faces come up more often) so that every total from 2 to 12 has the same chance, $\frac{1}{11}$? You can choose the twelve probabilities freely, so there seems to be plenty of room.

The answer is no, and the proof is a gem: it uses only three of the eleven totals and the inequality $x + \frac{1}{x} \geq 2$. It is a model of how to prove that something is *impossible*, which is often harder, and more interesting, than showing that something can be done.

Before you start

Ask yourself How would you prove $x + \frac{1}{x} \geq 2$ for $x > 0$? When is there equality?

Multiply by $x > 0$: the inequality becomes $x^2 + 1 \geq 2x$, that is $(x-1)^2 \geq 0$, which is true. Equality holds only when $x = 1$. (Alternatively, $x + \frac{1}{x} - 2 = (\sqrt{x} - \frac{1}{\sqrt{x}})^2$.) This is the AM–GM inequality in its simplest form: the average of x and $\frac{1}{x}$ is at least their geometric mean, 1.

Working it through

STAGE 1 The extreme totals

Ask yourself In how many ways can you throw a total of 2? A total of 12? Write their probabilities in terms of the q_i and r_j.

A total of 2 needs 1 on both dice, and 12 needs 6 on both, so $\mathrm{P}(2) = q_1 r_1$ and $\mathrm{P}(12) = q_6 r_6$.

Suppose, for contradiction, that all eleven totals are equally likely, each with probability $\frac{1}{11}$. Then

$$q_1 r_1 = \frac{1}{11}, \qquad q_6 r_6 = \frac{1}{11}.$$

In particular, all four of q_1, r_1, q_6, r_6 are positive.

STAGE 2 The middle total

Ask yourself A total of 7 can be made in six ways. Which two of them involve only the faces 1 and 6? Write their probabilities using the equations above.

The total 7 includes the outcomes (1, 6) and (6, 1), so

$$\mathrm{P}(7) \geq q_1 r_6 + q_6 r_1.$$

Now use the equations of Stage 1 to bring in a ratio. Since $q_1 = \frac{1}{11 r_1}$ and $q_6 = \frac{1}{11 r_6}$,

$$q_1 r_6 + q_6 r_1 = \frac{1}{11}\left(\frac{r_6}{r_1} + \frac{r_1}{r_6}\right) \geq \frac{1}{11} \times 2 = \frac{2}{11},$$

using $x + \frac{1}{x} \geq 2$ with $x = r_6/r_1 > 0$. So $\mathrm{P}(7) \geq \frac{2}{11}$, contradicting $\mathrm{P}(7) = \frac{1}{11}$. The totals cannot all be equally likely. ■

Pause and try Look at what made the proof work. Making 2 and 12 as likely as 7 forces the 1s and 6s to be fairly likely, and then the "mixed" outcomes (1, 6) and (6, 1) alone make 7 too likely. Figure 9.13.1 shows the cells involved.




Figure 9.13.1 The 36 outcomes of two dice, labelled by their totals. The proof uses only the four corner cells. The product of the two amber probabilities equals the product of the two dark teal ones, $q_1r_1q_6r_6 = q_1r_6 \cdot q_6r_1$, and by AM–GM the sum of the teal pair is at least twice the geometric mean of the amber pair.

STAGE 3 Two throws of one die

Ask yourself If one die is thrown twice, what is the probability that the two numbers are the same, in terms of the q_j? What is it for a fair die?

The two throws agree with probability $q_1^2 + q_2^2 + \cdots + q_6^2$. For a fair die this is $6 \times \frac{1}{36} = \frac{1}{6}$, which is already much bigger than $\frac{1}{36}$.

Ask yourself Can loading make $\sum q_j^2$ smaller than $\frac{1}{6}$? You know that $\sum q_j = 1$.

Write $q_j = \frac{1}{6} + e_j$, where the deviations e_j add to zero. Then

$$\sum q_j^2 = \sum\Big(\frac{1}{36} + \frac{e_j}{3} + e_j^2\Big) = \frac{1}{6} + \frac{1}{3}\sum e_j + \sum e_j^2 = \frac{1}{6} + \sum e_j^2 \geq \frac{1}{6}.$$

So the probability of a repeat is always at least $\frac{1}{6}$, with equality only for a fair die. It certainly cannot be less than $\frac{1}{36}$. ■

Looking back

- **Proof by contradiction.** We assumed all totals equally likely and derived an impossibility.
- **Use only what you need.** Of the eleven equations, three were enough. Choosing the right ones was the key step.
- **Inequalities prove impossibility.** AM–GM in part (i), and "a sum of squares is non-negative" in part (ii), ruled out whole families of possibilities at once.

What's really going on

A proof with polynomials

Here is a completely different proof, using the generating functions of Section 9.6. Let $Q(x) = q_1x + \cdots + q_6x^6$ and $R(x) = r_1x + \cdots + r_6x^6$. If all totals had probability $\frac{1}{11}$, then

$$Q(x)R(x) = \frac{x^2 + x^3 + \cdots + x^{12}}{11} = \frac{x^2}{11}(1 + x + \cdots + x^{10}).$$

Now $Q(x)/x$ is a polynomial of degree 5 with real coefficients, and every real polynomial of odd degree has a real root (its graph goes from $-\infty$ to $+\infty$, or the other way). So the left side has a real non-zero root. But $1 + x + \cdots + x^{10} = \frac{x^{11}-1}{x-1}$, whose roots are the 11th roots of unity other than 1, none of which is real. Contradiction. The same argument shows that no loading of two n-sided dice can make the totals equally likely whenever $n - 1$ is odd, that is, for n even. (For odd n the polynomial argument needs more work, but the AM–GM proof above works for every $n \geq 2$.)

Collision probability

The quantity $\sum q_j^2$ in part (ii), the chance that two independent draws agree, is called the *collision probability* of a distribution. Part (ii) shows that it is smallest for the uniform distribution. It measures how concentrated a distribution is: in ecology, $\sum q_j^2$ is Simpson's index of diversity, where q_j is the proportion of species j in a habitat; in economics, it is the Herfindahl index of market concentration; in cryptography, it measures how guessable a password distribution is. And it explains the remark at the end of Section 9.9: unequal birthday probabilities can only make shared birthdays more likely.

Impossibility proofs

Some of the most famous results in mathematics say that something cannot be done: no fraction squares to 2; no ruler-and-compass construction trisects every angle; no formula in radicals solves every quintic; no algorithm decides every mathematical statement. Each changed its subject, because to prove that *no* method works, you must understand the whole space of methods. The loaded dice problem is a miniature: to rule out every loading, we needed an invariant property (the AM–GM inequality) that every loading must obey.

The thinking skill

Prove impossibility by contradiction and inequality

To show that no choice of parameters works, assume one does, pick out the few conditions that clash, and use an inequality to show they cannot all hold.

At university, contradiction combined with inequalities is the standard way to prove that objects do not exist, in analysis, in combinatorics and in optimisation, where inequalities like AM–GM and Cauchy–Schwarz set limits on what can be achieved.

Try this next

1 **Cauchy–Schwarz.** Use the Cauchy–Schwarz inequality, $\left(\sum a_j b_j\right)^2 \le \sum a_j^2 \sum b_j^2$, with $a_j = q_j$ and $b_j = 1$, to give a one-line proof of part (ii).

2 **Two different dice.** If two *different* loaded dice are each thrown once, can the probability that they show the same number be less than $\frac{1}{6}$? Can it be 0?

3 **Uniform totals with other faces.** Show that two dice *can* give equally likely totals if their faces are numbered differently: try one die numbered $0, 1, 2, 3, 4, 5$ and the other $0, 6, 12, 18, 24, 30$. What totals arise? Why does this not contradict the question?

4 **Three dice.** Can three loaded six-sided dice, with faces 1 to 6, make all totals from 3 to 18 equally likely? (*Hint:* use the totals 3, 8, 13 and 18, and AM–GM for six numbers. Why does the polynomial proof fail this time?)

Further reading

- **J. Michael Steele, *The Cauchy–Schwarz Master Class*** (Cambridge University Press). A beautiful introduction to inequalities, starting from AM–GM.
- **Geoffrey Grimmett & Dominic Welsh, *Probability: An Introduction*** (Oxford University Press). Generating functions and their use in proofs like the one above.
- **Ian Stewart, *Galois Theory*** (CRC Press), the opening chapters. The most famous impossibility proof of all, for the quintic.

What We Have Learned

We began by saying that counting is the oldest mathematics there is, and that doing it cleverly is still one of the hardest. We asked three questions: how do you count without listing, when must something happen, and how do counting and probability help each other? Here is what the chapter's thirteen questions showed.

1. How do you count without listing?

The chapter built up a toolkit.

- **Recurrences.** Adding one line to the plane (Section 9.1) or one post to a fence (Section 9.3) changed the count in a simple way, giving a recurrence that could be solved. Recording a little extra information, the colour of the last post, made the recurrences close up.
- **Generating functions.** Packaging a sequence as a power series turned recurrences into algebra (Section 9.4), explained why $1/89 = 0.01123\ldots$, and turned counting combinations into multiplying polynomials: coins in Section 9.5, dice in Section 9.6, and balanced power sums in Section 9.7, where a repeated root at $x = 1$ did all the work.
- **Bijections.** The reflection principle (Section 9.10) counted bad queues by matching them with unrestricted paths, and complements of random sets turned subsets into disjoint pairs (Section 9.12).
- **Inclusion–exclusion.** Overcounting and correcting found the number of derangements (Section 9.8).
- **Double counting.** Counting regions with Euler's formula (Section 9.1) and pairs of subsets by sizes (Section 9.12) gave the same answers by different routes, and proved identities on the way.

2. When must something happen?

Several sections showed that size alone forces structure. Among six people there must be three mutual friends or three mutual strangers (Section 9.2), by the pigeonhole principle, though five people are not enough. Twenty-three people make a shared birthday more likely than not (Section 9.9), because the number of pairs grows like the square of the number of people. Random subsets of a large set almost always overlap (Section 9.12). And some things *cannot* happen, however you try: no loading of two dice makes every total equally likely (Section 9.13), and no numbering of dice other than Sicherman's reproduces the ordinary totals (Section 9.6).

3. How do counting and probability help each other?

In the second half of the chapter, probabilities were found by counting: derangements by inclusion–exclusion, the birthday problem by a product, the ballot problem by reflection. But the traffic went the other way too. Linearity of expectation gave the coupon collector's nH_n (Section 9.11) and the expected number of matches in a shuffle without any counting at all. Thinking of a random subset as n coin tosses (Section 9.12) turned counts like 3^n and $(m+1)^n$ into one-line calculations. And Erdős's probabilistic method (Section 9.2) proved that good colourings exist by showing that a random colouring works with positive probability. Two constants appeared from nowhere: e, in the limiting chance of a derangement, and Euler's γ, in the cost of completing a collection.

4. A toolkit for counting

Method	Where you used it
What changes when you add one	Lines (9.1); fence posts (9.3)
Pigeonhole principle	Ramsey (9.2)
Coupled recurrences	Fence posts (9.3)
Generating functions	9.4, 9.5, 9.6, 9.7, 9.13
Case analysis from the extremes	Strange dice (9.6); loaded dice (9.13)
Inclusion–exclusion	Derangements (9.8)
Complement and exponential approximation	Birthdays (9.9)
Reflection principle	Ballot problem (9.10)
Linearity of expectation	Coupons (9.11); matches (9.8)
One element at a time	Random subsets (9.12)
Contradiction with AM–GM	Loaded dice (9.13)

Where next

- The Poisson limit of derangements returns in Section 12.11, and generating functions of random variables drive the study of family trees in Section 12.15.
- Coupled recurrences become Markov chains in Section 12.16.
- The Gumbel distribution of the coupon collector describes the largest of many random quantities in Section 12.25.

If you want to go further before university

- **Miklós Bóna, *A Walk Through Combinatorics***: a friendly first course covering almost every idea in this chapter.
- **Herbert Wilf, *generatingfunctionology***: the classic on generating functions; free online.

- **Ronald Graham, Donald Knuth & Oren Patashnik, *Concrete Mathematics***: sums, recurrences and generating functions, with wit and many exercises.
- **Geoffrey Grimmett & Dominic Welsh, *Probability: An Introduction***: the counting side of first-year probability.

Whitehead called combinatorics "the slums of topology". This chapter has tried to show why nobody would say that now. Behind every clever count there was an idea that reaches far beyond it: a generating function, a bijection, a recurrence, an inequality. Those ideas connect counting with algebra, analysis and probability, and learning to look for them, rather than for one more trick, is what turns counting into mathematics.

Answers to Try This Next

These are brief answers and hints, not full solutions: they give the result and the key step, so that you can check your own work. Try each exercise properly before looking.

9.1 Lines Cutting the Plane

1. One circle gives 2 regions. The nth circle meets the others in $2(n-1)$ points, which cut it into $2(n-1)$ arcs, each splitting a region in two. So $R_n = R_{n-1} + 2(n-1)$, giving $n^2 - n + 2$.

2. Each line adds 1 plus the number of crossings on it, so the total is $1 + n +$ (number of crossings). The k parallel lines lose $\binom{k}{2}$ crossings: $1 + n + \binom{n}{2} - \binom{k}{2}$.

3. The nth plane is cut by the others in $n-1$ lines in general position, which divide it into $\frac{1}{2}(n^2 - n + 2)$ pieces, each splitting a region of space. Adding these to $\frac{1}{6}\big((n-1)^3 + 5(n-1) + 6\big)$ gives $\frac{1}{6}(n^3 + 5n + 6)$.

4. For $n = 6$ there are 31 regions, not 32. Inside the circle the graph has $V = n + \binom{n}{4}$ vertices (each crossing comes from four points) and $E = n + \binom{n}{2} + 2\binom{n}{4}$ edges (the arcs of the circle, plus the chords, each crossing adding two pieces). Euler's formula $V - E + F = 2$ counts the outside too, so the number of regions is $F - 1 = 1 + \binom{n}{2} + \binom{n}{4}$.

9.2 Ramsey's Theorem in Miniature

1. There are at most $6 \times 6 = 36$ bichromatic angles. Each triangle that is not one-colour contains exactly two, so there are at most 18 such triangles. Since $\binom{6}{3} = 20$, at least 2 triangles are one-colour.

2. A point has 16 edges, so at least 6 of one colour, say red, going to points $P_1, \ldots, P_6$. If any edge among the P_i is red, there is a red triangle. If not, the P_i form a two-coloured K_6, which has a one-colour triangle.

3. A point v has 9 edges, so either at least 4 are gold or at least 6 are silver. With 4 gold neighbours: a gold edge among them gives a gold triangle with v, and otherwise they form a silver K_4. With 6 silver neighbours: among them there is a one-colour triangle; if gold we are done, and if silver it forms a silver K_4 with v.

4. Label each term with (i, d): the lengths of the longest increasing and decreasing subsequences ending there. For two terms, the later one has a larger i or a larger d, so the labels are all different. If no subsequence of length 3 existed, there would be only 4 possible labels for 5 terms. Example with neither: $2, 1, 4, 3$.

9.3 Colouring Fence Posts

1. $r_{n+1} = s_n$ as before (post n is not red, so post $n+1$ can be red), and $r_n + s_n = (k-1)^{n-1}$, since each post after the first has $k-1$ choices. So $r_{n+1} + r_n = (k-1)^{n-1}$, and $r_n = \frac{(k-1)^{n-1} + (k-1)(-1)^{n-1}}{k}$. For the circle: k choices of colour for post 1, then a line of $n+1$ posts ending in the same colour, which gives $kr_{n+1} = (k-1)^n + (-1)^n(k-1)$.

2. $2, 3, 5, 8, \ldots$: with a_n ways, a line ending in white continues any line of $n-1$, and one ending in red continues a line of $n-2$ ending in white, so $a_n = a_{n-1} + a_{n-2}$. The answer is F_{n+2}.

3. 4 choices for the centre, then a 3-colouring of the n-cycle: $4\big(2^n + 2(-1)^n\big)$.

4. For $n = 4$: 18 colourings. A quarter-turn fixes none, and a half-turn fixes the 6 with opposite posts alike, so there are $\frac{1}{4}(18 + 0 + 6 + 0) = 6$ necklaces. For $n = 5$: 30 colourings, no rotation fixes any, so 6.

9.4 Sequences as Functions

1. The tail is at most $F_7 \sum_{r \geq 7} 2^{r-7} 10^{-r} = \frac{13 \times 10^{-7}}{0.8} < 2 \times 10^{-6}$. Since $\sum F_r 10^{-r} = \frac{10}{89}$ and the first six terms give 0.112358, $\frac{10}{89}$ lies between 0.112358 and 0.112360, so $\frac{1}{89} = 0.011235\ldots$. With two-digit blocks, $\sum F_r 100^{-r-1} = \frac{1}{9899}$.

2. Summing the recurrence gives $f(x)(1 - 7x + 10x^2) = 2 - 7x$, and $\frac{2-7x}{(1-2x)(1-5x)}$ splits as stated. For b_n: $10p + 5q = 40$ and $40p + 10q = 100$ give $p = 1$, $q = 6$; the roots of $t^2 = t + 6$ are 3 and -2, and $b_n = 4 \cdot 3^n + (-2)^n$.

3. $\sum n x^n = \frac{x}{(1-x)^2}$; at $x = \frac{1}{100}$ this is $\frac{100}{9801}$, so $\frac{1}{9801} = \sum n \cdot 100^{-n-1}$. The pattern goes wrong at the block 98: the block for 100 carries 1 into the 99, which carries into the 98, so the expansion runs $\ldots 96\,97\,99\,00\,01\ldots$.

4. Multiplying by $\frac{1}{1-x} = 1 + x + x^2 + \cdots$ makes the coefficient of x^n equal to $u_0 + \cdots + u_n$. The squares have generating function $\frac{x(1+x)}{(1-x)^3}$, so the partial sums have $\frac{x+x^2}{(1-x)^4}$, and the coefficient of x^n is $\binom{n+2}{3} + \binom{n+1}{3} = \frac{n(n+1)(2n+1)}{6}$.

9.5 Making Change

1. The coefficient of x^{24} counts $6a + 6b + 3c = 24$: for each $s = a + b$ from 0 to 4 there are $s + 1$ choices, giving 15. In general the coefficient of x^n is 0 unless $n = 3m$, and then it is $\frac{1}{2}(k+1)(k+2)$ with $k = \lfloor m/2 \rfloor$. With the extra factor the coefficients are partial sums: 55, 55 and 650 for x^{24}, x^{25} and x^{66}. They count the ways of paying with 1p and 3p coins and two different kinds of 6p coin.

2. Each part k is used once or not at all. The product telescopes: $\prod \frac{1-x^{2k}}{1-x^k}$ cancels every factor with an even exponent from the denominator, leaving $\prod \frac{1}{1-x^{\text{odd}}}$, the generating function for partitions into odd parts. For $n = 6$: 6, $5+1$, $4+2$, $3+2+1$, and $5+1$, $3+3$, $3+1+1+1$, $1+1+1+1+1+1$: four of each.

3. The values for $n = 0, 1, 2, \ldots$ are $1, 1, 2, 2, 3, 4, 5, 6, 7, 8, 10, \ldots$, and $\frac{(n+4)^2}{20}$ is always within $\frac{1}{2}$ of them (checked up to $n = 60$; a proof uses the partial fractions of $\frac{1}{(1-x)(1-x^2)(1-x^5)}$).
4. The last coin is 1p or 2p, so $c_n = c_{n-1} + c_{n-2}$: the Fibonacci numbers, $c_n = F_{n+1}$.

9.6 Strange Dice

1. Each die is x times a product of the factors $1 + x$, $1 + x$, $1 + x^2$, $1 + x^2$ with value 4 at $x = 1$, so it takes two of them. Apart from the standard pair, the only option is $x(1 + x)^2$ and $x(1 + x^2)^2$: faces $1, 2, 2, 3$ and $1, 3, 3, 5$.
2. The product is $2x^3 + x^4 + x^5 + 4x^6 + 3x^7 + 8x^8 + 6x^9 + 5x^{10} + 6x^{11}$, matching the table.
3. Yes: the Sicherman pair $1, 2, 2, 3, 3, 4$ and $1, 3, 4, 5, 6, 8$ together with a standard die.
4. $(x + x^2)(x + \cdots + x^8) = x^2(1 + x)^2(1 + x^2)(1 + x^4)$. The coin needs a factor with value 2: besides the standard choice $x(1 + x)$, it can be $x(1 + x^2)$ (faces $1, 3$, with the octahedron $1, 2, 2, 3, 5, 6, 6, 7$) or $x(1 + x^4)$ (faces $1, 5$, with $1, 2, 2, 3, 3, 4, 4, 5$).

9.7 Equal Sums of Powers

1. $\sum(c + b_k)^3 - \sum(c + a_k)^3 = 3c^2\big(\sum b_k - \sum a_k\big) + 3c\big(\sum b_k^2 - \sum a_k^2\big) + \sum b_k^3 - \sum a_k^3$, and the first two brackets vanish. Starting from $\{1, 4, 6, 7\}$, $\{2, 3, 5, 8\}$ with $c = 8$: $\{1, 4, 6, 7, 10, 11, 13, 16\}$ and $\{2, 3, 5, 8, 9, 12, 14, 15\}$.
2. A and $B \cup \{0\}$ both have 6 elements, sum 33 and sum of squares 253. Then $\sum_{a \in A}(n + a)^2 = \sum_{b \in B \cup \{0\}}(n + b)^2$, which splits $\{n^2, \ldots, (n + 11)^2\}$.
3. Expand $(x + c)^r$ by the binomial theorem: it involves only powers up to r, whose sums agree. The question's sets, with 0 added to the second, are $\{0, \ldots, 15\}$, and translating gives any 16 consecutive integers.
4. With $ABABABAB$, A gets about 1.676 and B about 1.042. With $ABBABAAB$, A gets about 1.560 and B about 1.158: still unequal, but much closer.

9.8 Nobody Gets Their Own Shirt

1. $\mathrm{E}(C_1) = \frac{1}{m}$, $\mathrm{Var}(C_1) = \frac{1}{m} - \frac{1}{m^2}$, and $\mathrm{Cov}(C_1, C_2) = \frac{1}{m(m-1)} - \frac{1}{m^2} = \frac{1}{m^2(m-1)}$. Then $\mathrm{E}(N) = 1$ and $\mathrm{Var}(N) = m\big(\frac{1}{m} - \frac{1}{m^2}\big) + m(m-1)\frac{1}{m^2(m-1)} = 1$. For $m = 4$: $\mathrm{P}(N = 0) = \frac{9}{24} = 0.375$, against $\mathrm{e}^{-1} \approx 0.368$.
2. Card 1 goes to some position i ($n - 1$ choices). If card i goes to position 1, the rest is a derangement of $n - 2$; if not, treat position 1 as card i's forbidden place, giving a derangement of $n - 1$. Check: $D_3 = 2(1 + 0) = 2$ and $D_4 = 3(2 + 1) = 9$.
3. Choose the k fixed cards and derange the rest: $\binom{n}{k}\frac{D_{n-k}}{n!} = \frac{1}{k!} \cdot \frac{D_{n-k}}{(n-k)!} = \frac{d_{n-k}}{k!}$, which tends to $\frac{\mathrm{e}^{-1}}{k!}$.

4. $30-(15+10+6)+(5+3+2)-1=8$, which equals $30\cdot\frac{1}{2}\cdot\frac{2}{3}\cdot\frac{4}{5}$: the inclusion–exclusion sum factorises (this is Euler's $\varphi(30)$).

9.9 The Birthday Problem

1. $\mathrm{P}(X=r)=\bigl(1-\frac{1}{n}\bigr)\cdots\bigl(1-\frac{r-2}{n}\bigr)\frac{r-1}{n}$, and these sum to 1. For the expectation, $\mathrm{P}(Y=r)$ is counted once in each of $\mathrm{P}(Y\ge 1),\dots,\mathrm{P}(Y\ge r)$. Write $Q_k=\mathrm{P}(X>k)=\prod_{j=1}^{k-1}(1-\frac{j}{n})$. Then $\mathrm{E}(X)=\sum(k+1)\frac{k}{n}Q_k$ and also $\mathrm{E}(X)=1+\sum Q_k$; using $\sum\frac{k}{n}Q_k=1$ and subtracting gives $\sum Q_k\bigl(1-\frac{k^2}{n}\bigr)=0$.

2. $\binom{k}{3}\approx 365^2\ln 2$ gives $k\approx 82$; the approximation ignores the dependence between triples, and the true answer is 88.

3. 14 people. (Each pair is "close" with probability $\frac{3}{365}$, and $\binom{k}{2}\frac{3}{365}\approx\ln 2$ gives $k\approx 14$; an exact count gives a probability of about 0.537 for 14 and 0.483 for 13.)

4. Two people share day i with probability p_i^2; add. By Cauchy–Schwarz, $1=\bigl(\sum p_i\bigr)^2\le 365\sum p_i^2$, with equality only when all p_i are equal.

9.10 Always Able to Give Change

1. The first person must have a £1 coin. Conditioning on the next few people splits into many cases (which coins you hold after each £2 customer), each finishing with the answers to parts (i) and (ii); collecting them gives $\frac{m-2}{m+1}$. The reflection principle does it in one line.

2. The first vote must be for A. Among the counts that start with A, the bad ones (those that reach a tie) correspond, by reflecting the part before the first tie, to the counts that start with B. So the number of good counts is $\binom{a+b-1}{a-1}-\binom{a+b-1}{a}$, and dividing by $\binom{a+b}{a}$ gives $\frac{a-b}{a+b}$.

3. $((()))$, $(()())$, $(())()$, $()(())$, $()()()$. A bijection: the first bracket pair splits a sequence into the part inside it and the part after it; correspondingly the triangle on a fixed edge splits the polygon into two smaller polygons. Recurse on both.

4. Each £5 customer needs four £1 coins in change, so the path goes up 1 or down 4. A path can jump past the barrier instead of touching it, and reflection no longer matches the bad paths with a simple set.

9.11 Collecting the Set

1. The expected waits are $\frac{n}{n}+\frac{n}{n-1}+\cdots+\frac{n}{n/2+1}=n(H_n-H_{n/2})\approx n(\ln n-\ln\frac{n}{2})=n\ln 2$.

2. Each card is missing with probability $(1-\frac{1}{n})^k$; add over the n cards. For 95%, $(1-\frac{1}{n})^k\approx \mathrm{e}^{-k/n}=0.05$, so $k\approx n\ln 20\approx 3n$.

3. After t packets a given card has been seen fewer than twice with probability about $\mathrm{e}^{-t/n}(1+\frac{t}{n})$. Both children finish roughly when the expected number of such

cards, $ne^{-t/n}(1+\frac{t}{n})$, falls to about 1; that gives $\frac{t}{n} \approx \ln n + \ln\ln n$, far less than $2\ln n$.

4. $\frac{1}{n+1} < \ln\frac{n+1}{n}$ (area under $\frac{1}{x}$ on $[n, n+1]$), so $H_n - \ln n$ decreases. By convexity the trapezium overestimates the area: $\ln\frac{n+1}{n} < \frac{1}{2}\left(\frac{1}{n} + \frac{1}{n+1}\right)$, so $H_n - \ln n - \frac{1}{2n}$ increases. With $n = 1$, $\gamma > \frac{1}{2}$; with $n = 4$, $\gamma < H_4 - \ln 4 \approx 0.697$.

9.12 Random Subsets

1. Each element must be in at least one set: $\left(\frac{3}{4}\right)^n$, and $\left(1 - 2^{-m}\right)^n$ in general.
2. $|S \setminus A_2|$ has the same distribution as $|A_2|$, and is independent of A_1, so the probability is that of $|A_1| + |S \setminus A_2| = n$. The pair $(A_1, S \setminus A_2)$ corresponds to a random subset of a $2n$-element set, of size n with probability $\binom{2n}{n}/4^n$. Stirling (or Wallis, Section 4.4) gives $\approx \frac{1}{\sqrt{\pi n}}$.
3. Each element goes in A, in B or in neither: 3^n. The only ordered pair with $A = B$ is $(\varnothing, \varnothing)$, so there are $\frac{3^n-1}{2} + 1 = \frac{3^n+1}{2}$ unordered pairs.
4. For $n = 3$, 3 sets (all the singletons, or all the pairs). For $n = 4$, 6 sets (all the pairs). In general $\binom{n}{\lfloor n/2 \rfloor}$ (Sperner's theorem).

9.13 Loaded Dice

1. $1 = \left(\sum q_j\right)^2 \le 6\sum q_j^2$, so $\mathrm{P}(\text{same}) = \sum q_j^2 \ge \frac{1}{6} > \frac{1}{36}$.
2. Yes, and yes: the probability is $\sum q_j r_j$, which is 0 if, say, one die always shows 1 and the other always 2.
3. The totals are $0, 1, \ldots, 35$, each exactly once: uniform. The question requires both dice to be numbered 1 to 6, and here they are not.
4. No. Write $\mathrm{P}(3) = q_1 r_1 s_1$ and $\mathrm{P}(18) = q_6 r_6 s_6$, both $\frac{1}{16}$. $\mathrm{P}(8) + \mathrm{P}(13)$ includes the six terms $q_1 r_1 s_6, \ldots, q_1 r_6 s_6$ (two 1s and a 6, or two 6s and a 1), whose product is $(\mathrm{P}(3)\mathrm{P}(18))^3$. AM–GM gives $\mathrm{P}(8) + \mathrm{P}(13) \ge 6 \cdot \frac{1}{16} > \frac{2}{16}$. The polynomial proof as given fails because $1 + x + \cdots + x^{15}$ *does* have a real root, -1. (It can be rescued: three real quintics need three real roots, and -1 is only a simple root.)