11.3 The Real Pendulum

WORKSHEET

The question

STEP III 1998, Question 11

Read the question before you go on. It is in the STEP Support Programme's question database:

step.maths.org/questions/98-s3-q11

Hints

Try the question on your own first. If you get stuck, read the hints *one at a time*, and go back to the question after each one. Hints marked *Pause and try* are the key steps.

Ask yourself The bob moves on a circle of radius l. What are its speed and its height (relative to the lowest point) in terms of θ and $\dot{\theta}$?

STAGE 2 The period as an integral

Ask yourself How can you write $\cos\theta - \cos\gamma$ in terms of half-angles? And how much of the period is spent going from $\theta = 0$ to $\theta = \gamma$?

STAGE 3 Removing the singularity

Ask yourself The integrand becomes infinite at $\theta = \gamma$. Write $k = \sin\frac{\gamma}{2}$ and substitute $\sin\frac{\theta}{2} = k\sin\phi$. What range of ϕ corresponds to $0 \le \theta \le \gamma$? What happens to the integrand?

STAGE 4 The first correction

Ask yourself For small k, expand $(1 - k^2\sin^2\phi)^{-1/2}$ by the binomial theorem. What is $\int_0^{\pi/2} \sin^2\phi \, d\phi$?

Try this next

1. **A tethered particle.** Try STEP III 2015, Question 9 (step.maths.org/questions/15-s3-q9): an oscillator whose period depends on its amplitude even for small oscillations. Why does it, when a pendulum's does not?

2. **The next term.** Carry the binomial expansion one term further and show that $P \approx 2\pi\sqrt{\frac{l}{g}}(1 + \frac{\gamma^2}{16} + \frac{11\gamma^4}{3072})$.

3. **The AGM by hand.** Use three steps of the AGM to estimate the period of a pendulum with amplitude $90°$, and compare with the value 1.1803 in the figure.

4. **Clocks.** A pendulum clock is regulated to keep perfect time with an amplitude of $4°$. If its amplitude increases to $5°$, how many seconds a day does it lose?

Further reading

- **David Morin, *Introduction to Classical Mechanics*** (Cambridge University Press), Chapter

4, which treats the large-amplitude pendulum.

- **Jonathan Borwein & Peter Borwein, *Pi and the AGM*** (Wiley). Gauss's discovery and its uses, including the fastest ways to compute π.
- **Dava Sobel, *Longitude*** (Fourth Estate). The story of timekeeping at sea, beyond the pendulum.

The full guided solution, and what the question leads to at university, is in Section 11.3:
https://beyondstep.keithmcnulty.org/ch11/real-pendulum.html