12.16 Markov Chains Forwards and Backwards

WORKSHEET

The question

STEP III 1996, Question 12

Read the question before you go on. It is in the STEP Support Programme's question database:

step.maths.org/questions/96-s3-q12

Hints

Try the question on your own first. If you get stuck, read the hints *one at a time*, and go back to the question after each one. Hints marked *Pause and try* are the key steps.

Ask yourself What does the condition "each row adds up to 1" mean? Do the columns add up to 1?

STAGE 2 The stationary distribution

Ask yourself We want $\mathbf{a} = \mathbf{a}P$ with $a_1 + a_2 + a_3 = 1$. Write out the equations. Which are the simplest?

STAGE 3 Running the chain backwards

Ask yourself $q_{ij} = \mathrm{P}(\text{previous } i \mid \text{current } j)$. Use Bayes' theorem, with the stationary distribution for both the previous and the current theorem.

Try this next

1. **A rabbit warren.** Try STEP III 2003, Question 13 (step.maths.org/questions/03-s3-q13): a random walk on the corners of a square with an exit at each corner, solved by first-step equations.
2. **Two steps.** If Professor Ecks's last theorem was new and correct, find the probabilities of the three types for the theorem after next.
3. **Eigenvalues.** Show that the eigenvalues of P are 1, 0.2 and 0.1.
4. **Reversible?** Check whether $a_i p_{ij} = a_j p_{ji}$ for Professor Ecks's chain. What does the answer tell you?

Further reading

- **James Norris, *Markov Chains*** (Cambridge University Press). The standard undergraduate text.
- **Olle Häggström, *Finite Markov Chains and Algorithmic Applications*** (Cambridge University Press). Short, and good on Markov chain Monte Carlo.
- **Brian Hayes, "First links in the Markov chain"**, *American Scientist* 101 (2013). Markov's Pushkin analysis, retold.

The full guided solution, and what the question leads to at university, is in Section 12.16:
https://beyondstep.keithmcnulty.org/ch12/markov.html