

# Topology

K

May 11, 2022



# Contents

<b>1</b>	<b>Topological Space</b>	<b>5</b>
1.1	Definitions and Theorems . . . . .	5
1.2	Proofs, Remarks, and Examples . . . . .	7
<b>2</b>	<b>Connected Spaces and Sets</b>	<b>9</b>
2.1	Definition and Theorems . . . . .	9
2.2	Proofs, Remarks, and Examples . . . . .	10
2.3	Exercises and Notes . . . . .	11
<b>3</b>	<b>Separation Axioms</b>	<b>13</b>
<b>4</b>	<b>Compact Spaces</b>	<b>15</b>
4.1	Proofs, Remarks, and Examples . . . . .	16



# Chapter 1

## Topological Space

### 1.1 Definitions and Theorems

**Definition 1** (Topological Space). A **topological space** is an **ordered pair**  $(X, \tau)$ , where  $X$  is a **set** and  $\tau$  is a **collection of subsets** that satisfies the following **axioms**.

1. The **empty set**  $\emptyset$  and the **entire set**  $X$  belongs to  $\tau$ .
2. Any **arbitrary union** of members of  $\tau$  belongs to  $\tau$ .
3. The **intersection** of **finite number** of members of  $\tau$  belongs to  $\tau$ .

The **collection**  $\tau$  is called a **topology** on  $X$  and the **elements** of  $\tau$  are called **open sets**. A **subset**  $A \subset X$  is said to be **closed** if its **complement**  $X \setminus A$  is **open**.

**Definition 2** (Continuous Maps). Let  $(X, \tau_X)$  and  $(Y, \tau_Y)$  be **topological spaces**. A **map**  $f : X \longrightarrow Y$  is said to be **continuous** if the preimage of an open subset is again open, i.e.

$$\text{for all } U \in \tau_Y \text{ it is } f^{-1}(U) \in \tau_X. \quad (1.1)$$

**Lemma 3.** The different definitions of continuity in a topological space and a metric space are equivalent, i.e. if  $X$  and  $Y$  are metric spaces, then  $f : X \longrightarrow Y$  is  $\epsilon$ - $\delta$ -continuous if and only if  $f$  is continuous.

**Definition 4** (Homeomorphism). Let  $X$  and  $Y$  be **topological spaces**.

1. A **map**  $f : X \longrightarrow Y$  is a **homeomorphism** if it has the following properties.
  - (a)  $f$  is **bijective**.
  - (b)  $f$  and the **inverse map**  $f^{-1}$  is **continuous**.
2. Two topological spaces  $X$  and  $Y$  are said to be **homeomorphic** if a homeomorphism exists.
3. We denote the set of all homeomorphisms from  $X$  to  $Y$  by  $\text{Homeo}(X, Y)$ . If  $Y = X$  we also write  $\text{Homeo}(X)$ .

**Definition 5** (Homeomorphism). Let  $(X, \tau)$  a topological space.

1.  $\mathcal{B} \subset \mathcal{O}$  is a **basis** of the topology, if any member of  $\mathcal{O}$  is the **union of subsets** from  $\mathcal{B}$ .
2.  $\mathcal{S} \subset \mathcal{O}$  is a **subbasis** of the topology, if any member of  $\mathcal{O}$  is the **union of finite intersections of subsets** from  $\mathcal{S}$ .

We say that  $\mathcal{B}$  and  $\mathcal{S}$  **generates**  $\mathcal{O}$  and write  $\overline{\mathcal{S}} = \overline{\mathcal{B}} = \mathcal{O}$ .

**Lemma 6.** Let  $\mathcal{S} \subset \mathcal{P}(X)$  be a **collection of subsets**, then there **exists exactly one** topology  $\tau \subset \mathcal{P}(X)$  of  $X$  such that

1.  $\mathcal{S} \subset \tau$
2. If  $\tau' \subset \mathcal{P}(X)$  a topology with  $\mathcal{S} \subset \tau'$ , then  $\tau \subset \tau'$ .

**Definition 7.** 1. Given  $(X, \tau)$  be a **topological space**,  $S \subset X$  a subset, the **subspace topology** (also the induced topology or the relative topology) on  $S$  is defined by

$$\tau_S = \{ S \cap U \mid U \in \tau \}.$$

2. Let  $(X, \tau_X)$  and  $(Y, \tau_Y)$  be two **topological spaces**. The product topology of  $X$  and  $Y$  is defined by

$$\tau_{X \times Y} := \{ U \times V \mid U \in \tau_X \text{ and } V \in \tau_Y \}.$$

3. Let  $(X, \tau_X)$  and  $(Y, \tau_Y)$  be two **topological spaces**. The topological sum of  $X$  and  $Y$  is defined by

$$\tau_{X \sqcup Y} := \{ U \sqcup V \mid U \in \tau_X \text{ and } V \in \tau_Y \}.$$

**Definition 8.** Let  $(X, \tau)$  be a topological space.

1. Given a **point**  $p \in X$ , a subset  $U \subset X$  is a neighborhood of  $p$  if there is an open subset  $V \in \tau$  such that  $p \in V$ . If such a neighborhood exists,  $p$  is called a interior point of  $U$ .
2. Let  $S \subset X$  be a subset. The interior of  $S$ , denoted by  $\overset{\circ}{S}$  or  $\text{int}(S)$ , is the **set** of all interior points of  $S$ .
3. Let  $S \subset X$  be a subset. The closure of  $S$ , denoted by  $\overline{S}$  or  $\text{cl}(S)$ , is defined by

$$\text{cl}(S) := X \setminus \text{int}(X \setminus S).$$

## 1.2 Proofs, Remarks, and Examples

**Example 8.1.** Let  $X$  be a set.

1.  $\tau = \mathcal{P}(X)$  is called the **discrete topology**. In this case,  $(X, \tau)$  is called the **discrete space**. It is the **finest topology** that can be defined on a set. (The set of all possible topologies on a given set forms a partially ordered set.)
2.  $\tau = \{\emptyset, \mathcal{P}(X)\}$  is called the **trivial topology**.
3. Let  $(X, d)$  be a **metric space**. Set

$$\tau_d := \{U \in \mathcal{P}(X) \mid U \text{ is an open subset in the metric space } (X, d)\}. \quad (1.2)$$

Recall that  $U$  being an open subset in the metric space  $(X, d)$  means that for all  $x \in U$  there is an  $r > 0$  such that  $B_d(x, r)$  is contained in  $U$ .

Here,  $\tau$  is a topology. In other words, a metric induces a topology.

(Proof as homework.)

4. The Zariski-topology.

**Example 8.2.** List of natural topologies.

1. On  $\mathbb{R}^n$  the canonical topology, called the Euclidean topology, is generated by the basis that is formed by open balls, i.e. open subsets of  $\mathbb{R}^n$  are arbitrary unions of open balls. In other words, if  $A \in \mathcal{O}_{\mathbb{R}^n}$  and  $I$  is an index set, then

$$A = \bigcup_{i \in I} B_r(p) = \bigcup_{i \in I} \{x \in \mathbb{R}^n \mid d(p, x) < r\}.$$

This definition agrees with the topology endowed on arbitrary metric spaces.

**Remark.** The set of all homeomorphisms of  $X$  to itself  $\text{Homeo}(X)$  is a group with composition as its operation.

**Remark.** This lemma does not hold for basis.

**Remark.** 1.  $\tau_{X \times Y}$  is the most coarse topology for which both of the projections are continuous.

2.  $\tau_{X \sqcup Y}$  is the finest topology for which both the inclusions are continuous.

Note about product topology:  $\{U \times V \mid U \in \mathcal{O}_X, V \in \mathcal{O}_Y\}$ ; often  $W \subset X \times Y \iff \forall (x, y) \in W \exists U_X \in \mathcal{O}_X, V_Y \in \mathcal{O}_Y, x \in U_X, y \in V_Y$





## Chapter 2

# Connected Spaces and Sets

### 2.1 Definition and Theorems

**Definition 9.** A topological space  $X$  is said to be connected, if one of the following equivalent conditions is met.

1.  $X$  is not a union of two disjoint sets.
2. The only subsets of  $X$  that are both open and closed (clopen) are the emptyset  $\emptyset$  and the entire set  $X$ .
3. The only subsets of  $X$  with empty boundary are the emptyset  $\emptyset$  and the entire set  $X$ .
4. All continuous maps from  $X$  to the two point space  $\{0, 1\}$  endowed with the discrete topology is constant.

**Lemma 10.** Any interval  $I \subset \mathbb{R}$  is connected.

**Definition 11.** A connected component of a topological space is a maximally connected subset  $X_0 \subseteq X$ , i.e.  $X_0$  connected and for all  $X_0 \subsetneq X_1$  then  $X_1$  is not connected.

**Proposition 12.** Connected components are closed subsets.

**Lemma 13** (Lemma 11). Let  $X$  be connected and  $f : X \rightarrow Y$  and locally constant, i.e. for all  $x \in X$  there exists a  $U_x \in \mathcal{O}_X$ ,  $x \in U_x$  such that  $f$  restricted on  $U_x$  is identical to  $f(x)$ , then  $f$  is constant.

**Definition 14.**  $X$  is said to be path connected, if for every pair of points  $x$  and  $x_0$  in  $X$  there is a continuous map (called path)  $\gamma : [0, 1] \rightarrow X$  with  $\gamma(0) = x_0$  and  $\gamma(1) = x$ .

**Lemma 15.** If  $X$  is path connected, then it is also connected.

## 2.2 Proofs, Remarks, and Examples

**Lemma 16.** Any interval  $I \subset \mathbb{R}$  is connected.

*Proof.* Fix an interval  $I \subset \mathbb{R}$ , and let  $A, B \subset \mathbb{R}$  be two nonempty, open and disjoint subsets such that  $A \sqcup B = I$ . Moreover, let  $a \in A$  and  $b \in B$  and assume without loss of generality that  $a < b$ . If we set

$$s := \inf \{ x \in B \mid a < x \}, \quad (2.1)$$

then  $s \in I$  because  $s$  is between  $a$  and  $b$  and we have  $[a, b] \subset I$ .

Now, on one side, we have  $s \in \text{cl}(B)$  and since the complement of  $B$  is an open subset  $A$ , so  $B = \text{cl}(B)$ . It is therefore  $s \in B$ .

But we also have  $s \in A$  because the infimum cannot be contained in an open set, but  $s \in I = A \sqcup B$ .  $\square$

**Example 16.1.** The general linear group  $\text{GL}_n(K)$  for a field  $K$  and  $n \in \mathbb{N}$  is not connected for  $K = \mathbb{R}$  and  $K = \mathbb{C}$ .

**Remark.** Let  $f : X \rightarrow Y$  be continuous and  $X$  be connected, then  $f(X) \subset Y$  is connected.

*Proof.* Let  $f(X) = A \sqcup B$  with  $A$  and  $B$  being two open disjoint sets.  $f^{-1}(A)$  and  $f^{-1}(B)$  are open since  $f$  is continuous. We also have  $f^{-1}(A) \cap f^{-1}(B) = f^{-1}(A \cap B) = \emptyset$  so  $f^{-1}(A) = \emptyset$  or  $f^{-1}(B) = \emptyset$ , so  $A = \emptyset$  or  $B = \emptyset$  and we are done.  $\square$

*Proof.*  $\square$

**Example 16.2.** For  $\mathbb{Q} \subset \mathbb{R}$  the connected components are points and those are not open.

*Proof.* Locally constant implies continuous with regards to the discrete topology on  $Y$ . Let  $x \in X$ ,  $X = f^{-1}(f(x)) \cup f^{-1}(Y \setminus \{f(x)\})$  is a disjoint union and since  $X$  is connected  $f^{-1}(Y \setminus \{f(x)\}) = \emptyset$ . Conclude  $f$  is identical to  $f(x)$ .  $\square$

**Application:**  $f : X \rightarrow \{0, 1\}$ ,  $X$  is connected,  $f$  locally constant, there is a  $x \in X$  such that  $f(x) = 1$ , then  $f$  is identical to 1.

*Proof.* Let  $A$  and  $B$  two disjoint open sets such that  $A \sqcup B = X$ , and let  $a \in A$  and  $b \in B$ . Let  $\gamma : [0, 1] \rightarrow X$  be continuous path with  $\gamma(0) = a$  and  $\gamma(1) = b$ . We have that  $\gamma^{-1}$   $\square$

**Remark.** The converse statement is not true in general.

**Example 16.3.**  $X = \{ (x, \sin(\frac{1}{x})) \mid x > 0 \} \cup \{0\} \times [-1, 1] \subset \mathbb{R}^2$  is connected but not path connected.

*Proof.* Homework  $\square$

**Remark.** missing

## 2.3 Exercises and Notes

If  $X$  and  $Y$  are two connected topological spaces, then their product  $X \times Y$  is also connected.

*Proof.*

□



## Chapter 3

# Separation Axioms

Literature: Groessere Liste in Sten, Seibeck

**Definition 17** ( $T_1$  Space). Let  $X$  be a topological space.

1. We say that two points  $x$  and  $y$  can be separated if each lies in a neighborhood that does **not** contain the other point.
2. A topological space  $X$  is a  $T_1$  space if any two distinct points in  $X$  are separated.

**Proposition 18.** Let  $X$  be a topological space. Then, the following are equivalent.

1.  $X$  is a  $T_1$  space.
2. Points are closed in  $X$ , i.e. given any  $x \in X$ , the singleton set  $\{x\}$  is a closed set.

**Definition 19** ( $T_2$  Space). Let  $X$  be a topological space.

1. Points  $x$  and  $y$  in  $X$  can be separated by neighborhood if there exists a neighborhood  $U$  of  $x$  and a neighborhood  $V$  of  $y$  such that  $U$  and  $V$  are disjoint, i.e.  $U \cap V = \emptyset$ .
2. A topological space  $X$  is a  $T_2$  space if any two distinct points in  $X$  are separated by neighborhood.

**Proposition 20.** Let  $X$  be a topological space. Then, the following are equivalent.

1.  $X$  is a  $T_2$  space.
2. Any singleton set  $\{x\}$  is the intersection of all closed neighborhoods of  $x$ .
3. The diagonal  $\Delta = \{(x, x) \mid x \in X\}$  is closed as a subset of the product space  $X \times X$ .

**Proposition 21.**  $T_2$  spaces are also  $T_1$  spaces.



## Chapter 4

# Compact Spaces

**Definition 22.** 1. A topological space  $X$  is called compact if each of its open cover has a finite subcover.

2. A topological space  $X$  is called sequentially compact if every sequence in  $X$  has a convergent subsequence whose limit is in  $X$ .

**Theorem 23.** Satz 17

**Theorem 24.** Let  $A \subset \mathbb{R}^n$  be a subset.  $A$  is compact if and only if it is closed and bounded.

**Theorem 25.** Let  $X$  be a  $T_2$  space. If a subset  $K \subset X$  is compact, then it is closed.

**Theorem 26.** Let  $X$  and  $Y$  be topological spaces,  $X$  compact, and  $Y$  be a  $T_2$  space. If  $f : X \rightarrow Y$  is bijective and continuous, then the inverse function  $f^{-1}$  is continuous.

## 4.1 Proofs, Remarks, and Examples

**Lemma 27.**  $[0, 1] \subset \mathbb{R}$  is compact.