

Commutative Ring Theory

Kei Thoma

May 3, 2023

Contents

1	Introduction and Motivation	5
2	Metric Spaces	7
3	Topological Spaces	9
4	Products, Sequential Continuity, and Nets	11
5	Compactness	13

Chapter 1

Introduction and Motivation

Chapter 2

Metric Spaces

Chapter 3

Topological Spaces

Chapter 4

Products, Sequential Continuity, and Nets

Lemma 1 (Lemma 4.15). In any space X , a subset $A \subset X$ is open if and only if every point $x \in A$ has a neighbourhood $\mathcal{V} \subset X$ that is contained in A .

Proof. “ \Rightarrow ”: If A is open, then A itself can be taken as the desired neighbourhood of every $x \in A$. “ \Leftarrow ”: Let every point $x \in A$ have a neighbourhood $\mathcal{V} \subset X$ that is contained in A . Denote the open sets of these neighbourhoods by \mathcal{U}_x . Then, A is the union of all these open sets \mathcal{U}_x and thus open. \square

Lemma 2 (Lemma 4.16). In any first-countable topological space X , a subspace $A \subset X$ is not open if and only if there exists a point $x \in A$ and a sequence $x_n \in X \setminus A$ such that $x_n \rightarrow x$.

Proof. “ \Leftarrow ”: (Proof by contraposition.) If $A \subset X$ is open, then for every $x \in A$ and sequence $x_n \in X$ converging to x , we cannot have $x_n \in X \setminus A$ for all n since A is a neighbourhood of x . This is true so far for all topological spaces, with or without first-countability axiom, but the latter will be needed to prove the converse.

“ \Rightarrow ”: So suppose now that $A \subset X$ is not open, which by Lemma 4.15, means there exists a point $x \in A$ such that no neighbourhood $\mathcal{V} \subset X$ of x is contained in A . Fix a countable neighbourhood base $\mathcal{U}_1, \mathcal{U}_2, \dots$ for x . XXX

Observe that since none of the neighbourhoods \mathcal{U}_n can be contained in A , there exists a sequence of points

$$x_n \in \mathcal{U}_n \text{ such that } x_n \notin A.$$

This sequence converges to x since every neighbourhood $\mathcal{V} \subset X$ of x contains one of \mathcal{U}_N , implying that for all $n \geq N$,

$$x_n \in \mathcal{U}_n \subset \mathcal{U}_N \subset \mathcal{V}.$$

\square

Definition 3. A **directed set** $(I, <)$ consists of a set I with a partial order $<$ such that for every pair $\alpha, \beta \in I$, there exists an element $\gamma \in I$ with $\gamma < \alpha$ and $\gamma < \beta$.

Definition 4. Given a space X , a net $\{x_\alpha\}_{\alpha \in I}$ in X is a function $I \rightarrow X : \alpha \mapsto x_\alpha$ where $(I, <)$ is a directed set.

Chapter 5

Compactness

Definition 5. A subset $A \subset X$ is compact if every open cover of A has a finite subcover, i.e. given an arbitrary open cover $\{\mathcal{U}_\alpha\}_{\alpha \in I}$ of A , one can always find a finite subset $\{\alpha_1, \dots, \alpha_N\} \subset I$ such that $A \subset \mathcal{U}_{\alpha_1} \cup \dots \cup \mathcal{U}_{\alpha_N}$. We say that X itself is a compact space if X is compact subset of itself.