

Exercise Sheet 2

Exercise 1

Solution

1. $\mathbb{Z} \times \mathbb{Z}$ is not a Dedekind domain as it is not even an integral domain. Take $(1, 0) \in \mathbb{Z} \times \mathbb{Z}$ and $(0, 1) \in \mathbb{Z} \times \mathbb{Z}$ for example. $(1, 0) \cdot (0, 1) = (0, 0)$ even though we chose nonzero elements.
2. $\mathbb{Z}[X]/(X^2 + 3)$ is not a Dedekind domain as it is not integrally closed.

First, define a ring homomorphism $\varphi : \mathbb{Z}[X] \rightarrow \mathbb{Z}$ that substitutes X with $\sqrt{-3}$. We have $\varphi(\mathbb{Z}[X]) = \mathbb{Z}[\sqrt{-3}]$ and $\ker(\varphi) = X^2 + 3$. With the isomorphism theorem, we have $\mathbb{Z}[X]/(X^2 + 3) \cong \mathbb{Z}[\sqrt{-3}]$.

Consider

$$\alpha := \frac{1}{2} + \frac{1}{2}\sqrt{-3} \in \text{Quot}(\mathbb{Z}[\sqrt{-3}]) = \mathbb{Q}(\sqrt{-3}). \quad (1)$$

From example 3.2.5. (script), we know that

$$\mathcal{O}_{\mathbb{Q}(\sqrt{-3})} = \mathbb{Z}[\alpha]. \quad (2)$$

Therefore, α is integral over \mathbb{Z} and hence over $\mathbb{Z}[\sqrt{-3}]$ as well, but it does not lie in $\mathbb{Z}[\sqrt{-3}]$. We conclude $\mathbb{Z}[\sqrt{-3}]$ and with it $\mathbb{Z}[X]/(X^2 + 3)$ is not integrally closed.

3. $\mathbb{F}_{11}[X]$ is a Dedekind domain.

From remark 1.0.3. / 2. (script), we have that the ring of polynomials in one variable over a field is a Euclidean ring, so $\mathbb{F}_{11}[X]$ is a Euclidean ring. Every Euclidean ring is a principal ideal domain (remark 1.0.3. / 3. from the script) and every principal ideal domain is a Dedekind domain (example 4.1.10. from the script). Hence, $\mathbb{F}_{11}[X]$ is a Dedekind domain.

4. $\mathbb{R}[X, Y]$ is not a Dedekind domain because