

Topology

K

May 11, 2022

Contents

1	Topological Space	5
1.1	Definitions and Theorems	5
1.2	Proofs, Remarks, and Examples	7
2	Connected Spaces and Sets	9
2.1	Definition and Theorems	9
2.2	Proofs, Remarks, and Examples	10
3	Separation Axioms	11

Chapter 1

Topological Space

1.1 Definitions and Theorems

Definition 1 (Topological Space). A **topological space** is an **ordered pair** (X, τ) , where X is a **set** and τ is a **collection of subsets** that satisfies the following **axioms**.

1. The **empty set** \emptyset and the **entire set** X belongs to τ .
2. Any **arbitrary union** of members of τ belongs to τ .
3. The **intersection** of **finite number** of members of τ belongs to τ .

The **collection** τ is called a **topology** on X and the **elements** of τ are called **open sets**. A **subset** $A \subset X$ is said to be **closed** if its **complement** $X \setminus A$ is **open**.

Definition 2 (Continuous Maps). Let (X, τ_X) and (Y, τ_Y) be **topological spaces**. A **map** $f : X \longrightarrow Y$ is said to be **continuous** if the preimage of an open subset is again open, i.e.

$$\text{for all } U \in \tau_Y \text{ it is } f^{-1}(U) \in \tau_X. \quad (1.1)$$

Lemma 3. The different definitions of continuity in a topological space and a metric space are equivalent, i.e. if X and Y are metric spaces, then $f : X \longrightarrow Y$ is ϵ - δ -continuous if and only if f is continuous.

Definition 4 (Homeomorphism). Let X and Y be **topological spaces**.

1. A **map** $f : X \longrightarrow Y$ is a **homeomorphism** if it has the following properties.
 - (a) f is **bijective**.
 - (b) f and the **inverse map** f^{-1} is **continuous**.
2. Two topological spaces X and Y are said to be **homeomorphic** if a homeomorphism exists.
3. We denote the set of all homeomorphisms from X to Y by $\text{Homeo}(X, Y)$. If $Y = X$ we also write $\text{Homeo}(X)$.

Definition 5 (Homeomorphism). Let (X, τ) a topological space.

1. $\mathcal{B} \subset \mathcal{O}$ is a **basis** of the topology, if any member of \mathcal{O} is the **union of subsets** from \mathcal{B} .
2. $\mathcal{S} \subset \mathcal{O}$ is a **subbasis** of the topology, if any member of \mathcal{O} is the **union of finite intersections of subsets** from \mathcal{S} .

We say that \mathcal{B} and \mathcal{S} **generates** \mathcal{O} and write $\overline{\mathcal{S}} = \overline{\mathcal{B}} = \mathcal{O}$.

Lemma 6. Let $\mathcal{S} \subset \mathcal{P}(X)$ be a **collection of subsets**, then there **exists exactly one** topology $\tau \subset \mathcal{P}(X)$ of X such that

1. $\mathcal{S} \subset \tau$
2. If $\tau' \subset \mathcal{P}(X)$ a topology with $\mathcal{S} \subset \tau'$, then $\tau \subset \tau'$.

Definition 7. 1. Given (X, τ) be a **topological space**, $S \subset X$ a subset, the **subspace topology** (also the induced topology or the relative topology) on S is defined by

$$\tau_S = \{ S \cap U \mid U \in \tau \}.$$

2. Let (X, τ_X) and (Y, τ_Y) be two **topological spaces**. The product topology of X and Y is defined by

$$\tau_{X \times Y} := \{ U \times V \mid U \in \tau_X \text{ and } V \in \tau_Y \}.$$

3. Let (X, τ_X) and (Y, τ_Y) be two **topological spaces**. The topological sum of X and Y is defined by

$$\tau_{X \sqcup Y} := \{ U \sqcup V \mid U \in \tau_X \text{ and } V \in \tau_Y \}.$$

Definition 8. Let (X, τ) be a topological space.

1. Given a **point** $p \in X$, a subset $U \subset X$ is a neighborhood of p if there is an open subset $V \in \tau$ such that $p \in V$. If such a neighborhood exists, p is called a interior point of U .
2. Let $S \subset X$ be a subset. The interior of S , denoted by \mathring{S} or $\text{int}(S)$, is the **set** of all interior points of S .
3. Let $S \subset X$ be a subset. The closure of S , denoted by \overline{S} or $\text{cl}(S)$, is defined by

$$\text{cl}(S) := X \setminus \text{int}(X \setminus S).$$

1.2 Proofs, Remarks, and Examples

Example 8.1. Let X be a [set](#).

1. $\tau = \mathcal{P}(X)$ is called the [discrete topology](#). In this case, (X, τ) is called the [discrete space](#). It is the [finest topology](#) that can be defined on a set. (The set of all possible topologies on a given set forms a partially ordered set.)
2. $\tau = \{\emptyset, \mathcal{P}(X)\}$ is called the [trivial topology](#).
3. Let (X, d) be a [metric space](#). Set

$$\tau_d := \{ U \in X \mid U \text{ is a open subset in the metric space } (X, d) \}. \quad (1.2)$$

Recall that U being an open subset in the metric space (X, d) means that for all $x \in U$ there is an $r > 0$ such that $B_d(x, r)$ is contained in U .

Here, τ is a topology. In other words, a metric induces a topology.

(Proof as homework.)

4. The Zariski-topology.

Remark. The set of all homeomorphisms of X to itself $\text{Homeo}(X)$ is a group with composition as its operation.

Remark. This lemma does not hold for basis.

Remark. 1. $\tau_{X \times Y}$ is the most coarse topology for which both of the projections are continuous.

2. $\tau_{X \sqcup Y}$ is the finest topology for which both the inclusions are continuous.

Note about product topology: $\{ U \times V \mid U \in \mathcal{O}_X, V \in \mathcal{O}_Y \}$; often $W \subset X \times Y \iff \forall (x, y) \in W \exists U_X \in \mathcal{O}_X, V_Y \in \mathcal{O}_Y, x \in U_X, y \in V_Y$

Chapter 2

Connected Spaces and Sets

2.1 Definition and Theorems

Definition 9. A topological space X is said to be connected, if one of the following equivalent conditions is met.

1. X is not a union of two disjoint sets.
2. The only subsets of X that are both open and closed (clopen) are the emptyset \emptyset and the entire set X .
3. The only subsets of X with empty boundary are the emptyset \emptyset and the entire set X .
4. All continuous maps from X to the two point space $\{0, 1\}$ endowed with the discrete topology is constant.

Lemma 10. Any interval $I \subset \mathbb{R}$ is connected.

Definition 11. A connected component of a topological space is a maximally connected subset $X_0 \subseteq X$, i.e. X_0 connected and for all $X_0 \subsetneq X_1$ then X_1 is not connected.

Proposition 12. Connected components are closed subsets.

Lemma 13 (Lemma 11). Let X be connected and $f : X \rightarrow Y$ and locally constant, i.e. for all $x \in X$ there exists a $U_x \in \mathcal{O}_X$, $x \in U_x$ such that f restricted on U_x is identical to $f(x)$, then f is constant.

Definition 14. X is said to be path connected, if for every pair of points x and x_0 in X there is a continuous map (called path) $\gamma : [0, 1] \rightarrow X$ with $\gamma(0) = x_0$ and $\gamma(1) = x$.

Lemma 15. If X is path connected, then it is also connected.

2.2 Proofs, Remarks, and Examples

Proof. Fix an interval $I \subset \mathbb{R}$, and let $A, B \subset \mathbb{R}$ be two nonempty, open and disjoint subsets such that $A \sqcup B = I$. Moreover, let $a \in A$ and $b \in B$ and assume without loss of generality that $a < b$. If we set

$$s := \inf \{ x \in B \mid a < x \}, \quad (2.1)$$

then $s \in [a, b] \subset I$ because I is an interval. \square

Example 15.1. The general linear group $\mathrm{GL}_n(K)$ for a field K and $n \in \mathbb{N}$ is not connected for $K = \mathbb{R}$ and $K = \mathbb{C}$.

Remark. Let $f : X \rightarrow Y$ be continuous and X be connected, then $f(X) \subset Y$ is connected.

Proof. Let $f(X) = A \sqcup B$ with A and B being two open disjoint sets. $f^{-1}(A)$ and $f^{-1}(B)$ are open since f is continuous. We also have $f^{-1}(A) \cap f^{-1}(B) = f^{-1}(A \cap B) = \emptyset$ so $f^{-1}(A) = \emptyset$ or $f^{-1}(B) = \emptyset$, so $A = \emptyset$ or $B = \emptyset$ and we are done. \square

Proof. \square

Example 15.2. For $\mathbb{Q} \subset \mathbb{R}$ the connected components are points and those are not open.

Proof. Locally constant implies continuous with regards to the discrete topology on Y . Let $x \in X$, $X = f^{-1}(f(x)) \cup f^{-1}(Y \setminus \{f(x)\})$ is a disjoint union and since X is connected $f^{-1}(Y \setminus \{f(x)\}) = \emptyset$. Conclude f is identical to $f(x)$. \square

Application: $f : X \rightarrow \{0, 1\}$, X is connected, f locally constant, there is a $x \in X$ such that $f(x) = 1$, then f is identical to 1.

Proof. Let A and B two disjoint open sets such that $A \sqcup B = X$, and let $a \in A$ and $b \in B$. Let $\gamma : [0, 1] \rightarrow X$ be continuous path with $\gamma(0) = a$ and $\gamma(1) = b$. We have that γ^{-1} \square

Remark. The converse statement is not true in general.

Example 15.3. $X = \{ (x, \sin(\frac{1}{x})) \mid x > 0 \} \cup \{0\} \times [-1, 1] \subset \mathbb{R}^2$ is connected but not path connected.

Proof. Homework \square

Remark. missing

Chapter 3

Separation Axioms

Literature: Groessere Liste in Sten, Seibeck

Definition 16 (T_1 Space). Let X be a topological space.

1. We say that two points x and y can be separated if each lies in a neighborhood that does **not** contain the other point.
2. A topological space X is a T_1 space if any two distinct points in X are separated.

Proposition 17. Let X be a topological space. Then, the following are equivalent.

1. X is a T_1 space.
2. Points are closed in X , i.e. given any $x \in X$, the singleton set $\{x\}$ is a closed set.

Definition 18 (T_2 Space). Let X be a topological space.

1. Points x and y in X can be separated by neighborhood if there exists a neighborhood U of x and a neighborhood V of y such that U and V are disjoint, i.e. $U \cap V = \emptyset$.
2. A topological space X is a T_2 space if any two distinct points in X are separated by neighborhood.

Proposition 19. Let X be a topological space. Then, the following are equivalent.

1. X is a T_2 space.
2. Any singleton set $\{x\}$ is the intersection of all closed neighborhoods of x .
3. The diagonal $\Delta = \{(x, x) \mid x \in X\}$ is closed as a subset of the product space $X \times X$.

Proposition 20. T_2 spaces are also T_1 spaces.