

Chapter 1

Commutative Rings

Exercise 1.1. Let $\varphi : A \longrightarrow B$ be a ring homomorphism, $\mathfrak{a}_1, \mathfrak{a}_2, \mathfrak{a}_3$ ideals in A , and $\mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3$ ideals of B . Prove the following statements.

1. $(\mathfrak{a}_1 + \mathfrak{a}_2)^e = (\mathfrak{a}_1)^e + (\mathfrak{a}_2)^e$.

Proof. We show $(\mathfrak{a}_1 + \mathfrak{a}_2)^e \subseteq (\mathfrak{a}_1)^e + (\mathfrak{a}_2)^e$. Let $x \in (\mathfrak{a}_1 + \mathfrak{a}_2)^e$, then we have for some index set I

$$x = \sum_{i \in I} \lambda_i x_i, \quad (1.1)$$

where $\lambda_i \in B$ and $x_i \in \varphi(\mathfrak{a}_1 + \mathfrak{a}_2)$ for all $i \in I$. For each $i \in I$ it is $x_i = \varphi(\mu_{i,1}a_{i,1} + \mu_{i,2}a_{i,2})$, hence

$$x = \sum_{i \in I} \lambda_i \varphi(\mu_{i,1}a_{i,1} + \mu_{i,2}a_{i,2}) \quad (1.2)$$

$$= \sum_{i \in I} \lambda_i (\varphi(\mu_{i,1}a_{i,1}) + \varphi(\mu_{i,2}a_{i,2})) \quad (\text{by linearity}) \quad (1.3)$$

$$= \sum_{i \in I} \lambda_i (\mu_{i,1}\varphi(a_{i,1}) + \mu_{i,2}\varphi(a_{i,2})) \quad (\text{by linearity}) \quad (1.4)$$

$$= \sum_{i \in I} \lambda_i \mu_{i,1} \varphi(a_{i,1}) + \lambda_i \mu_{i,2} \varphi(a_{i,2}) \quad (\text{by distributivity}) \quad (1.5)$$

$$= \sum_{i \in I} \lambda_i \mu_{i,1} \varphi(a_{i,1}) + \sum_{i \in I} \lambda_i \mu_{i,2} \varphi(a_{i,2}) \quad (\text{reordering the sum}). \quad (1.6)$$

$$(1.7)$$

The last term is exactly the elements expressed by $\mathfrak{a}_1^e + \mathfrak{a}_2^e$, therefore, $(\mathfrak{a}_1 + \mathfrak{a}_2)^e \subseteq (\mathfrak{a}_1)^e + (\mathfrak{a}_2)^e$.

I think the above proof should work into both directions. \square