


Lecture 2

Linear transformations

and Diagonalization



카카오톡 단톡방 MATLAB KR

<https://open.kakao.com/o/gKBr7Zde>

1. Linear transformation

1.1 Standard matrix for a linear transformation

So, what is a matrix?

- Five perspectives on viewing matrices
 - Collection of data (numeric)
 - Collection of vectors (geometric)
 - A system of linear equations (algebraic)
 - **A linear transformation (operational)**
 - A tangent space of a function (differential)

Matrix as a linear transformation

$$A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}$$

$$A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 + 1x_2 \\ 0x_1 + 2x_2 \\ 1x_1 + 1x_2 \end{bmatrix}$$

- The matrix A represents a function


$$T_A: \mathbb{R}^2 \rightarrow \mathbb{R}^3$$

- T_A is *linear*, i.e.,

$$T_A(u + v) = T_A(u) + T_A(v)$$

$$T_A(cu) = cT_A(u)$$

Matrix as a linear transformation



- In fact, we *do not need* a matrix.
- All we need is the *images of basis vectors*.
- Standard basis of \mathbb{R}^2

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Standard matrix

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\begin{aligned} T(\mathbf{x}) &= x_1 T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + x_2 T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\ &= [T(\mathbf{e}_1) \quad T(\mathbf{e}_2)] \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\ &= \underset{\substack{\uparrow \\ \text{Standard matrix for } T}}{A} \mathbf{x} \end{aligned}$$

- Every linear transformation from \mathbb{R}^n to \mathbb{R}^m is represented by an $m \times n$ matrix.
- The inverse also holds.

- Standard matrix for T , denoted by $[T] \Rightarrow T(x) = [T]x$



Examples of linear transformations

- Stretch / compression
 - Shear
 - Rotation
 - Reflection
-
- lab: `example_matrix_is_a_linear_transformation.m`
-
- Q. Is translation a *linear* transformation? (cf. [Affine transformation](#))

Invertibility of a square matrix

- An $n \times n$ matrix A is invertible if and only if
 - $Ax = 0$ has only a trivial solution, i.e., $\ker(T_A) = \{0\}$.
 - $\det(A) \neq 0$
 - A set of column (row) vectors of A forms a basis of \mathbb{R}^n , i.e., $\text{Im}(T) = \mathbb{R}^n$.
 - $\text{rank}(A) = n$, $\text{nullity}(A) = 0$
 - T_A is 1-1 and onto.
- Q. Can an $m \times n$ matrix represent a bijective function?


Direction of stretching



- lab: `example_matrix_is_a_linear_transformation.m`

1.2 Projection and least square

Projection formula



- Projection of x onto v


$$\text{proj}_v x = \frac{x \cdot v}{\|v\|^2} v$$

$$= (x \cdot v) v \quad \text{if } \|v\| = 1$$

- Every vector x in \mathbb{R}^n is *uniquely* expressed as

$$x = \text{proj}_v x + \text{proj}_{v^\perp} x$$

Projection operator



- Projection operator onto $\text{span}\{v\}$


$$T(x) = \text{proj}_v x = \left(\frac{1}{v^T v} v v^T \right) x$$

- Standard matrix for T

$$P = \frac{1}{v^T v} v v^T \quad (= v v^T \text{ if } \|v\| = 1)$$

- Q. $\text{rank}(P) = ?$

Projection onto a *subspace*



- Projection of $x \in \mathbb{R}^n$ onto $W \leq \mathbb{R}^n$

$$\text{proj}_W x = M(M^T M)^{-1} M^T x$$

- where the column vectors of M form a basis of W .

- lab: example_projection_operator.m

Least square and projection operation

Linear least squares [\[edit\]](#)

Main article: [Linear least squares](#)

A regression model is a linear one when the model comprises a [linear combination](#) of the parameters, i.e.,

$$f(x, \beta) = \sum_{j=1}^m \beta_j \phi_j(x),$$

where the function ϕ_j is a function of x .^[12]

Letting $X_{ij} = \phi_j(x_i)$ and putting the independent and dependent variables in matrices X and Y , respectively, we can compute the least squares in the following way. Note that D is the set of all data.^{[12][13]}

$$\begin{aligned} L(D, \beta) &= \|Y - X\beta\|^2 = (Y - X\beta)^\top (Y - X\beta) = Y^\top Y - Y^\top X\beta - \beta^\top X^\top Y + \beta^\top X^\top X\beta \\ &= Y^\top Y - X^\top Y\beta - X^\top Y\beta + X^\top X\beta^2 \end{aligned}$$

The gradient of the loss is:

$$\frac{\partial L(D, \beta)}{\partial \beta} = \frac{\partial (Y^\top Y - X^\top Y\beta - X^\top Y\beta + X^\top X\beta^2)}{\partial \beta} = -2X^\top Y + 2X^\top X\beta$$


Setting the gradient of the loss to zero and solving for β , we get:^{[13][12]}

$$-2X^\top Y + 2X^\top X\beta = 0 \Rightarrow X^\top Y = X^\top X\beta$$

$$\hat{\beta} = (X^\top X)^{-1} X^\top Y$$

$$\text{proj}_W x = M(M^\top M)^{-1} M^\top x$$

Ordinary Least Squares




$$c_0 + c_1 x_1 + e_1 = y_1$$

$$c_0 + c_1 x_2 + e_2 = y_2$$

$$c_0 + c_1 x_3 + e_3 = y_3$$

$$\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \end{bmatrix} + \begin{bmatrix} e_1 \\ e_2 \\ e_3 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

$$A x + e = y$$



- Least square method


≡ To find x and \hat{y} that minimize $\|e\|$

≡ Projecting y onto $\text{col}(A)$ which is \hat{y}

$$\text{proj}_{\text{col}(A)} y = A(A^T A)^{-1} A^T y$$

- lab: example_least_square_is_projection.m

Linear regression example(?)



- lab: example_triangle_game.m

Orthonormal basis

- $v \in \mathbb{R}^n, \|v\| = 1$

➔ Projection operator onto $\text{span}\{v\}$

$$T(x) = (vv^T)x$$



- Standard matrix for T

- M is an orthonormal basis of $W \leq \mathbb{R}^n$

➔ Projection operator onto $\text{col}(M)$

$$T(x) = M(M^T M)^{-1} M^T x$$

$$= MM^T x$$



Standard matrix for T

Why do we care orthonormal bases?

- [Thm] If $\{v_1, \dots, v_k\}$ is an orthonormal basis for $W \leq \mathbb{R}^n$, then the orthogonal projection of a vector $x \in \mathbb{R}^n$ onto W is expressed as

$$\text{proj}_W x = (x \cdot v_1)v_1 + (x \cdot v_2)v_2 + \dots + (x \cdot v_k)v_k$$

- [Thm] If $\{v_1, \dots, v_k\}$ is an orthonormal basis for $W \leq \mathbb{R}^n$, then any $w \in W$ is expressed as

$$w = (w \cdot v_1)v_1 + (w \cdot v_2)v_2 + \dots + (w \cdot v_k)v_k$$


Gram-Schmidt and QR decomposition

- From a basis $\{w_1, \dots, w_k\}$ for $W \leq \mathbb{R}^n$, an orthonormal basis of W can be constructed as follows:
 - 1. Normalize w_1 to u_1 such that $|u_1| = 1$.
 - 2. Calculate $v_2 = w_2 - \text{proj}_{u_1} w_2$ and normalize to u_2 .
 - 3. Calculate $v_3 = w_3 - \text{proj}_{\text{span}\{u_1, u_2\}} w_3$ and normalize to u_3 .
 - 4. Repeat to v_k and u_k .
- Rmk. $\text{span}\{w_1, \dots, w_l\} = \text{span}\{u_1, \dots, u_l\}$ for any $l \leq k$.
- lab: example_Gram_Schmidt_and_qr_decomposition.m

2. Coordinate and change of basis

2.1 Vectors and coordinates

Making the familiar *Unfamiliar*



$$\begin{bmatrix} mg \sin \theta \\ -mg \cos \theta \end{bmatrix} \text{ vs } \begin{bmatrix} 0 \\ -mg \end{bmatrix}$$

- What is a vector?
 - An arrow?
 - A tuple of numbers?
 - Something with magnitude and direction?
- Vector \neq Coordinate
- Standard basis is not that *special*.

Coordinate of a vector

- [Def] If $B = \{v_1, \dots, v_k\}$ is an ordered basis for $W \leq \mathbb{R}^n$, and if

$$w = a_1 v_1 + \dots + a_k v_k$$

- then we call a_1, \dots, a_k the *coordinate* of w with respect to B , and

$$[w]_B = \begin{bmatrix} a_1 \\ \vdots \\ a_k \end{bmatrix}$$

- the coordinate matrix for w with respect to B .

Why do we care orthonormal bases?

- [Thm] If B is an orthonormal basis for a k -dim. $W \leq \mathbb{R}^n$, and if u , v , and w are vectors in W such that

$$[u]_B = \begin{bmatrix} u_1 \\ \vdots \\ u_k \end{bmatrix}, [v]_B = \begin{bmatrix} v_1 \\ \vdots \\ v_k \end{bmatrix}, [w]_B = \begin{bmatrix} w_1 \\ \vdots \\ w_k \end{bmatrix}$$

- then

- (a) $\|w\| = \sqrt{w_1^2 + w_2^2 + \cdots + w_k^2} = \|[w]_B\|$

- (b) $u \cdot v = u_1 v_1 + u_2 v_2 + \cdots + u_k v_k = [u]_B \cdot [v]_B$

Change of basis

- How to change from $[w]_B$ to $[w]_{B'}$?
- [Thm] Given $w \in \mathbb{R}^n$, two bases of \mathbb{R}^n $B = \{v_1, \dots, v_n\}$ and $B' = \{v'_1, \dots, v'_n\}$,

$$[w]_{B'} = P_{B \rightarrow B'} [w]_B$$

- where $P_{B \rightarrow B'}$ is the coordinate change matrix from B to B'

$$P_{B \rightarrow B'} = [[v_1]_{B'}, \dots, [v_n]_{B'}]$$

- Rmk. $P_{B' \rightarrow B} = (P_{B \rightarrow B'})^{-1}$
- Q. $P_{B \rightarrow B} = ?$


Change of basis

- If S is the standard basis of \mathbb{R}^n , then $P_{B \rightarrow S} = [[v_1]_S, \dots, [v_n]_S] = [v_1, \dots, v_n]$.
- Rmk. Change of basis from B to B' = from B to S , and then from S to B' .

$$P_{B \rightarrow B'} = P_{S \rightarrow B'} P_{B \rightarrow S} = (P_{B' \rightarrow S})^{-1} P_{B \rightarrow S} = [[v'_1]_S, \dots, [v'_n]_S]^{-1} [[v_1]_S, \dots, [v_n]_S]$$

- [Thm] If B and B' are orthonormal bases for \mathbb{R}^n , then the coordinate change matrices $P_{B \rightarrow B'}$ and $P_{B' \rightarrow B}$ are orthogonal.

Change of basis – example




$$S = \{e_1, e_2\}$$

$$B = \{v_1, v_2\}$$


- How to interpret $V = [v_1, v_2]$?
- $V = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$
- 1) Rotation matrix
- 2) Coordinate change matrix $P_{B \rightarrow S}$

Change of basis




2.2 Matrix representation of linear transformations

Do linear transformations depend on bases?




$$\begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix}_S$$




$$\begin{bmatrix} 5 & 0 \\ 0 & 1 \end{bmatrix}_B$$

Do linear transformations depend on bases?



$$\frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix}_S$$



$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}_B$$

Do linear transformations depend on bases?

$$x \xrightarrow{T} T(x)$$

$$[x]_B \xrightarrow{A} A[x]_B$$

- The coordinate of x depends on the basis.
- T transforms a vector to another vector.
- What does the matrix A transform?
 - Vectors?
 - Coordinates of vectors?
- Is the matrix A for T unique?

Do linear transformations depend on bases?

- A matrix represents a linear transformation with respect to a *specific* basis.
- *Matrix representation* of a linear transformation depends on the *choice of basis*.
- Standard matrix: Matrix representation of a linear transformation with respect to the *standard* basis.

$$\begin{aligned} T(\mathbf{x}) &= x_1 T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + x_2 T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\ &= [T(\mathbf{e}_1) \quad T(\mathbf{e}_2)] \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\ &= \quad \quad A \quad \quad \mathbf{x} \end{aligned}$$

Matrix representation of linear transformations

- [Thm] Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear operator, $B = \{v_1, \dots, v_n\}$ be a basis for \mathbb{R}^n . Then $A = [[T(v_1)]_B, \dots, [T(v_n)]_B]$ is called the matrix for T with respect to the basis B , denoted by $[T]_B$, and satisfies

$$[T(x)]_B = A[x]_B = [T]_B[x]_B$$

- Q. What is a vector? (hint: not a tuple)
- Q. What is a matrix? (hint: not a rectangular arrangement of numbers)
- Q. What is a linear transformation? (hint: What is the linearity?)
- Q. How to get $[T]_B$ from $[T]_{B'}$?

Matrix representation of linear transformations

$$\begin{array}{ccc}
 [x]_S & \xrightarrow{[T]_S = A} & [T(x)]_S \\
 \downarrow P^{-1} & & \uparrow P \\
 [x]_B & \xrightarrow{[T]_B} & [T(x)]_B
 \end{array}$$

$$x \xrightarrow{T} T(x)$$

$$[x]_B \xrightarrow{[T]_B} [T(x)]_B = [T]_B [x]_B$$

- Recall that the coordinate change matrix from an ordered basis $B = \{v_1, \dots, v_n\}$ to the standard basis S is

$$P_{B \rightarrow S} = [v_1, \dots, v_n]$$



- Thus, the followings hold.

$$[T]_S = P[T]_B P^{-1}$$

$$[T]_B = P^{-1}[T]_S P$$

Matrix representation of linear transformations

- Example




$$S = \{e_1, e_2\}, B = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$$

$$P_{B \rightarrow S} = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

$$[T]_S = \begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 5 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}^{-1}$$


Matrix representation of linear transformations



- Example


$$S = \{e_1, e_2\}, B = \left\{ \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix} \right\}$$

$$P_{B \rightarrow S} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$



$$\begin{aligned} [T]_S &= \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix} \\ &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^{-1} \end{aligned}$$

Matrix representation of linear transformations



- For generalization, given a coordinate change matrix P from B to B' , the followings hold.

$$[T]_{B'} = P[T]_B P^{-1}$$

$$[T]_B = P^{-1}[T]_{B'} P$$

Matrix representation of linear transformations


- Bases of domain and codomain of a linear transformation may be *different*.
- [Thm] Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, $B = \{v_1, \dots, v_n\}$ be a basis for \mathbb{R}^n , $B' = \{u_1, \dots, u_m\}$ be a basis for \mathbb{R}^m . Then

$$A = \left[[T(v_1)]_{B'}, \dots, [T(v_n)]_{B'} \right]$$

- is the matrix for T with respect to the bases B and B' , denoted by $[T]_B^{B'}$, and satisfies

$$[T(x)]_{B'} = A[x]_B = [T]_B^{B'} [x]_B$$

Matrix representation of linear transformations




$$[T]_{B_1}^{B'_1} = U[T]_{B_2}^{B'_2}V^{-1} = [\text{Id}]_{B'_2}^{B'_1}[T]_{B_2}^{B'_2}[\text{Id}]_{B_1}^{B_2}$$

$$[T]_{B_1}^{B'_1}[x]_{B_1} = [\text{Id}]_{B'_2}^{B'_1}[T]_{B_2}^{B'_2}[\text{Id}]_{B_1}^{B_2}[x]_{B_1}$$

Matrix representation of linear transformations

- Rmk. Since the matrix representation depends on the both bases of domain and codomain, an identity matrix may **not** represent an identity transformation.



$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$$

$$[T]_S^B = [\text{Id}]_S^B [T]_S$$

3. Diagonalization

Some quote

- "This chapter begins the "*second half*" of linear algebra.

The first half was about $Ax = b$.

The new problem $Ax = \lambda x$ will still be solved making it *diagonal* if possible."

- Linear Algebra and Its Applications, Gilbert Strang.

Cengage

Fourth Edition

LINEAR ALGEBRA AND ITS APPLICATIONS



Gilbert Strang



Cengage Digital App includes
• Practice Exams



This edition is licensed for sale only in the United States. Reproduction, resale, or distribution of this edition outside the United States is prohibited.
UNIVERSITY MICROFILMS
PROHIBITED

3.1 Eigenvalues and Eigenvectors

Eigenvalues and Eigenvectors

- [Def] If A is an $n \times n$ matrix, then a scalar λ is called an eigenvalue of A if there is a **nonzero** vector x such that $Ax = \lambda x$. Such a vector x is called an eigenvector of A corresponding to λ .
- Rmk. Eigenvalues are the solutions of $|\lambda I - A| = 0$.
- [Def] For an $n \times n$ matrix A , $|\lambda I - A|$ is called the *characteristic polynomial* of A .
- Q. Does a rotation matrix have eigenvalues?

Eigenvalues and Eigenvectors



- Example


$$S = \{e_1, e_2\}, B = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$$

$$P_{B \rightarrow S} = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$


$$[T]_S = \begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 5 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}^{-1}$$

- What are the eigenvalues and eigenvectors of $[T]_S$?


Eigenvalues and Eigenvectors




$$S = \{e_1, e_2\} \quad \begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix}$$



$$B = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} \quad \begin{bmatrix} 5 & 0 \\ 0 & 1 \end{bmatrix}$$



Eigenvalues and Eigenvectors



$$S = \{e_1, e_2\}, B = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$$

$$P_{B \rightarrow S} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

$$[T]_S = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}^{-1}$$

- lab: example_direction_of_eigenvectors.m

Linear independence of eigenvectors

- [Thm] If v_1, \dots, v_k are eigenvectors of a matrix A that corresponding to distinct eigenvalues $\lambda_1, \dots, \lambda_k$, then the set $\{v_1, \dots, v_k\}$ is linearly independent.
- [Cor] If an $n \times n$ matrix A has n distinct eigenvalues, then the set of eigenvectors of A forms a basis of \mathbb{R}^n .
- Q. What if an $n \times n$ matrix A have less than n eigenvalues?

Multiplicity of eigenvalues and eigenspaces

- [Def] If the characteristic polynomial of an $n \times n$ matrix A is factored as

$$|\lambda I - A| = (\lambda - \lambda_1)^{m_1} (\lambda - \lambda_2)^{m_2} \dots (\lambda - \lambda_k)^{m_k}$$

- then m_i is called the **algebraic multiplicity** of the eigenvalue λ_i .

- [Def] If λ_0 is an eigenvalue of A , the solution space of

$$(\lambda_0 I - A)x = 0$$

- is called the **eigenspace** of A corresponding to λ_0 .
- The dimension of this eigenspace is called the **geometric multiplicity** of λ_0 .

Multiplicity of eigenvalues and eigenspaces

$$A = \begin{bmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{bmatrix}$$



- Find the...
 - Characteristic polynomial of A
 - Eigenvalues and eigenvectors
 - Algebraic multiplicity and geometric multiplicity of each eigenvalue
 - Eigenspace for each eigenvalue
- lab: example_eigenvalue_problem.m

Invertibility of a square matrix

- An $n \times n$ matrix A is invertible if and only if
 - $Ax = 0$ has only a trivial solution, i.e., $\ker(T_A) = \{0\}$.
 - $\det(A) \neq 0$
 - A set of column (row) vectors of A forms a basis of \mathbb{R}^n , i.e., $\text{Im}(T) = \mathbb{R}^n$.
 - $\text{rank}(A) = n$, $\text{nullity}(A) = 0$
 - T_A is 1-1 and onto.
 - 0 is not an eigenvalue of A .

3.2 Matrix similarity

What is the matrix similarity?



$$S = \{e_1, e_2\}$$

$$\begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix}$$

$$\begin{array}{ccc} [x]_S & \xrightarrow{[T]_S} & [T(x)]_S \\ \downarrow P^{-1} & & \uparrow P \\ [x]_B & \xrightarrow{[T]_B} & [T(x)]_B \end{array}$$

$$B = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$$

$$\begin{bmatrix} 5 & 0 \\ 0 & 1 \end{bmatrix}$$

Matrix similarity







- [Def] If A and C are square matrices with the same size, then we say that C is similar to A , denote $A \sim C$, if there exists an invertible matrix P such that $C = P^{-1}AP$.
- [Thm] If A and C are two matrix representations of the same linear operator, then A is similar to C . The inverse is also true.
- [Cor] If $A \sim C$, then $C \sim A$.

Similar matrices are *similar*

- [Thm] Similar matrices have similarity invariants such as:
 - the same determinant.
 - the same rank.
 - the same nullity.
 - the same trace.
 - the same characteristic polynomial.
 - the same eigenvalues with the same algebraic and geometric multiplicities.
- Q. Are the properties above for a matrix or a linear operator?
- Q. If two matrices have same properties above, are they similar?
- lab: example_similarity_check.m


3.3 Eigenvalue Decomposition

Motivation of Diagonalization



The Diagonalization Problem

- [Def] For a square matrix A , if there exists an invertible matrix P such that $P^{-1}AP$ is *diagonal*, then A is said to be **diagonalizable**, and P is said to diagonalize A .
- [Thm] An $n \times n$ matrix A is diagonalizable
 $\Leftrightarrow A$ has n linearly independent eigenvectors.
 \Leftrightarrow Eigenvectors of A form a basis of \mathbb{R}^n .

$$A = PDP^{-1}$$


Column vectors are eigenvectors. Diagonal terms are eigenvalues.

Diagonalizability of matrices

- [Thm] An $n \times n$ matrix with n distinct real eigenvalues is diagonalizable.
- [Thm] If A is an $n \times n$ matrix, then
 - (a) The geometric multiplicity of an eigenvalue of A is less than or equal to its algebraic multiplicity.
 - (b) A is diagonalizable.
 - \Leftrightarrow The sum of the geometric multiplicities of its eigenvalues is n .
 - \Leftrightarrow The geometric multiplicity of each eigenvalue of A is the same as its algebraic multiplicity.
- Rmk. Diagonalizability does not mean distinct eigenvalues.

Diagonalizability of matrices: example

$$A = \begin{bmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{bmatrix}$$

- Is A diagonalizable?
- If so, find an invertible matrix P and a diagonal matrix D such that $A = PDP^{-1}$.
- lab: example_eigenvalue_problem.m

We still care the orthonormal bases


$$\begin{array}{ccc} [x]_S & \xrightarrow{[T]_S = A} & [T(x)]_S \\ \downarrow P^{-1} & & \uparrow P \\ [x]_B & \xrightarrow{[T]_B} & [T(x)]_B \end{array}$$

- It is almost always better to have an orthonormal basis.
- When can a square matrix A be decomposed like this:


$$A = PDP^T$$

The Orthogonal Diagonalization Problem

- [Def] For a square matrix A , if there exists an *orthogonal* matrix P such that $P^T A P$ is *diagonal*, then A is said to be **orthogonally diagonalizable**, and P is said to orthogonally diagonalize A .



$$A = P D P^T$$



Column vectors
are eigenvectors.

Diagonal terms are
eigenvalues.

Orthogonal diagonalizability of matrices

- [Thm] An $n \times n$ matrix A is orthogonally diagonalizable.
 \Leftrightarrow There exists an orthonormal set of n eigenvectors of A .
 $\Leftrightarrow A$ is symmetric.
- [Thm] If a square matrix A is symmetric, then eigenvectors from different eigenspaces are orthogonal.
- Rmk. For any square matrix, eigenvectors from *different eigenspaces* are *linearly independent*.


Spectral decomposition

- If a matrix A is orthogonally diagonalizable, then

$$\begin{aligned} A &= P D P^T \\ &= \begin{bmatrix} | & & | \\ p_1 & \cdots & p_n \\ | & & | \end{bmatrix} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \begin{bmatrix} - & p_1^T & - \\ & \vdots & \\ - & p_n^T & - \end{bmatrix} \\ &= \lambda_1 p_1 p_1^T + \cdots + \lambda_n p_n p_n^T \end{aligned}$$

- which is called a spectral decomposition (or an **eigenvalue decomposition**).


Geometric interpretation of EVD



$$\begin{aligned} A &= \begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}^T \\ &= 1 \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} + 5 \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \end{aligned}$$





- lab: example_geometric_interpretation_of_EVD.m

eigshow



- Q. Equation of the ellipse?
- Q. Direction of major/minor axes?
- lab: eigshow.m

Geometric interpretation of diagonalization



- lab: example_diagonalization.m

3.4 Singular Value Decomposition

Motivation of SVD

- Even if a square matrix A is not symmetric, A may be decomposed as $A = PDP^{-1}$ where P is not orthogonal.
- However, an orthogonal matrix or an orthonormal basis is too good to abandon.
- What do we have to throw away?
 - 1) $A = PHP^T$ where P is orthogonal and H is upper-triangular. (Hessenberg, Schur)
 - 2) $A = U\Sigma V^T$ where U, V are orthogonal and Σ is diagonal. (SVD)
- Can we even decompose a non-square matrix in the form of $A = U\Sigma V^T$?


Singular Value Decomposition; SVD

- [Thm] If A is an $n \times n$ matrix of rank k , then A can be factored as

$$A = U \Sigma V^T$$
$$= \begin{bmatrix} | & & | \\ u_1 & \cdots & u_n \\ | & & | \end{bmatrix} \begin{bmatrix} \sigma_1 & & & & \\ & \ddots & & & \\ & & \sigma_k & & \\ & & & 0 & \\ & & & & \ddots \\ & & & & & 0 \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ & \vdots & \\ - & v_n^T & - \end{bmatrix}$$


- where U and V are $n \times n$ *orthogonal* matrices and Σ is an $n \times n$ *diagonal* matrix whose main diagonal has k positive entries and $n - k$ zeros.

Singular Value Decomposition; SVD







- Rmk. Domain and codomain of Σ have different bases.

eigshow



- Q. Equation of the ellipse?
- Q. Direction of major/minor axes?
- lab: eigshow.m

Geometric interpretation of SVD



- lab: example_SVD.m

Singular Value Decomposition; SVD

$$A = U\Sigma V^T = \begin{bmatrix} | & & | \\ u_1 & \cdots & u_n \\ | & & | \end{bmatrix} \begin{bmatrix} \sigma_1 & & & \\ & \ddots & & \\ & & \sigma_k & \\ & & & 0 \\ & & & & \ddots \\ & & & & & 0 \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ & \vdots & \\ - & v_n^T & - \end{bmatrix}$$

- Rmk.
 - $A^T A = V D V^T$ where D is diagonal and $\text{diag}(D) \geq 0$.
 - $\Sigma^2 = D$.
 - Main diagonal of Σ are in nonincreasing order.
 - σ_i : singular values of A
 - u_i : left singular vectors of A
 - v_i : right singular vectors of A

SVD of a general matrix

- [Thm] If A is an $m \times n$ matrix of rank k , then A can be factored as

$$A = U\Sigma V^T$$

$$= \begin{bmatrix} | & & | & | & & | \\ u_1 & \cdots & u_k & u_{k+1} & \cdots & u_m \\ | & & | & | & & | \end{bmatrix} \begin{bmatrix} \sigma_1 & & & & & \\ & \ddots & & & & \\ & & \sigma_k & & & \\ \hline & & & 0_{k \times (n-k)} & & \\ 0_{(m-k) \times k} & & & & 0_{(m-k) \times (n-k)} & \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ & \vdots & \\ - & v_k^T & - \\ - & v_{k+1}^T & - \\ & \vdots & \\ - & v_n^T & - \end{bmatrix}$$

$m \times m$

$m \times n$

$n \times n$

Reduced SVD


$$\begin{aligned}
 A = U\Sigma V^T &= \left[\begin{array}{c|c|c} | & & | \\ u_1 & \cdots & u_k \\ | & & | \end{array} \right] \begin{array}{c} | \\ u_{k+1} \\ | \end{array} \cdots \begin{array}{c} | \\ u_m \\ | \end{array} \left[\begin{array}{c|c} \boxed{\begin{array}{ccc} \sigma_1 & & \\ & \ddots & \\ & & \sigma_k \end{array}} & 0_{k \times (n-k)} \\ \hline 0_{(m-k) \times k} & 0_{(m-k) \times (n-k)} \end{array} \right] \left[\begin{array}{c|c|c} - & v_1^T & - \\ & \vdots & \\ - & v_k^T & - \\ \hline - & v_{k+1}^T & - \\ & \vdots & \\ - & v_n^T & - \end{array} \right] \\
 &= \left[\begin{array}{c|c|c} | & & | \\ u_1 & \cdots & u_k \\ | & & | \end{array} \right] \left[\begin{array}{ccc} \sigma_1 & & \\ & \ddots & \\ & & \sigma_k \end{array} \right] \left[\begin{array}{c|c|c} - & v_1^T & - \\ & \vdots & \\ - & v_k^T & - \end{array} \right] = U_1 \Sigma_1 V_1^T \\
 &\quad m \times k \quad \quad k \times k \quad \quad k \times n
 \end{aligned}$$

Reduced singular value expansion

$$A = U_1 \Sigma_1 V_1^T = \begin{bmatrix} | & & | \\ u_1 & \cdots & u_k \\ | & & | \end{bmatrix} \begin{bmatrix} \sigma_1 & & \\ & \ddots & \\ & & \sigma_k \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ & \vdots & \\ - & v_k^T & - \end{bmatrix}$$


$$= \sigma_1 u_1 v_1^T + \cdots + \sigma_k u_k v_k^T$$

Image compression using SVD




- lab: example_cameraman_image_compression.m

Principal Component Analysis (PCA)



- Data for height and weight
- What is the principal axis?
- How to divide into two groups?

Principal Component Analysis (PCA)




$$X =$$

| height (cm) | weight (kg) |
|-------------|-------------|
| 170 | 70 |
| 150 | 45 |
| 160 | 55 |
| 180 | 60 |
| 172 | 80 |
| 100 | 17 |
| 110 | 20 |
| 90 | 20 |
| 105 | 18 |
| 88 | 21 |

$$X_c = X - \text{mean}(X)$$

$$Y = X_c V \quad \rightarrow \text{Covariance matrix of } Y \text{ should be diagonal.}$$

Principal Component Analysis (PCA)



$$Y^T Y = \begin{bmatrix} \text{var}(y_1) & 0 \\ 0 & \text{var}(y_2) \end{bmatrix}$$



$$Y^T Y = V^T X_c^T X_c V = D$$

$$X_c^T X_c = V D V^T$$

$$X_c = U \Sigma V$$


- Column vectors of V are principal axes.
- lab: example_PCA_simple.m

Principal Component Analysis (PCA)




- 300 fruit images are given.
- Each image is 100 x 100.
- How to divide the fruit images into three groups?

Principal Component Analysis (PCA)





$$X_c = U \Sigma V^T$$

300 × 300 300 × 300 300 × 10000




Principal Component Analysis (PCA)



- lab: example_PCA_fruit.m

3.5 Pseudoinverse

Inverse of a *non-square* matrix



- Let A be an $m \times n$ matrix A with full row rank ($m < n$).
 - Is there an $n \times m$ matrix B such that $BA = I_n$?
 - Is there an $n \times m$ matrix B such that $AB = I_m$?

Inverse of a *non-square* matrix

$$\begin{array}{ccc}
 [x]_S & \xrightarrow{[T]_S = A} & [T(x)]_S \\
 \downarrow V^T & & \uparrow U \\
 [x]_V & \xrightarrow{[T]_V^U = \Sigma} & [T(x)]_U
 \end{array}$$

$$A = U \Sigma V^T$$

$\begin{matrix} \nearrow & \uparrow & \nwarrow \\ m \times m & m \times n & n \times n \end{matrix}$

- Reversing the direction does not work because Σ is not square.

$$\begin{array}{ccc}
 [x]_S & \xrightarrow{[T]_S = A} & [T(x)]_S \\
 \downarrow V_1^T & & \uparrow U_1 \\
 [x]_V & \xrightarrow{[T]_V^U = \Sigma_1} & [T(x)]_U
 \end{array}$$

$$A = U_1 \Sigma_1 V_1^T$$

$\begin{matrix} \nearrow & \uparrow & \nwarrow \\ m \times n & n \times n & n \times n \end{matrix}$

- Reversing the direction *works* for the *reduced* SVD because A has full row rank.


Inverse of a *non-square* matrix

- Let A be an $m \times n$ matrix, $A = U_1 \Sigma_1 V_1^T$ be the SVD of A , $A^+ = V_1 \Sigma_1^{-1} U_1^T$.
- If A is full row rank, $AA^+ = I_m$, which is called the right inverse of A .
- If A is full column rank, $A^+A = I_n$, which is called the left inverse of A .
- [Thm] If A is an $m \times n$ matrix with full column rank, then
$$A^+ = (A^T A)^{-1} A^T$$
- If A has full row rank, then
$$A^+ = A^T (AA^T)^{-1}$$

Properties of pseudoinverse


- If A^+ is the pseudoinverse of an $m \times n$ matrix A , then:
 - (a) $AA^+A = A$
 - (b) $A^+AA^+ = A^+$
 - (c) $(AA^+)^T = AA^+$
 - (d) $(A^+A)^T = A^+A$
 - (e) $(A^T)^+ = (A^+)^T$
 - (f) $A^{++} = A$

Pseudoinverse solves the Least Square Prob.



$$\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \end{bmatrix} + \begin{bmatrix} e_1 \\ e_2 \\ e_3 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

$$A \quad x \quad + \quad e \quad = \quad y$$



$$\text{proj}_{\text{col}(A)} y = A(A^T A)^{-1} A^T y$$


$$\hat{y} = Ax$$

$$x = (A^T A)^{-1} A^T y = A^+ y$$

- $AA^+ x$ is the projection of x onto $\text{col}(A)$.
- Best fit is projecting y onto $\text{col}(A)$.

- lab: example_least_square_is_projection.m

Condition number



- What matters is Σ which stretches/compresses a vector in each direction.
- [Def] **Condition number** of a matrix A is the ratio of the largest singular value of A to the smallest singular value of A .


$$\text{cond}(A) = \frac{\sigma_1}{\sigma_k}$$

- Condition number of A is too large.
 - \equiv Some dimension of $\text{Im}(A)$ is negligible.
 - $\equiv A$ is near-singular.



**Multibody vibration problem
is an eigenvalue problem.**

Compliant mechanisms



$$\delta_x = \frac{PL}{EA}$$


$$\delta_y = \frac{QL^3}{3EI} + \frac{ML^2}{2EI}$$

$$\theta_z = \frac{QL^2}{2EI} + \frac{ML}{EI}$$


$$\begin{bmatrix} \delta_x \\ \delta_y \\ \theta_z \end{bmatrix} = \begin{bmatrix} \frac{L}{EA} & 0 & 0 \\ 0 & \frac{L^3}{3EI} & \frac{L^2}{2EI} \\ 0 & \frac{L^2}{2EI} & \frac{L}{EI} \end{bmatrix} \begin{bmatrix} P \\ Q \\ M \end{bmatrix}$$

$$x = Cf \quad f = Kx$$

2-DOF system



(a)




$$m_1 \ddot{x}_1(t) + (k_1 + k_2)x_1(t) - k_2x_2(t) = 0$$

$$x_1(t) = X_1 \cos(\omega t + \phi)$$

$$m_2 \ddot{x}_2(t) - k_2x_1(t) + (k_2 + k_3)x_2(t) = 0$$

$$x_2(t) = X_2 \cos(\omega t + \phi)$$

2-DOF system



$$M\ddot{x} + Kx = 0$$

$$(K - \omega^2 M)x = 0$$


$$KM^{-1}x = \omega^2 x$$

```
[V, D] = eig(K*inv(M));
```

```
[V, D] = eig(inv(sqrt(M))*K*inv(sqrt(M)));
```

```
[V, D] = eig(K, M);
```

Multi-mass Compliant Mechanism



$$M\ddot{x} + Kx = 0$$

```
[V, D] = eig(K*inv(M));
```

```
[V, D] = eig(inv(sqrt(M))*K*inv(sqrt(M)));
```

```
[V, D] = eig(K, M);
```

See also.

<https://ocw.snu.ac.kr/node/30658>