

Part 1: Face - bones, muscles and vessels

133 questions in 3 sections. Answers and short explanations are at the end.

Section	Questions
Osteology of the face and skull	1 to 54
Muscles of the face, mastication and the TMJ	55 to 95
Blood supply and lymphatics of the face	96 to 133

How to use these questions. Each question has four statements and **only one** is correct, as in the real exam. Some questions ask for the statement that is FALSE (marked EXCEPT); read the stem carefully. Pick one answer, then check the key. The questions follow the lecture slides; where a slide and the textbooks disagree, the explanation says so.

Osteology of the face and skull

1. How many bones make up the skull?

- ☐ A. 20: 8 cranial and 12 facial
- ☐ B. 24: 10 cranial and 14 facial
- ☐ C. 22: 8 cranial and 14 facial
- ☐ D. 22: 14 cranial and 8 facial

2. Which of the following is a bone of the facial skeleton, not of the neurocranium?

- ☐ A. Ethmoid
- ☐ B. Vomer
- ☐ C. Temporal
- ☐ D. Sphenoid

3. Which of the following is a PAIRED facial bone?

- ☐ A. Vomer
- ☐ B. Mandible
- ☐ C. Ethmoid
- ☐ D. Inferior nasal concha

4. The unpaired facial bones are:

- ☐ A. Mandible and vomer
- ☐ B. Vomer and sphenoid
- ☐ C. Ethmoid and sphenoid
- ☐ D. Mandible and ethmoid

5. The neurocranium: all of the following are correct EXCEPT:

- ☐ A. Contains 8 bones
- ☐ B. Is divided into the calvaria and the base
- ☐ C. Includes the zygomatic bone
- ☐ D. Includes the frontal bone

6. The joint responsible for the 'no' (rotation) movement of the head is:

- ☐ A. Atlanto-axial joint
- ☐ B. Atlanto-occipital joint
- ☐ C. Temporomandibular joint
- ☐ D. Intervertebral joint C2-C3

7. The sella turcica is found on:

- ☐ A. The petrous part of the temporal bone
- ☐ B. The greater wing of the sphenoid
- ☐ C. The cribriform plate of the ethmoid
- ☐ D. The upper surface of the body of the sphenoid

8. Which opening is NOT in the greater wing of the sphenoid?

- ☐ A. Foramen spinosum
- ☐ B. Optic canal
- ☐ C. Foramen rotundum
- ☐ D. Foramen ovale

9. Match: the foramen rotundum transmits:

- ☐ A. V3 (mandibular nerve)
- ☐ B. V2 (maxillary nerve)
- ☐ C. V1 (ophthalmic nerve)
- ☐ D. The middle meningeal artery

10. The foramen ovale transmits:

- ☐ A. The maxillary nerve
- ☐ B. The mandibular nerve (V3)
- ☐ C. The middle meningeal artery
- ☐ D. The optic nerve

11. The foramen spinosum transmits:

- ☐ A. The facial nerve
- ☐ B. The internal carotid artery
- ☐ C. The middle meningeal vessels
- ☐ D. The mandibular nerve

12. The structures that separate the anterior from the middle cranial fossa are:

- ☐ A. The lesser wings of the sphenoid
- ☐ B. The petrous parts of the temporal bones
- ☐ C. The crista galli
- ☐ D. The greater wings of the sphenoid

13. The superior orbital fissure lies:

- ☐ A. Between the lesser and greater wings of the sphenoid
- ☐ B. In the frontal bone
- ☐ C. Between the greater wing and the maxilla
- ☐ D. In the ethmoid bone

14. The pterygoid processes belong to the:

- ☐ A. Temporal bone
- ☐ B. Maxilla
- ☐ C. Sphenoid
- ☐ D. Palatine bone

15. The cribriform plate of the ethmoid:

- ☐ A. Transmits the olfactory nerve fibres
- ☐ B. Forms the upper part of the nasal septum
- ☐ C. Forms the floor of the orbit
- ☐ D. Is pierced by the optic nerve

16. The crista galli gives attachment to:

- ☐ A. The falx cerebelli
- ☐ B. The diaphragma sellae
- ☐ C. The tentorium cerebelli
- ☐ D. The falx cerebri

17. Which nasal concha is NOT part of the ethmoid?

- ☐ A. Inferior nasal concha
- ☐ B. Superior nasal concha
- ☐ C. Middle nasal concha
- ☐ D. Supreme nasal concha

18. The bony nasal septum is formed by:

- ☐ A. The perpendicular plate of the ethmoid and the vomer
- ☐ B. The nasal bones and the lacrimal bones
- ☐ C. The vomer and the inferior nasal conchae
- ☐ D. The palatine bones and the sphenoid

19. The temporalis muscle arises from the:

- ☐ A. Squamous part of the temporal bone (temporal fossa)
- ☐ B. Petrous part of the temporal bone
- ☐ C. Mastoid process
- ☐ D. Styloid process

20. The internal acoustic meatus is found in the:

- ☐ A. Occipital bone
- ☐ B. Petrous part of the temporal bone
- ☐ C. Squamous part of the temporal bone
- ☐ D. Greater wing of the sphenoid

21. Sternocleidomastoid attaches to the:

- ☐ A. Mastoid process
- ☐ B. Zygomatic process
- ☐ C. Styloid process
- ☐ D. Pterygoid process

22. Which muscle does NOT attach to the styloid process?

- ☐ A. Stylohyoid
- ☐ B. Stylopharyngeus
- ☐ C. Styloglossus
- ☐ D. Masseter

23. The stylomastoid foramen transmits:

- ☐ A. The glossopharyngeal nerve
- ☐ B. The internal carotid artery
- ☐ C. The vestibulocochlear nerve
- ☐ D. The facial nerve

24. The head of the mandible articulates with the:

- ☐ A. Occipital condyle
- ☐ B. Mandibular fossa of the temporal bone
- ☐ C. Zygomatic bone
- ☐ D. Greater wing of the sphenoid

25. The pterion overlies the:

- ☐ A. Internal carotid artery
- ☐ B. Superficial temporal artery
- ☐ C. Sigmoid sinus
- ☐ D. Anterior branch of the middle meningeal artery

26. A young man is hit on the side of the head with a bat, has a lucid interval, then deteriorates. The structure most likely torn is:

- ☐ A. Superficial temporal vein
- ☐ B. Internal carotid artery
- ☐ C. Anterior branch of the middle meningeal artery
- ☐ D. Facial artery

27. Which bone does the zygomatic bone NOT articulate with?

- ☐ A. Frontal
- ☐ B. Maxilla
- ☐ C. Mandible
- ☐ D. Temporal

28. Which is NOT a process of the zygomatic bone?

- ☐ A. Frontal process
- ☐ B. Alveolar process
- ☐ C. Temporal process
- ☐ D. Maxillary process

29. The zygomatic arch is formed by:

- ☐ A. The zygomatic process of the temporal bone and the temporal process of the zygomatic bone
- ☐ B. The zygomatic bone only
- ☐ C. The frontal process of the zygomatic bone and the maxilla
- ☐ D. Mainly the maxilla

30. The zygomatic nerve passes through foramina in the:

- ☐ A. Zygomatic bone
- ☐ B. Maxilla
- ☐ C. Ethmoid
- ☐ D. Frontal bone

31. Which muscle attaches to the zygomatic bone?

- ☐ A. Zygomaticus major
- ☐ B. Mentalis
- ☐ C. Buccinator
- ☐ D. Temporalis

32. The largest paranasal sinus is in the:

- ☐ A. Sphenoid
- ☐ B. Maxilla
- ☐ C. Frontal bone
- ☐ D. Ethmoid

33. Which is NOT a process of the maxilla?

- ☐ A. Alveolar
- ☐ B. Frontal
- ☐ C. Zygomatic
- ☐ D. Pterygoid

34. The upper teeth are held by the:

- ☐ A. Palatine process of the maxilla
- ☐ B. Frontal process of the maxilla
- ☐ C. Zygomatic process of the maxilla
- ☐ D. Alveolar process of the maxilla

35. The hard palate is formed by:

- ☐ A. Pterygoid processes and maxillae
- ☐ B. Palatine processes of the maxillae and horizontal plates of the palatine bones
- ☐ C. Vomer and palatine bones
- ☐ D. Horizontal plates of the palatine bones only

36. The infraorbital nerve emerges onto the face through the:

- ☐ A. Zygomaticofacial foramen
- ☐ B. Supraorbital notch
- ☐ C. Infraorbital foramen of the maxilla
- ☐ D. Mental foramen

37. The palatine bone: all of the following are correct EXCEPT:

- ☐ A. Forms the roof of the nasal cavity
- ☐ B. Has a perpendicular plate in the lateral nasal wall
- ☐ C. Is L-shaped
- ☐ D. Has a horizontal plate forming the posterior hard palate

38. The vomer forms:

- ☐ A. The lower and posterior part of the bony nasal septum
- ☐ B. The floor of the orbit
- ☐ C. The roof of the nasal cavity
- ☐ D. The lateral wall of the nose

39. The lacrimal bone lies in the:

- ☐ A. Medial wall of the orbit
- ☐ B. Lateral wall of the orbit
- ☐ C. Floor of the orbit
- ☐ D. Roof of the orbit

40. The bridge of the nose is formed by the:

- ☐ A. Nasal bones
- ☐ B. Vomer
- ☐ C. Inferior conchae
- ☐ D. Lacrimal bones

41. The only movable bone of the skull is the:

- ☐ A. Maxilla
- ☐ B. Zygomatic bone
- ☐ C. Mandible
- ☐ D. Hyoid

42. The mental foramen is located:

- ☐ A. Below the orbit
- ☐ B. On the inner surface of the ramus
- ☐ C. On the coronoid process
- ☐ D. On the outer surface of the mandibular body below the premolars

43. The inferior alveolar nerve enters the mandible through the:

- ☐ A. Mandibular notch
- ☐ B. Mandibular foramen
- ☐ C. Foramen ovale
- ☐ D. Mental foramen

44. The mandibular notch lies between the:

- ☐ A. Angle and the body
- ☐ B. Coronoid process and condylar process
- ☐ C. Mental protuberance and the symphysis
- ☐ D. Two mylohyoid lines

45. The temporalis inserts on the:

- ☐ A. Mental protuberance
- ☐ B. Condylar process
- ☐ C. Angle of the mandible
- ☐ D. Coronoid process

46. The submandibular gland lies in a fossa on the mandible that is:

- ☐ A. Above the mylohyoid line
- ☐ B. On the coronoid process
- ☐ C. Below the mylohyoid line
- ☐ D. On the outer surface below the premolars

47. Which bone does NOT take part in the floor (base) of the cranial cavity?

- ☐ A. Ethmoid
- ☐ B. Sphenoid
- ☐ C. Parietal
- ☐ D. Occipital

48. The foramen ovale lies in the:

- ☐ A. Posterior cranial fossa
- ☐ B. Anterior cranial fossa
- ☐ C. Middle cranial fossa
- ☐ D. Orbit

49. The jugular foramen lies in the:

- ☐ A. Pterygopalatine fossa
- ☐ B. Anterior cranial fossa
- ☐ C. Posterior cranial fossa
- ☐ D. Middle cranial fossa

50. Which opening is NOT in the posterior cranial fossa?

- ☐ A. Hypoglossal canal
- ☐ B. Foramen magnum
- ☐ C. Foramen ovale
- ☐ D. Jugular foramen

51. The paranasal sinuses (from the osteology lecture): all of the following are correct EXCEPT:

- ☐ A. Are four pairs: frontal, ethmoidal, sphenoid, maxillary
- ☐ B. The maxilla is the only facial bone that contains a sinus
- ☐ C. The maxillary sinus is the largest
- ☐ D. The sphenoid sinus is the largest

52. Which is NOT a proposed function of the paranasal sinuses?

- ☐ A. Giving resonance to the voice
- ☐ B. Lightening the skull
- ☐ C. Producing cerebrospinal fluid
- ☐ D. Humidifying inspired air

53. The pituitary gland lies in the:

- ☐ A. Cribriform plate
- ☐ B. Foramen magnum
- ☐ C. Petrous temporal bone
- ☐ D. Sella turcica of the sphenoid

54. The foramen magnum lies in the:

- ☐ A. Sphenoid
- ☐ B. Parietal bone
- ☐ C. Temporal bone
- ☐ D. Occipital bone

Muscles of the face, mastication and the TMJ

55. Muscles of facial expression: all of the following are correct EXCEPT:

- ☐ A. Are subcutaneous, lying in the superficial fascia
- ☐ B. Insert into the skin
- ☐ C. Are supplied by the facial nerve (CN VII)
- ☐ D. Are supplied by the mandibular nerve

56. Muscles of facial expression act on the openings of the face as:

- ☐ A. Sphincters and dilators
- ☐ B. Constrictors of the pharynx
- ☐ C. Protractors of the tongue
- ☐ D. Elevators of the mandible

57. Which muscle near the eye is NOT supplied by the facial nerve?

- ☐ A. Levator palpebrae superioris
- ☐ B. Frontalis
- ☐ C. Corrugator supercilii
- ☐ D. Orbicularis oculi

58. The two bellies of occipitofrontalis are joined by the:

- ☐ A. Superficial temporal fascia
- ☐ B. Platysma
- ☐ C. Galea aponeurotica
- ☐ D. Temporal fascia

59. The frontal belly of occipitofrontalis:

- ☐ A. Closes the eye
- ☐ B. Raises the eyebrows and wrinkles the forehead
- ☐ C. Has a bony origin on the frontal bone
- ☐ D. Draws the scalp backward

60. Gentle, involuntary closure of the eyelids (blinking) is produced by:

- ☐ A. Corrugator supercilii
- ☐ B. Orbital part of orbicularis oculi
- ☐ C. Levator palpebrae superioris
- ☐ D. Palpebral part of orbicularis oculi

61. Forceful, voluntary closure of the eye is produced by:

- ☐ A. Superior tarsal muscle
- ☐ B. Procerus
- ☐ C. Palpebral part of orbicularis oculi
- ☐ D. Orbital part of orbicularis oculi

62. The 'frowning muscle' is:

- ☐ A. Risorius
- ☐ B. Mentalis
- ☐ C. Procerus
- ☐ D. Corrugator supercilii

63. Transverse wrinkles over the bridge of the nose are made by:

- ☐ A. Procerus
- ☐ B. Nasalis, transverse part
- ☐ C. Corrugator supercilii
- ☐ D. Depressor septi

64. Which is NOT a muscle of the nose?

- ☐ A. Buccinator
- ☐ B. Nasalis
- ☐ C. Depressor septi
- ☐ D. Procerus

65. The alar part of nasalis:

- ☐ A. Elevates the upper lip
- ☐ B. Opens the nostril
- ☐ C. Depresses the septum
- ☐ D. Compresses the nasal aperture

66. Narrowing and closing the mouth, as in whistling, is the action of:

- ☐ A. Risorius
- ☐ B. Levator anguli oris
- ☐ C. Orbicularis oris
- ☐ D. Zygomaticus major

67. The buccinator is supplied by:

- ☐ A. The hypoglossal nerve
- ☐ B. The glossopharyngeal nerve
- ☐ C. The facial nerve
- ☐ D. The buccal nerve of V3

68. The pterygomandibular raphe separates the buccinator from:

- ☐ A. The orbicularis oris
- ☐ B. The masseter
- ☐ C. The medial pterygoid
- ☐ D. The superior constrictor of the pharynx

69. Keeping food between the teeth during chewing is a function of:

- ☐ A. Platysma
- ☐ B. Buccinator
- ☐ C. Orbicularis oris
- ☐ D. Masseter

70. The smiling muscle that elevates the angle of the mouth from the zygomatic bone is:

- ☐ A. Zygomaticus major
- ☐ B. Risorius
- ☐ C. Depressor anguli oris
- ☐ D. Mentalis

71. The muscle that arises from the masseteric fascia and pulls the angle of the mouth laterally is:

- ☐ A. Zygomaticus minor
- ☐ B. Risorius
- ☐ C. Levator anguli oris
- ☐ D. Buccinator

72. Levator labii superioris alaeque nasi arises from:

- ☐ A. The mandible
- ☐ B. The frontal process of the maxilla
- ☐ C. The nasal bone
- ☐ D. The zygomatic arch

73. Levator anguli oris arises from:

- ☐ A. The masseteric fascia
- ☐ B. The zygomatic arch
- ☐ C. The mandible
- ☐ D. The canine fossa of the maxilla

74. The 'pouting muscle' is:

- ☐ A. Risorius
- ☐ B. Procerus
- ☐ C. Corrugator supercilii
- ☐ D. Mentalis

75. Which muscle belongs to the UPPER group of mouth muscles?

- ☐ A. Mentalis
- ☐ B. Levator anguli oris
- ☐ C. Depressor labii inferioris
- ☐ D. Depressor anguli oris

76. Platysma is supplied by the:

- ☐ A. Mandibular nerve
- ☐ B. Marginal mandibular branch
- ☐ C. Cervical branch of the facial nerve
- ☐ D. Ansa cervicalis

77. Which nerve is NOT a terminal branch of the facial nerve in the face?

- ☐ A. Auriculotemporal
- ☐ B. Marginal mandibular
- ☐ C. Buccal
- ☐ D. Temporal

78. Mentalis is supplied by which branch of the facial nerve?

- ☐ A. Zygomatic
- ☐ B. Marginal mandibular
- ☐ C. Cervical
- ☐ D. Temporal

79. A patient cannot close the right eye, cannot raise the right eyebrow and the right corner of the mouth droops. The lesion is:

- ☐ A. Right lower motor neuron facial palsy (e.g. Bell's palsy)
- ☐ B. Right trigeminal nerve lesion
- ☐ C. Left upper motor neuron lesion
- ☐ D. Right oculomotor palsy

80. The muscles of mastication are supplied by the:

- ☐ A. Mandibular nerve (V3)
- ☐ B. Hypoglossal nerve
- ☐ C. Maxillary nerve (V2)
- ☐ D. Facial nerve

81. Temporalis: all of the following are correct EXCEPT:

- ☐ A. Its posterior fibres retract the mandible
- ☐ B. Inserts on the coronoid process and anterior border of the ramus
- ☐ C. Elevates the mandible
- ☐ D. Inserts on the angle of the mandible

82. The masseter arises from the:

- ☐ A. Lateral pterygoid plate
- ☐ B. Maxillary tuberosity
- ☐ C. Temporal fossa
- ☐ D. Zygomatic arch

83. The lateral pterygoid inserts on the:

- ☐ A. Medial surface of the ramus
- ☐ B. Coronoid process
- ☐ C. Pterygoid fovea of the neck, the TMJ capsule and disc
- ☐ D. Angle of the mandible

84. The only muscle of mastication that helps open the mouth is:

- ☐ A. Medial pterygoid
- ☐ B. Masseter
- ☐ C. Lateral pterygoid
- ☐ D. Temporalis

85. The medial pterygoid inserts on the:

- ☐ A. Pterygoid fovea
- ☐ B. Medial surface of the ramus
- ☐ C. Articular disc
- ☐ D. Coronoid process

86. Actions of the medial pterygoid: all of the following are correct EXCEPT:

- ☐ A. Elevates the mandible
- ☐ B. Depresses the mandible against resistance
- ☐ C. Grinding when acting alternately
- ☐ D. Protrudes the mandible (both sides)

87. Which muscle does NOT elevate the mandible?

- ☐ A. Temporalis
- ☐ B. Masseter
- ☐ C. Medial pterygoid
- ☐ D. Lateral pterygoid

88. Depression of the mandible (opening the mouth) is mostly caused by:

- ☐ A. Masseter
- ☐ B. Medial pterygoid
- ☐ C. Temporalis
- ☐ D. Gravity

89. The articular surfaces of the TMJ are covered by:

- ☐ A. Elastic cartilage
- ☐ B. Hyaline cartilage
- ☐ C. Fibrocartilage
- ☐ D. Synovial membrane only

90. Which is NOT a ligament of the TMJ?

- ☐ A. Pterygomandibular raphe
- ☐ B. Stylomandibular ligament
- ☐ C. Sphenomandibular ligament
- ☐ D. Lateral ligament

91. The lateral (temporomandibular) ligament prevents:

- ☐ A. Lateral grinding
- ☐ B. Posterior dislocation
- ☐ C. Anterior dislocation
- ☐ D. Elevation

92. Protrusion and retraction of the mandible take place in:

- ☐ A. The atlanto-axial joint
- ☐ B. The lower cavity of the TMJ
- ☐ C. The symphysis menti
- ☐ D. The upper cavity of the TMJ

93. TMJ dislocation is usually:

- ☐ A. Medial
- ☐ B. Anterior, with the mouth stuck open
- ☐ C. Posterior, with the mouth closed
- ☐ D. Superior into the cranial fossa

94. Relations of the TMJ: all of the following are correct EXCEPT:

- ☐ A. Laterally: parotid gland, fascia and skin
- ☐ B. Superiorly: middle cranial fossa with the temporal lobe
- ☐ C. Medially: maxillary artery and vein and auriculotemporal nerve
- ☐ D. Posteriorly: the coronoid process

95. Which nerve is at risk in a high-energy TMJ injury?

- ☐ A. Auriculotemporal nerve
- ☐ B. Hypoglossal nerve
- ☐ C. Lingual nerve
- ☐ D. Optic nerve

Blood supply and lymphatics of the face

96. The right common carotid artery arises from:

- ☐ A. The subclavian artery
- ☐ B. The arch of the aorta directly
- ☐ C. The brachiocephalic trunk
- ☐ D. The external carotid

97. The common carotid artery bifurcates at the level of:

- ☐ A. The angle of the mandible (C2)
- ☐ B. The upper border of the thyroid cartilage (C3-C4)
- ☐ C. The sternoclavicular joint
- ☐ D. The cricoid cartilage (C6)

98. At its origin, the external carotid artery lies:

- ☐ A. Anteromedial to the internal carotid
- ☐ B. Posterolateral to the internal carotid
- ☐ C. Behind the internal jugular vein
- ☐ D. Inside the parotid gland

99. The external carotid artery ends:

- ☐ A. In the carotid triangle as the facial and lingual arteries
- ☐ B. In the parotid gland as the superficial temporal and maxillary arteries
- ☐ C. At the foramen spinosum
- ☐ D. At the angle of the mandible as the occipital artery

100. Which structure passes between the external and internal carotid arteries?

- ☐ A. Hypoglossal nerve
- ☐ B. Facial nerve
- ☐ C. Internal jugular vein
- ☐ D. Glossopharyngeal nerve

101. Within the parotid gland the external carotid artery is crossed by the:

- ☐ A. Phrenic nerve
- ☐ B. Facial nerve
- ☐ C. Vagus nerve
- ☐ D. Lingual nerve

102. Which artery is NOT a branch of the external carotid?

- ☐ A. Superior thyroid
- ☐ B. Ophthalmic
- ☐ C. Facial
- ☐ D. Lingual

103. The smallest branch of the external carotid is:

- ☐ A. Ascending pharyngeal
- ☐ B. Superior thyroid
- ☐ C. Posterior auricular
- ☐ D. Occipital

104. The terminal branches of the external carotid artery are:

- ☐ A. Facial and lingual
- ☐ B. Maxillary and facial
- ☐ C. Occipital and posterior auricular
- ☐ D. Superficial temporal and maxillary

105. The superior laryngeal artery is a branch of the:

- ☐ A. Facial artery
- ☐ B. Superior thyroid artery
- ☐ C. Inferior thyroid artery
- ☐ D. Lingual artery

106. The lingual artery is divided into three parts by the:

- ☐ A. Digastric
- ☐ B. Genioglossus
- ☐ C. Hyoglossus
- ☐ D. Mylohyoid

107. Which is NOT a branch of the lingual artery?

- ☐ A. Deep lingual
- ☐ B. Superior laryngeal
- ☐ C. Suprahyoid
- ☐ D. Dorsal lingual

108. The facial artery: all of the following are correct EXCEPT:

- ☐ A. Arises in the carotid triangle at the level of the hyoid
- ☐ B. Is closely related to the submandibular gland
- ☐ C. Ends as the superficial temporal artery
- ☐ D. Crosses the mandible at the anterior border of the masseter

109. Which is a CERVICAL branch of the facial artery?

- ☐ A. Lateral nasal
- ☐ B. Angular
- ☐ C. Superior labial
- ☐ D. Submental

110. The termination of the facial artery is the:

- ☐ A. Superior labial artery
- ☐ B. Angular artery
- ☐ C. Lateral nasal artery
- ☐ D. Submental artery

111. The occipital artery: all of the following are correct EXCEPT:

- ☐ A. Runs under the posterior belly of digastric
- ☐ B. Is a terminal branch of the ECA
- ☐ C. Supplies sternocleidomastoid and the posterior scalp
- ☐ D. Gives meningeal branches to the dura

112. The smallest branch of the external carotid gives the inferior tympanic artery. It is the:

- ☐ A. Occipital artery
- ☐ B. Posterior auricular artery
- ☐ C. Superior thyroid artery
- ☐ D. Ascending pharyngeal artery

113. The stylomastoid artery, entering the stylomastoid foramen, is a branch of the:

- ☐ A. Occipital artery
- ☐ B. Superficial temporal artery
- ☐ C. Posterior auricular artery
- ☐ D. Maxillary artery

114. The transverse facial artery is a branch of the:

- ☐ A. Lingual artery
- ☐ B. Maxillary artery
- ☐ C. Superficial temporal artery
- ☐ D. Facial artery

115. The maxillary artery: all of the following are correct EXCEPT:

- ☐ A. Arises behind the neck of the mandible in the parotid
- ☐ B. Is the larger terminal branch of the ECA
- ☐ C. Enters the skull through the foramen ovale
- ☐ D. Is divided into three parts by the lateral pterygoid

116. Which structure is NOT supplied by the maxillary artery?

- ☐ A. Upper and lower jaws
- ☐ B. Meninges
- ☐ C. Muscles of mastication
- ☐ D. Thyroid gland

117. The middle meningeal artery enters the skull through the:

- ☐ A. Foramen rotundum
- ☐ B. Jugular foramen
- ☐ C. Foramen ovale
- ☐ D. Foramen spinosum

118. Which artery supplies the jaw, muscles of mastication and meninges?

- ☐ A. Maxillary
- ☐ B. Facial
- ☐ C. Superficial temporal
- ☐ D. Superior thyroid

119. The angular vein is formed by:

- ☐ A. Facial and retromandibular veins
- ☐ B. Supratrochlear and supraorbital veins
- ☐ C. Lingual and pharyngeal veins
- ☐ D. Superficial temporal and maxillary veins

120. The common facial vein drains into the:

- ☐ A. External jugular vein
- ☐ B. Subclavian vein
- ☐ C. Pterygoid plexus
- ☐ D. Internal jugular vein

121. The retromandibular vein: all of the following are correct EXCEPT:

- ☐ A. Its anterior division joins the facial vein
- ☐ B. Its posterior division joins the posterior auricular vein to form the external jugular
- ☐ C. Is formed by the superficial temporal and maxillary veins
- ☐ D. Is formed by the facial and angular veins

122. The maxillary vein is formed from:

- ☐ A. The angular vein
- ☐ B. The facial vein
- ☐ C. The pterygoid venous plexus
- ☐ D. The cavernous sinus directly

123. The external jugular vein drains into the:

- ☐ A. Sigmoid sinus
- ☐ B. Internal jugular vein
- ☐ C. Brachiocephalic vein directly
- ☐ D. Subclavian vein

124. The internal jugular vein: all of the following are correct EXCEPT:

- ☐ A. Joins the subclavian vein to form the brachiocephalic vein
- ☐ B. Continues the cavernous sinus
- ☐ C. Receives the facial, lingual and superior and middle thyroid veins
- ☐ D. Leaves the skull through the jugular foramen

125. A facial infection can spread to the cavernous sinus through:

- ☐ A. The ophthalmic veins and the pterygoid plexus
- ☐ B. The lymphatics of the submental nodes
- ☐ C. The facial artery
- ☐ D. The external jugular vein

126. A patient squeezes a boil beside the nose and days later develops proptosis, ophthalmoplegia and fever. The most likely complication is:

- ☐ A. Cavernous sinus thrombosis
- ☐ B. Parotitis
- ☐ C. Maxillary sinusitis
- ☐ D. Bell's palsy

127. The deep facial vein connects the facial vein with the:

- ☐ A. Sigmoid sinus
- ☐ B. Superior sagittal sinus
- ☐ C. Pterygoid venous plexus
- ☐ D. Subclavian vein

128. Lymph from the forehead, lateral eyelids and conjunctiva drains first to:

- ☐ A. Occipital nodes
- ☐ B. Preauricular (superficial parotid) nodes
- ☐ C. Submental nodes
- ☐ D. Submandibular nodes

129. Lymph from the cheeks, upper lip and lateral nose drains first to:

- ☐ A. Submandibular nodes
- ☐ B. Preauricular nodes
- ☐ C. Supraclavicular nodes
- ☐ D. Submental nodes

130. Lymph from the lower lip, chin and tip of the tongue drains first to:

- ☐ A. Submandibular nodes
- ☐ B. Preauricular nodes
- ☐ C. Submental nodes
- ☐ D. Retropharyngeal nodes

131. Final lymphatic drainage of the face: all of the following are correct EXCEPT:

- ☐ **A.** Ends in the axillary nodes
- ☐ **B.** Through the deep and superficial cervical nodes
- ☐ **C.** Through the jugular trunks
- ☐ **D.** Into the right lymphatic duct on the right

132. Which statement is correct?

- ☐ **A.** The facial and occipital arteries are terminal branches of the ECA
- ☐ **B.** The jugulodigastric node is part of the deep cervical nodes
- ☐ **C.** The lingual artery is a branch of the internal carotid
- ☐ **D.** The retromandibular vein drains only into the subclavian vein

133. Within the parotid gland, from superficial to deep:

- ☐ **A.** External carotid artery, facial nerve, retromandibular vein
- ☐ **B.** Retromandibular vein, facial nerve, external carotid
- ☐ **C.** Facial nerve, external carotid, retromandibular vein
- ☐ **D.** Facial nerve, retromandibular vein, external carotid artery

Answer key

The single correct answer for each question, with a short explanation where it helps.

Osteology of the face and skull

1	C	Neurocranium 8 + viscerocranium 14.
2	B	Cranial: frontal, 2 parietal, 2 temporal, occipital, sphenoid, ethmoid.
3	D	Paired: maxilla, zygomatic, palatine, nasal, lacrimal, inferior nasal concha.
4	A	Ethmoid and sphenoid are cranial bones.
5	C	Neurocranium = calvaria (skull cap) + base; 8 bones. The zygomatic bone is facial.
6	A	Atlanto-occipital = nodding ('yes').
7	D	It houses the pituitary gland.
8	B	The optic canal is in the lesser wing.
9	B	'Standing Room Only': superior orbital fissure V1, rotundum V2, ovale V3.
10	B	Foramen ovale = V3. Spinosum = middle meningeal artery.
11	C	Foramen spinosum: middle meningeal artery and vein (the artery is a branch of the maxillary artery).
12	A	The petrous ridges separate the middle and posterior fossae.
13	A	Superior orbital fissure: between lesser and greater wings. Inferior orbital fissure: between greater wing and maxilla.
14	C	Each has a medial and lateral plate.
15	A	It forms the roof of the nasal cavity.
16	D	Upward midline projection of the ethmoid.
17	A	The inferior concha is a separate bone.
18	A	The septal cartilage completes it in front.
19	A	
20	B	
21	A	The mastoid lies behind the external acoustic meatus.
22	D	Styloid muscles: styloglossus, stylohyoid, stylopharyngeus.
23	D	CN VII leaves the skull through the stylomastoid foramen.
24	B	
25	D	Junction of frontal, parietal, squamous temporal and greater wing of sphenoid.
26	C	Fracture at the pterion tears the middle meningeal artery: extradural haematoma with a lucid interval.
27	C	It articulates with frontal, sphenoid, temporal and maxilla.
28	B	
29	A	Classic question, also on the 2026 mock.
30	A	Zygomatico-orbital, zygomaticofacial and zygomaticotemporal foramina.
31	A	Masseter and zygomaticus minor also attach to it.
32	B	
33	D	The pterygoid processes belong to the sphenoid; the maxilla has frontal, zygomatic, alveolar and palatine processes.

34	D	8 sockets on each side.
35	B	
36	C	
37	A	The roof of the nasal cavity is the cribriform plate.
38	A	It is unpaired.
39	A	
40	A	
41	C	
42	D	
43	B	Site of the inferior alveolar nerve block.
44	B	
45	D	
46	C	The sublingual fossa lies above the line.
47	C	Base: frontal, ethmoid, sphenoid, temporal, occipital.
48	C	
49	C	
50	C	The foramen ovale is in the middle cranial fossa.
51	D	The maxillary sinus is the largest.
52	C	
53	D	
54	D	

Muscles of the face, mastication and the TMJ

55	D	Mastication = V3; facial expression = VII.
56	A	
57	A	Levator palpebrae superioris = CN III (its smooth superior tarsal part is sympathetic).
58	C	
59	B	It arises from the galea; no bony attachment.
60	D	
61	D	
62	D	Corrugator draws the eyebrows medially and downward.
63	A	
64	A	
65	B	The transverse part compresses.
66	C	
67	C	The buccal nerve of V3 is only sensory to the cheek.
68	D	Buccinator in front, superior constrictor behind.
69	B	
70	A	
71	B	

72	B	
73	D	It raises the angle of the mouth while smiling.
74	D	
75	B	
76	C	
77	A	The auriculotemporal is a branch of V3.
78	B	
79	A	Whole half-face including forehead = LMN lesion of CN VII. An UMN (stroke) lesion spares the forehead.
80	A	
81	D	Temporalis inserts on the coronoid process; masseter on the angle.
82	D	It inserts on the angle and ramus up to the coronoid.
83	C	
84	C	
85	B	
86	B	
87	D	
88	D	Digastric, geniohyoid and mylohyoid assist against resistance.
89	C	
90	A	
91	B	
92	D	The lower cavity is the hinge (elevation/depression).
93	B	
94	D	Posteriorly: tympanic plate of the external acoustic meatus.
95	A	The facial nerve is also close to the joint.

Blood supply and lymphatics of the face

96	C	Left common carotid arises directly from the arch.
97	B	
98	A	It ends lateral to the ICA.
99	B	
100	D	Also styloglossus, stylopharyngeus and the pharyngeal branch of the vagus.
101	B	
102	B	The ophthalmic is the first branch of the internal carotid.
103	A	
104	D	
105	B	
106	C	
107	B	The superior laryngeal artery is a branch of the superior thyroid artery.
108	C	It ends as the angular artery.
109	D	Also ascending palatine, tonsillar and glandular.

110	B	
111	B	The terminal branches are the superficial temporal and maxillary.
112	D	
113	C	
114	C	
115	C	Its middle meningeal branch uses the foramen spinosum; the maxillary artery itself enters the pterygopalatine fossa.
116	D	
117	D	
118	A	Question from the Anatomy 3 mock test.
119	B	Formed at the medial angle of the eye; continues as the facial vein.
120	D	
121	D	It is formed by the superficial temporal and maxillary veins.
122	C	In the infratemporal fossa; joins the superficial temporal vein to form the retromandibular vein.
123	D	
124	B	It continues the sigmoid sinus.
125	A	
126	A	Spread through the facial/angular vein → ophthalmic veins → cavernous sinus (III, IV, VI, V1 affected).
127	C	
128	B	
129	A	
130	C	Then to the deep cervical chain.
131	A	Facial lymph ends in the cervical nodes and jugular trunks.
132	B	Adapted from the 2026 mock exam.
133	D	Classic ordering of the parotid contents.