

Part 2: Oral cavity, paranasal sinuses, orbit and orbital appendages

117 questions in 4 sections. Answers and short explanations are at the end.

Section	Questions
Oral cavity, tongue, palate and salivary glands	1 to 41
Nose and paranasal sinuses	42 to 67
The orbit: osteology and myology	68 to 92
Orbital appendages: eyelids, conjunctiva, lacrimal apparatus	93 to 117

How to use these questions. Each question has four statements and **only one** is correct, as in the real exam. Some questions ask for the statement that is FALSE (marked EXCEPT); read the stem carefully. Pick one answer, then check the key. The questions follow the lecture slides; where a slide and the textbooks disagree, the explanation says so.

Oral cavity, tongue, palate and salivary glands

1. Posteriorly, the oral cavity ends at:

- ☐ A. The palatopharyngeal folds
- ☐ B. The oropharyngeal isthmus (palatoglossal folds)
- ☐ C. The epiglottis
- ☐ D. The laryngeal inlet

2. With the jaws closed, the vestibule communicates with the oral cavity proper:

- ☐ A. Behind the third molars
- ☐ B. Through the oral fissure
- ☐ C. Through the sublingual papilla
- ☐ D. Between the central incisors

3. The lateral wall of the vestibule (the cheek) is formed by:

- ☐ A. Masseter
- ☐ B. Orbicularis oris
- ☐ C. Buccinator
- ☐ D. Mylohyoid

4. The parotid duct opens:

- ☐ A. Into the vestibule, opposite the upper second molar
- ☐ B. On the sublingual papilla
- ☐ C. Into the vestibule, opposite the lower second molar
- ☐ D. Along the sublingual fold

5. The floor of the oral cavity proper is formed mainly by:

- ☐ A. Masseter
- ☐ B. Mylohyoid
- ☐ C. Buccinator
- ☐ D. Hyoglossus

6. On the floor of the mouth: all of the following are correct EXCEPT:

- ☐ A. The sublingual gland raises the sublingual fold
- ☐ B. The submandibular duct opens on a papilla beside the frenulum
- ☐ C. The frenulum connects the tongue to the floor in the midline
- ☐ D. The parotid duct opens beside the frenulum

7. Sensation from the mucosa of the cheek is carried by:

- ☐ A. The greater palatine nerve
- ☐ B. The lingual nerve
- ☐ C. The buccal nerve (V3)
- ☐ D. The buccal branch of the facial nerve

8. Motor supply to buccinator and orbicularis oris is from:

- ☐ A. The glossopharyngeal nerve
- ☐ B. The mandibular nerve
- ☐ C. The facial nerve
- ☐ D. The hypoglossal nerve

9. The tongue: all of the following are correct EXCEPT:

- ☐ A. Is divided into halves by a median septum
- ☐ B. Is a mass of striated muscle covered with mucosa
- ☐ C. Is made of smooth muscle
- ☐ D. Has oral, pharyngeal and root parts

10. The sulcus terminalis:

- ☐ A. Divides the dorsum into anterior two thirds and posterior one third
- ☐ B. Lies on the ventral surface
- ☐ C. Has the foramen caecum at the tip of the tongue
- ☐ D. Has the vallate papillae behind it

11. The foramen caecum of the tongue marks:

- ☐ A. The centre of the lingual tonsil
- ☐ B. The opening of the submandibular duct
- ☐ C. The opening of the sublingual ducts
- ☐ D. The upper end of the thyroglossal duct

12. Which is NOT found on the anterior two thirds of the dorsum of the tongue?

- ☐ A. Fungiform papillae
- ☐ B. Vallate papillae
- ☐ C. Filiform papillae
- ☐ D. The lingual tonsil

13. Which is found on the ventral surface of the tongue?

- ☐ A. The lingual tonsil
- ☐ B. The plica fimbriata
- ☐ C. The foramen caecum
- ☐ D. The vallate papillae

14. The intrinsic muscles of the tongue:

- ☐ A. Move the tongue as a whole
- ☐ B. Alter the shape of the tongue
- ☐ C. Arise from the mandible
- ☐ D. Are supplied by the lingual nerve

15. Which is NOT an extrinsic muscle of the tongue?

- ☐ A. Styloglossus
- ☐ B. Hyoglossus
- ☐ C. Mylohyoid
- ☐ D. Genioglossus

16. Protrusion of the tongue is produced by:

- ☐ A. Styloglossus
- ☐ B. Genioglossus
- ☐ C. Palatoglossus
- ☐ D. Hyoglossus

17. Which muscle retracts the tongue?

- ☐ A. Styloglossus
- ☐ B. Geniohyoid
- ☐ C. Genioglossus
- ☐ D. Mylohyoid

18. Which muscle depresses the tongue?

- ☐ A. Levator veli palatini
- ☐ B. Hyoglossus
- ☐ C. Styloglossus
- ☐ D. Palatoglossus

19. General sensation of the anterior two thirds of the tongue is carried by:

- ☐ A. Hypoglossal (XII)
- ☐ B. Chorda tympani (VII)
- ☐ C. Lingual nerve (V3)
- ☐ D. Glossopharyngeal (IX)

20. Taste from the anterior two thirds of the tongue is carried by:

- ☐ A. Vagus
- ☐ B. Chorda tympani (VII)
- ☐ C. Lingual nerve (V3) fibres of its own
- ☐ D. Glossopharyngeal nerve

21. The posterior third of the tongue receives general sensation and taste from:

- ☐ A. Chorda tympani
- ☐ B. Lingual nerve
- ☐ C. Glossopharyngeal nerve (IX)
- ☐ D. Hypoglossal nerve

22. The base of the tongue (near the epiglottis) is supplied by:

- ☐ A. The internal laryngeal nerve (X)
- ☐ B. The lingual nerve
- ☐ C. The facial nerve
- ☐ D. The hypoglossal nerve

23. Which tongue muscle is NOT supplied by the hypoglossal nerve?

- ☐ A. Hyoglossus
- ☐ B. Styloglossus
- ☐ C. Palatoglossus
- ☐ D. Genioglossus

24. A patient protrudes the tongue and it deviates to the left. The lesion is in the:

- ☐ A. Right hypoglossal nerve
- ☐ B. Right facial nerve
- ☐ C. Left hypoglossal nerve
- ☐ D. Left lingual nerve

25. The main artery of the tongue is:

- ☐ A. The maxillary artery
- ☐ B. The lingual artery
- ☐ C. The superior thyroid artery
- ☐ D. The ophthalmic artery

26. Lymph from the tip of the tongue drains to:

- ☐ A. Submandibular nodes of the same side only
- ☐ B. Axillary nodes
- ☐ C. Submental nodes, on both sides
- ☐ D. Preauricular nodes

27. The hard palate is formed by:

- ☐ A. The pterygoid plates and the palatine bones
- ☐ B. The maxillae only
- ☐ C. The vomer and the palatine bones
- ☐ D. Palatine processes of the maxillae and horizontal plates of the palatine bones

28. The palatine aponeurosis is the expanded tendon of:

- ☐ A. Tensor veli palatini
- ☐ B. Levator veli palatini
- ☐ C. Musculus uvulae
- ☐ D. Palatoglossus

29. Tensor veli palatini is supplied by:

- ☐ A. The nerve to medial pterygoid (V3)
- ☐ B. The hypoglossal nerve
- ☐ C. The pharyngeal plexus
- ☐ D. The facial nerve

30. Which palatine muscle is NOT supplied by the pharyngeal plexus?

- ☐ A. Levator veli palatini
- ☐ B. Palatopharyngeus
- ☐ C. Tensor veli palatini
- ☐ D. Palatoglossus

31. Levator veli palatini: all of the following are correct EXCEPT:

- ☐ A. Arises from the petrous temporal bone and auditory tube
- ☐ B. Raises the soft palate
- ☐ C. Is supplied by V3
- ☐ D. Inserts into the palatine aponeurosis

32. Palatopharyngeus inserts on:

- ☐ A. The posterior border of the thyroid cartilage
- ☐ B. The spine of the sphenoid
- ☐ C. The pterygoid hamulus
- ☐ D. The body of the hyoid bone

33. Which is NOT one of the three major salivary glands?

- ☐ A. Sublingual
- ☐ B. Lacrimal
- ☐ C. Parotid
- ☐ D. Submandibular

34. The parotid gland is:

- ☐ A. Purely serous
- ☐ B. The smallest major salivary gland
- ☐ C. Mixed, with mucous units predominating
- ☐ D. Purely mucous

35. Which vein is formed within the parotid gland?

- ☐ A. The retromandibular vein
- ☐ B. The lingual vein
- ☐ C. The facial vein
- ☐ D. The internal jugular vein

36. Postganglionic secretomotor fibres reach the parotid in:

- ☐ A. The chorda tympani
- ☐ B. The lingual nerve
- ☐ C. The greater petrosal nerve
- ☐ D. The auriculotemporal nerve

37. The parotid duct:

- ☐ A. Opens opposite the lower second molar
- ☐ B. Pierces the buccinator
- ☐ C. Pierces the masseter
- ☐ D. Opens on the sublingual papilla

38. The submandibular gland is:

- ☐ A. Mixed, with mucous units predominating
- ☐ B. Purely mucous
- ☐ C. Mixed, with serous units predominating
- ☐ D. Purely serous

39. The submandibular duct opens:

- ☐ A. On the sublingual papilla beside the frenulum
- ☐ B. By many small ducts along the sublingual fold
- ☐ C. Into the vestibule
- ☐ D. Opposite the upper second molar

40. Secretomotor fibres to the submandibular gland synapse in:

- ☐ A. The pterygopalatine ganglion
- ☐ B. The otic ganglion
- ☐ C. The ciliary ganglion
- ☐ D. The submandibular ganglion

41. The sublingual gland: all of the following are correct EXCEPT:

- ☐ A. Drains by many small ducts of Rivinus along the sublingual fold
- ☐ B. Is purely serous
- ☐ C. Is the smallest major salivary gland
- ☐ D. Lies above the mylohyoid

Nose and paranasal sinuses

42. The paranasal sinuses:

- ☐ A. All drain into the nasal cavity
- ☐ B. Are fully developed at birth
- ☐ C. Are five in number
- ☐ D. Are lined by stratified squamous epithelium

43. The nasal conchae divide each nasal cavity into several spaces. Which is NOT one of them?

- ☐ A. Superior meatus
- ☐ B. Nasopharynx
- ☐ C. Middle meatus
- ☐ D. Inferior meatus

44. The middle nasal meatus lies:

- ☐ A. Between the superior and middle conchae
- ☐ B. Between the inferior concha and the floor
- ☐ C. Above the superior concha
- ☐ D. Under the middle concha (between middle and inferior conchae)

45. Which does NOT drain into the middle meatus?

- ☐ A. Anterior and middle ethmoidal cells
- ☐ B. Frontal sinus
- ☐ C. Posterior ethmoidal cells
- ☐ D. Maxillary sinus

46. The posterior ethmoidal cells drain into:

- ☐ A. The inferior meatus
- ☐ B. The middle meatus
- ☐ C. The spheno-ethmoidal recess
- ☐ D. The superior meatus

47. The sphenoidal sinus drains into:

- ☐ A. The spheno-ethmoidal recess
- ☐ B. The maxillary sinus
- ☐ C. The middle meatus
- ☐ D. The inferior meatus

48. The nasolacrimal duct opens into:

- ☐ A. The nasopharynx
- ☐ B. The middle meatus
- ☐ C. The inferior meatus
- ☐ D. The superior meatus

49. The frontal sinus drains into:

- ☐ A. The spheno-ethmoidal recess
- ☐ B. The middle meatus, via the frontonasal duct
- ☐ C. The inferior meatus
- ☐ D. The superior meatus

50. The frontal sinus:

- ☐ A. Drains into the spheno-ethmoidal recess
- ☐ B. Is rudimentary at birth and usually well developed by 7-8 years
- ☐ C. Is fully developed at birth
- ☐ D. Is supplied by V2

51. The ethmoidal sinus: all of the following are correct EXCEPT:

- ☐ A. Is a single large cavity
- ☐ B. Forms the ethmoidal labyrinth between the orbits and nasal fossae
- ☐ C. Has 3 to 18 cells on each side
- ☐ D. Has anterior, middle and posterior cells

52. The ethmoid bulla is produced by:

- ☐ A. The posterior ethmoidal cells
- ☐ B. The middle ethmoidal cells
- ☐ C. The inferior concha
- ☐ D. The sphenoidal sinus

53. Which nerve supplies the ethmoidal sinus?

- ☐ A. The glossopharyngeal nerve
- ☐ B. The facial nerve
- ☐ C. The chorda tympani
- ☐ D. The anterior ethmoidal nerve (V1)

54. Lymph from the posterior ethmoidal cells drains to:

- ☐ A. Preauricular nodes
- ☐ B. Submental nodes
- ☐ C. Retropharyngeal nodes
- ☐ D. Submandibular nodes

55. The maxillary sinus: all of the following are correct EXCEPT:

- ☐ A. Is the largest paranasal sinus
- ☐ B. Is pyramidal
- ☐ C. Fills the body of the maxilla
- ☐ D. Opens into the inferior meatus

56. The roof of the maxillary sinus is:

- ☐ A. The floor of the orbit
- ☐ B. The hard palate
- ☐ C. The alveolar process
- ☐ D. The lateral wall of the nose

57. The maxillary sinus is supplied by:

- ☐ A. The ophthalmic nerve (V1)
- ☐ B. The mandibular nerve (V3)
- ☐ C. The facial nerve
- ☐ D. Branches of V2 (infraorbital and superior alveolar)

58. The maxillary sinus ostium opens:

- ☐ A. At the hiatus semilunaris, in the middle meatus
- ☐ B. In the inferior meatus
- ☐ C. In the superior meatus
- ☐ D. In the spheno-ethmoidal recess

59. A patient has upper molar 'toothache' with nasal congestion and cheek pain. The best explanation:

- ☐ A. Sphenoid sinusitis
- ☐ B. Ethmoid sinusitis
- ☐ C. Frontal sinusitis
- ☐ D. Maxillary sinusitis: the sinus floor is the maxillary alveolus and shares V2 supply

60. Which lies directly above the sphenoidal sinus?

- ☐ A. The nasopharynx
- ☐ B. The cavernous sinus
- ☐ C. The frontal sinus
- ☐ D. The pituitary in the hypophyseal fossa

61. The sphenoidal sinuses: all of the following are correct EXCEPT:

- ☐ A. Are supplied by V1 and V2
- ☐ B. Are two cavities separated by a septum
- ☐ C. Always have a midline septum
- ☐ D. Lie in the body of the sphenoid

62. Trans-sphenoidal surgery reaches the pituitary because:

- ☐ A. The pituitary lies just above the sphenoidal sinus, which opens into the nose
- ☐ B. The ethmoid sinus lies between the sphenoid sinus and the pituitary
- ☐ C. The pituitary lies inside the maxillary sinus
- ☐ D. The pituitary lies in the anterior cranial fossa

63. Which statement about the paranasal sinuses is correct?

- ☐ A. The sphenoidal sinus opens in the spheno-ethmoidal recess
- ☐ B. The sphenoid septum is always in the midline
- ☐ C. The maxillary ostium is in the inferior meatus
- ☐ D. The frontal sinuses lie on the sides of the nasal cavity

64. The choanae:

- ☐ A. Open the nasal cavities into the nasopharynx
- ☐ B. Are the nostrils
- ☐ C. Open into the oral cavity
- ☐ D. Are separated by the perpendicular plate of the ethmoid

65. The inferior nasal concha:

- ☐ A. Belongs to the ethmoid
- ☐ B. Is an independent bone
- ☐ C. Is the smallest concha
- ☐ D. Has the middle meatus beneath it

66. Which nasal concha is a separate bone?

- ☐ A. Middle concha
- ☐ B. Supreme concha
- ☐ C. Inferior concha
- ☐ D. Superior concha

67. Which does NOT form part of the nasal septum?

- ☐ A. Vomer
- ☐ B. Palatine bone
- ☐ C. Perpendicular plate of the ethmoid
- ☐ D. Septal cartilage

The orbit: osteology and myology

68. The apex of the orbit lies at:

- ☐ A. The supraorbital notch
- ☐ B. The inferior orbital fissure
- ☐ C. The lacrimal fossa
- ☐ D. The optic canal

69. Orbital axes: all of the following are correct EXCEPT:

- ☐ A. The medial walls are roughly parallel
- ☐ B. The orbital axes make about 45 degrees with each other
- ☐ C. The lateral walls meet at about 90 degrees
- ☐ D. The visual axes meet at 45 degrees

70. How many bones form the orbit?

- ☐ A. 6: 3 cranial and 3 facial
- ☐ B. 7: 4 cranial and 3 facial
- ☐ C. 5, all cranial
- ☐ D. 7: 3 cranial and 4 facial

71. The roof of the orbit is formed by:

- ☐ A. The frontal bone and the lacrimal bone
- ☐ B. The orbital plate of the frontal bone and the lesser wing of the sphenoid
- ☐ C. The cribriform plate of the ethmoid
- ☐ D. The greater wing of the sphenoid and the zygomatic bone

72. Which bone does NOT form part of the floor of the orbit?

- ☐ A. Zygomatic bone
- ☐ B. Orbital surface of the maxilla
- ☐ C. Orbital process of the palatine
- ☐ D. Lacrimal bone

73. Which does NOT form part of the medial wall of the orbit?

- ☐ A. Greater wing of the sphenoid
- ☐ B. Orbital plate of the ethmoid
- ☐ C. Frontal process of the maxilla
- ☐ D. Lacrimal bone

74. The lateral wall of the orbit is formed by:

- ☐ A. The lacrimal bone and the greater wing of the sphenoid
- ☐ B. The maxilla and the zygomatic bone
- ☐ C. The zygomatic bone and the greater wing of the sphenoid
- ☐ D. The lesser wing of the sphenoid and the zygomatic bone

75. Which wall of the orbit is the thinnest?

- ☐ A. Lateral
- ☐ B. Medial (lamina papyracea)
- ☐ C. Roof
- ☐ D. Floor

76. Which wall is the strongest?

- ☐ A. Lateral
- ☐ B. Floor
- ☐ C. Medial
- ☐ D. Roof

77. The orbital floor:

- ☐ A. Is formed by the greater wing of the sphenoid
- ☐ B. Gives origin to the superior oblique
- ☐ C. Is the roof of the maxillary sinus
- ☐ D. Is the strongest wall of the orbit

78. The fossa for the lacrimal gland lies:

- ☐ A. In the medial wall
- ☐ B. In the roof, superomedially
- ☐ C. In the roof, anterolaterally
- ☐ D. In the floor

79. The supraorbital notch lies on:

- ☐ A. The superior orbital margin (frontal bone)
- ☐ B. The zygomatic bone
- ☐ C. The maxilla
- ☐ D. The inferior orbital margin

80. The inferior orbital margin is formed by:

- ☐ A. The zygomatic bone and the maxilla
- ☐ B. The frontal bone and the maxilla
- ☐ C. The lacrimal bone and the zygomatic bone
- ☐ D. The maxilla only

81. The optic canal transmits:

- ☐ A. The optic nerve and the ophthalmic veins
- ☐ B. The optic nerve and the nasociliary nerve
- ☐ C. The optic nerve and the ophthalmic artery
- ☐ D. The ophthalmic artery and the abducens nerve

82. Which passes through the superior orbital fissure INSIDE the tendinous ring?

- ☐ A. Lacrimal nerve
- ☐ B. Trochlear nerve
- ☐ C. Frontal nerve
- ☐ D. Abducens nerve

83. Which does NOT pass through the superior orbital fissure OUTSIDE the tendinous ring?

- ☐ A. Superior ophthalmic vein
- ☐ B. Lacrimal nerve
- ☐ C. Abducens nerve
- ☐ D. Frontal nerve

84. The inferior orbital fissure connects the orbit to:

- ☐ A. The anterior cranial fossa
- ☐ B. The pterygopalatine fossa
- ☐ C. The nasal cavity
- ☐ D. The middle cranial fossa

85. The nasolacrimal canal runs from:

- ☐ A. The lacrimal fossa to the inferior meatus
- ☐ B. The lacrimal fossa to the middle meatus
- ☐ C. The orbit to the maxillary sinus
- ☐ D. The lacrimal gland to the superior meatus

86. The anterior and posterior ethmoidal foramina lie:

- ☐ A. In the floor
- ☐ B. In the lateral wall
- ☐ C. At the junction of the floor and lateral wall
- ☐ D. At the junction of the roof and medial wall

87. The maxillary nerve enters the orbit through:

- ☐ A. The inferior orbital fissure
- ☐ B. The optic canal
- ☐ C. The supraorbital notch
- ☐ D. The superior orbital fissure

88. The lateral rectus is supplied by:

- ☐ A. CN V1
- ☐ B. CN III
- ☐ C. CN IV
- ☐ D. CN VI

89. Which extraocular muscle does NOT arise at the apex of the orbit?

- ☐ A. Superior oblique
- ☐ B. Lateral rectus
- ☐ C. Levator palpebrae superioris
- ☐ D. Inferior oblique

90. Levator palpebrae superioris is supplied by:

- ☐ A. CN III
- ☐ B. Sympathetic fibres only
- ☐ C. CN VII
- ☐ D. CN IV

91. The four recti:

- ☐ A. Arise from the orbital floor
- ☐ B. Are all supplied by CN III
- ☐ C. Arise from the common tendinous ring
- ☐ D. Insert behind the equator

92. In an orbital floor blowout fracture, the muscle most likely trapped is:

- ☐ A. Lateral rectus
- ☐ B. Superior oblique
- ☐ C. Superior rectus
- ☐ D. Inferior rectus

Orbital appendages: eyelids, conjunctiva, lacrimal apparatus

93. Which is NOT a content of the orbit?

- ☐ A. The eyeball and extraocular muscles
- ☐ B. The ciliary ganglion and ophthalmic vessels
- ☐ C. The lacrimal gland and orbital fat
- ☐ D. The pterygopalatine ganglion

94. The superior tarsal (Müller's) muscle is:

- ☐ A. Smooth muscle supplied by sympathetic fibres
- ☐ B. Striated muscle supplied by CN III
- ☐ C. Striated muscle supplied by CN VII
- ☐ D. Smooth muscle supplied by CN III

95. The eyelid: all of the following are correct EXCEPT:

- ☐ A. Has a lateral ciliary part carrying the eyelashes
- ☐ B. Has no muscle layer
- ☐ C. Is 2 to 3 mm thick
- ☐ D. Has five layers

96. The layers of the eyelid from superficial to deep:

- ☐ A. Skin, muscle, conjunctiva, tarsal plate
- ☐ B. Conjunctiva, muscle, tarsal plate, skin
- ☐ C. Skin and cilia, loose connective tissue, muscle, tarsal plate, palpebral conjunctiva
- ☐ D. Skin, tarsal plate, muscle, conjunctiva

97. The glands of Zeis are:

- ☐ A. Mucin-secreting goblet cells
- ☐ B. The source of the aqueous layer of tears
- ☐ C. Sebaceous glands near the lashes
- ☐ D. Modified apocrine sweat glands

98. The glands of Moll are:

- ☐ A. Modified apocrine sweat glands
- ☐ B. Mucin-producing glands
- ☐ C. Sebaceous glands
- ☐ D. Embedded in the tarsal plates

99. The eyelashes: all of the following are correct EXCEPT:

- ☐ A. Act as sensory receptors
- ☐ B. Help prevent evaporation of the tear film
- ☐ C. Produce tears
- ☐ D. Protect the eye from foreign particles

100. The tarsal plates:

- ☐ A. Contain the glands of Moll
- ☐ B. Are present in the upper lid only
- ☐ C. Contain the meibomian glands
- ☐ D. Are hyaline cartilage

101. The orbital septum:

- ☐ A. Connects the tarsi to the orbital margins
- ☐ B. Lies behind the orbital fat
- ☐ C. Is a muscle layer of the lid
- ☐ D. Is part of the conjunctiva

102. The conjunctiva: all of the following are correct EXCEPT:

- ☐ A. Covers the cornea completely
- ☐ B. Protects against microbial infection
- ☐ C. Palpebral conjunctiva reflects at the fornices onto the eyeball as bulbar conjunctiva
- ☐ D. Lubricates the eye and lids

103. The mucin layer of the tear film comes from:

- ☐ A. Conjunctival goblet cells
- ☐ B. The meibomian glands
- ☐ C. The lacrimal gland
- ☐ D. The glands of Moll

104. The oily layer of the tear film:

- ☐ A. Comes from goblet cells
- ☐ B. Comes from the lacrimal gland
- ☐ C. Prevents evaporation
- ☐ D. Attaches the tear film to the eye

105. The lacrimal gland lies:

- ☐ A. In the lacrimal fossa of the lacrimal bone
- ☐ B. In the medial wall of the orbit
- ☐ C. On the orbital floor
- ☐ D. In the superolateral orbit, in the lacrimal fossa of the frontal bone

106. Relations of the lacrimal gland: all of the following are correct EXCEPT:

- ☐ A. Superior: zygomatic process of the frontal bone
- ☐ B. Anterior: orbital septum
- ☐ C. Inferolateral: medial rectus
- ☐ D. Posterior: orbital fat

107. The ducts of the lacrimal gland open into:

- ☐ A. The lacrimal sac
- ☐ B. The inferior meatus
- ☐ C. The puncta
- ☐ D. The superior conjunctival fornix

108. The lacrimal gland is supplied by:

- ☐ A. The superficial temporal artery
- ☐ B. The lacrimal artery, from the ophthalmic artery
- ☐ C. The facial artery
- ☐ D. The maxillary artery directly

109. Lymph from the lacrimal gland drains first to:

- ☐ A. Retropharyngeal nodes
- ☐ B. Superficial parotid nodes
- ☐ C. Submental nodes
- ☐ D. Axillary nodes

110. Order of tear flow:

- ☐ A. Lacrimal gland → lacrimal sac → puncta → middle meatus
- ☐ B. Lacrimal gland → fornix → conjunctival sac → puncta → lacrimal sac → nasolacrimal duct → inferior meatus
- ☐ C. Puncta → lacrimal gland → inferior meatus
- ☐ D. Lacrimal gland → nasolacrimal duct → puncta

111. Tears are drawn into the puncta by:

- ☐ A. Active secretion by the conjunctiva
- ☐ B. Capillary action
- ☐ C. Active pumping by the lacrimal gland
- ☐ D. Ciliary beating

112. The sensory nerve of the lacrimal gland is:

- ☐ A. The greater petrosal nerve
- ☐ B. The zygomatic nerve
- ☐ C. The lacrimal nerve, a branch of V1
- ☐ D. The chorda tympani

113. Secretomotor fibres to the lacrimal gland synapse in:

- ☐ A. The submandibular ganglion
- ☐ B. The otic ganglion
- ☐ C. The pterygopalatine ganglion
- ☐ D. The ciliary ganglion

114. Sympathetic fibres to the lacrimal gland:

- ☐ A. Stimulate secretion
- ☐ B. Come from the superior cervical ganglion via the deep petrosal nerve
- ☐ C. Synapse in the pterygopalatine ganglion
- ☐ D. Travel in the greater petrosal nerve

115. The ciliary ganglion: all of the following are correct EXCEPT:

- ☐ A. Gives the short ciliary nerves
- ☐ B. Is a parasympathetic ganglion relaying fibres of CN VII
- ☐ C. Also transmits sympathetic fibres from the internal carotid plexus
- ☐ D. Also transmits sensory fibres from the nasociliary nerve

116. A facial nerve lesion at the geniculate ganglion causes, on the same side:

- ☐ A. Loss of corneal sensation
- ☐ B. Loss of taste only
- ☐ C. Facial paralysis and a dry eye
- ☐ D. Facial paralysis with normal tear secretion

117. A patient has mild ptosis and a small pupil on one side. The most likely lesion:

- ☐ A. Oculomotor nerve
- ☐ B. Trochlear nerve
- ☐ C. Facial nerve
- ☐ D. Sympathetic supply (Horner syndrome)

Answer key

The single correct answer for each question, with a short explanation where it helps.

Oral cavity, tongue, palate and salivary glands

1	B	
2	A	There is a gap behind the last molars even with the teeth together.
3	C	Covered by skin outside and mucosa inside.
4	A	Stensen's duct pierces the buccinator.
5	B	
6	D	
7	C	The buccal nerve (V3) is sensory; the buccal branch of VII is motor.
8	C	
9	C	
10	A	Vallate papillae sit just in front of the sulcus; the foramen caecum is at its apex.
11	D	
12	D	The posterior third has no papillae but the lingual tonsil.
13	B	
14	B	Moving the tongue is the role of the extrinsic muscles.
15	C	Mylohyoid forms the floor of the mouth; it is not a tongue muscle.
16	B	
17	A	Styloglossus (and hyoglossus) retract; genioglossus protrudes.
18	B	Styloglossus and palatoglossus elevate; hyoglossus depresses.
19	C	Taste of the anterior two thirds = chorda tympani.
20	B	Chorda tympani fibres travel with the lingual nerve.
21	C	
22	A	
23	C	Palatoglossus is supplied by the pharyngeal plexus (vagus).
24	C	The healthy right genioglossus pushes the tongue toward the paralysed left side.
25	B	
26	C	
27	D	
28	A	
29	A	It is the only palatine muscle on V3.
30	C	Tensor veli palatini is the odd one out: V3.
31	C	
32	A	
33	B	
34	A	
35	A	

36	D	CN IX, lesser petrosal, otic ganglion, auriculotemporal (V3).
37	B	It crosses (does not pierce) the masseter, then pierces buccinator.
38	C	The 2026 mock exam had 'purely mucous' as a false option.
39	A	
40	D	
41	B	The sublingual gland is mixed, mainly mucous.

Nose and paranasal sinuses

42	A	Rudimentary at birth; enlarge at 6-7 years and after puberty.
43	B	The nasopharynx lies behind the choanae; the fourth space is the sphenoid recess.
44	D	
45	C	Also on the 2026 mock. Posterior ethmoidal → superior meatus.
46	D	
47	A	The recess lies above and behind the superior concha.
48	C	
49	B	Supplied by supraorbital and supratrochlear nerves (V1).
50	B	
51	A	
52	B	
53	D	
54	C	Anterior and middle cells drain to submandibular nodes.
55	D	Its ostium is in the middle meatus.
56	A	Apex: the zygomatic process of the maxilla.
57	D	
58	A	
59	D	
60	D	Cavernous sinus is lateral; nasopharynx is below.
61	C	The septum is often off midline (also a 2026 mock option).
62	A	
63	A	From the 2026 mock exam.
64	A	The choanae are separated by the vomer.
65	B	It is the largest/most extended. Superior and middle conchae belong to the ethmoid.
66	C	Superior, middle (and supreme) conchae belong to the ethmoid.
67	B	From the 2026 mock exam.

The orbit: osteology and myology

68	D	
69	D	In the primary position the visual axes are parallel; it is the orbital axes that diverge.
70	D	Cranial: frontal, sphenoid, ethmoid. Facial: zygomatic, lacrimal, maxilla, palatine.
71	B	Past exam Q1 tested the roof and the 'cribriform plate' distractor.
72	D	

73	A	The greater wing is in the lateral wall; the medial wall has the body of the sphenoid.
74	C	It is the strongest wall.
75	B	
76	A	
77	C	
78	C	The trochlear fossa is the superomedial one.
79	A	
80	A	
81	C	The superior ophthalmic vein uses the superior orbital fissure.
82	D	CN IV passes outside the ring.
83	C	The abducens passes inside the ring.
84	B	The superior orbital fissure joins the orbit to the middle cranial fossa.
85	A	
86	D	
87	A	It exits at the infraorbital foramen.
88	D	LR6 SO4.
89	D	It arises from the front of the orbital floor (maxilla).
90	A	Its smooth-muscle part (Müller's) is sympathetic.
91	C	
92	D	

Orbital appendages: eyelids, conjunctiva, lacrimal apparatus

93	D	That ganglion lies in the pterygopalatine fossa, not the orbit.
94	A	
95	B	
96	C	
97	C	Glands of Moll are modified apocrine sweat glands.
98	A	
99	C	
100	C	
101	A	
102	A	
103	A	
104	C	The mucin layer attaches the film to the eye.
105	D	The lacrimal bone holds the lacrimal sac, not the gland.
106	C	Inferolaterally the gland rests on the lateral rectus.
107	D	
108	B	
109	B	

110	B	
111	B	Gravity and blinking carry tears across the cornea; capillary action draws them into the puncta.
112	C	
113	C	Postganglionic fibres then travel with V2 → zygomatic → lacrimal nerve.
114	B	
115	B	The ciliary ganglion relays CN III fibres.
116	C	The greater petrosal carries lacrimal secretomotor fibres.
117	D	Loss of the sympathetic superior tarsal muscle and dilator pupillae.