Generalized Fiducial Inference for Hybrid Discrete-Continuous Count Data

Kendall Thomas

Doctoral Candidate, Statistics and Operations Research

Joint work with Dr. Jan Hannig



The University
of North Carolina
at Chapel Hill

UNC Research Week 2026

Motivation

- **Quantile cube**: time spent in 100 velocity × acceleration × angle bins per athlete–match session

- Counts range from under 100 to several thousand deciseconds:
  - NB is exact but costly
  - a Normal approximation fails where counts are small relative to their dispersion

- **Goal:** uncertainty for 500 regression and dispersion parameters, without a prior





Fig. from [1] Thomas & Hannig (2026), Journal of Quantitative Analysis in Sports

The Data

- GPS tracking data from one season for a professional women's soccer team (IRB 25-1234)
  - Sampling at 100 Hz with access at 10 Hz: timestamp, longitude, latitude
- Retained athlete-match sessions with ≥ 60 minutes played in ≥ 3 matches
  - Total: 17 athletes, 26 matches → 216 athlete-match sessions
- Covariates:
  - standardized age (mean 27.3 yrs, SD 2.1; range 22.2 – 33.1)
  - position (6 defenders, 4 midfielders, 7 forwards)



The Model

- **Negative Binomial (NB) shape** $r_{ij} = \mu_{ij}\nu_j$ (mean × dispersion) decides how close count i in bin j is to Normal, not the count itself

- **Our Approach**: assign each count by a 95% lower bound $\hat{r}_{ij}^{L}$ on its shape, cutoff $\omega = 10$

$$y_{ij} \sim \begin{cases} \mathrm{NB}\left(\alpha_{ij}t_i, \dfrac{\nu_j}{1+\nu_j}\right) & \text{if } \hat{r}_{ij}^{L} < \omega, \\ \mathcal{N}\left(\mu_{ij} = \dfrac{\alpha_{ij}t_i}{\nu_j}, \sigma_{ij}^2 = \dfrac{\alpha_{ij}t_i(\nu_j+1)}{\nu_j^2}\right) & \text{if } \hat{r}_{ij}^{L} \geq \omega \end{cases}$$

where $\alpha_{ij} = exp({X_i}^{\mathsf{T}}\beta_j)$ and $\boldsymbol{\theta} = ({\beta_1}^{\mathsf{T}}, \ldots, {\beta_{100}}^{\mathsf{T}}, \nu_1, \ldots, \nu_{100})^{\mathsf{T}}$

The Generalized Fiducial Density

- Generalized Fiducial Inference (GFI) expresses the data through a structural equation $Y = G(U,\ \theta)$ involving the parameters $\boldsymbol{\theta}$ and auxiliary variables $\mathbf{U}$ with known distribution[2, 3]

- **Theorem 2**: The Joint Generalized Fiducial Density takes the form

$$r(\boldsymbol{\theta} \mid \mathbf{y}) \propto L(\mathbf{y} \mid \boldsymbol{\theta})\, J(\boldsymbol{\theta}, \mathbf{y_C})$$

- **Proposition 3 (Block-Diagonal Jacobian):** The Jacobian decomposes into one block per movement bin, where each block depends only on that bin's own parameters $\theta_j = \left(\beta_j^{\mathrm{T}}, \nu_j\right)$.

$$J_C(\boldsymbol{\theta}) = \partial \mathrm{Y_C} / \partial \boldsymbol{\theta} = \mathrm{diag}(J_1, J_2, \ldots, J_{100})$$


$$J(\boldsymbol{\theta}, \mathbf{y_C}) = \sqrt{\det(\mathrm{n}^{-1} J_C(\boldsymbol{\theta})^{\mathrm{T}} J_C(\boldsymbol{\theta}))} = \prod_{j=1}^{100} \sqrt{\det\left(n^{-1} J_j^{\mathrm{T}} J_j\right)}$$

Fitting the Model


- **Lower-bound shape rule**: count i of bin j is Normal when

$$\hat{r}_{ij}^{L} = \hat{\mu}_{ij}\hat{v}_j e^{-1.645\hat{s}_j} \geq \omega = 10$$

  - $\hat{\mu}_{ij}$: pooled rate × exposure; $\hat{v}_j$: moment estimate; $\hat{s}_j$: athlete-bootstrap SD
- **Algorithm**: block-diagonal Jacobian, so each bin is fit independently
  - 74 active bins: Hamiltonian Monte Carlo, 4 × 4,000 draws, all $\hat{R} < 1.01$
  - 26 all-NB bins: exact discrete construction
- **Multiple testing**: Benjamini–Hochberg[4] FDR 0.05 per covariate




Simulation Results




Coverage at every nominal level, borderline regime

- NB regression (intercept, continuous covariate, 3-level group, exposure); n = 50 to 500; 2,000 replications
- Average coverage of 95% intervals: **94.6%** (93–96%); calibrated at every level (left, borderline regime)



- Partition by shape (right): shape rules ≈ 0.95; mean rule ≈ 0.92; realized counts fall to 0.71

Results: Age and Position



- Declared (BH, FDR 0.05, working model): age **18** bins; forward **23** and midfielder **48** (vs. defender)
- Position effects reach $|\beta| = 0.29$ (midfielder, highest velocity); age effects stay below 0.09 per SD (2.1 yrs) and follow an acceleration gradient

Calibration Check

- **17 athletes, not 216 sessions:** reassign athletes' labels at random (every session of an athlete keeps its permuted label), refit all 100 bins, repeat 100 times

Covariate	Bins declared	Median under permutation
Age	18	28.5
Forward	23	31
Midfielder	48	27.5

- Significance under the working model is not calibrated at the athlete level: the independent units are the 17 athletes
- The declared bins show where the fitted model places its largest, best-resolved contrasts: descriptive, not established

Conclusion

Exact where it matters, approximate where it is safe.

Hybrid GFI

exact NB where the shape is small, Normal where it is large

block-diagonal Jacobian

Bernstein–von Mises

Simulations

near-nominal coverage

partition by estimated shape, not by the mean

Application

position effects exceed age effects

descriptive at the athlete level (17 athletes)

Acknowledgements

- I extend my sincere thanks to the professional women's soccer team, coaching staff, and support personnel who facilitate data collection and generously share their time and expertise, enabling the extension of my research.



The University of North Carolina at Chapel Hill

Thank you!
Questions?

Email: ketho@unc.edu

Website: https://kelanethomas.github.io/home/

Full Paper

quantile-cube Python Package Available on PyPI!

References

1. Thomas, K. L. and Hannig, J. (2026), "Movement Dynamics in Elite Female Soccer Athletes: The Quantile Cube Approach", *Journal of Quantitative Analysis in Sports*.
2. Hannig, J. (2009), "On Generalized Fiducial Inference", *Statistica Sinica,* 19, 491-544.
3. Hannig, J., Iyer, H., Lai, R. C. S. and Lee, T. C. M. (2016), "Generalized fiducial inference: A review and new results," *Journal of the American Statistical Association,* 111(515), 1346–1361.
4. Benjamini, Y. and Hochberg, Y. (1995), "Controlling the false discovery rate: A practical and powerful approach to multiple testing," *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(1), 289–300.