

Amniote Mass

Erin Keleske

3/22/2018

Species experience tradeoffs for where there energy is spent. In this brief analysis, we will consider one such relationship: energy spent on personal growth and energy spent on the growth of offspring.

Initial data download —

Raw data can be found here. and originated from:

Nathan P. Myhrvold, Elita Baldrige, Benjamin Chan, Dhileep Sivam, Daniel L. Freeman, and S. K. Morgan Ernest. 2015. An amniote life-history database to perform comparative analyses with birds, mammals, and reptiles. *Ecology* 96:3109.<http://dx.doi.org/10.1890/15-0846.1>


This data includes common life history data on species from the class Aves, Mammalia, and Reptilia.

Data exploration and visualisation —


Assuming females of a given species have reached “adult body mass” by the time of sexual maturity, the relationship between these two variables can be used to calculate the approximate growth rate from birth to adulthood.

First, let’s explore the relationship between these variables. Since there’s such a wide range of sizes within the full amniote data, from the Dwarf Gecko to the Blue Whale, the data have been subset into classes and plotted separately.


```
## `geom_smooth()` using method = 'gam'
```



```
## `geom_smooth()` using method = 'gam'
```




```
## `geom_smooth()` using method = 'gam'
```




To represent investment in reproduction, we can look at a similar relationship between the weight at birth or hatching divided by gestation period.

Again, data was subset into classes. However, there is no available data on Aves gestation period.

```
## `geom_smooth()` using method = 'gam'
```



```
## `geom_smooth()` using method = 'gam'
```



Relationship between variables —

Distribution of a variable among groups —