

Multivariate anisotropic wavelets on the torus

Ronny Bergmann

AG Bildverarbeitung
TU Kaiserslautern

October 2nd, 2013

Workshop on
Advances in Image Processing
Anweiler am Trifels



FELIX KLEIN
ZENTRUM FÜR
MATHEMATIK

Introduction

Periodic wavelets were first defined for the univariate case [PT95]

- based on shifts by $2\pi/N$, $N \in \mathbb{N}$
- fast wavelet transform [Se98]
- (localised) de la Vallée Poussin mean based wavelets [Se98]


For the multivariate generalization

- a function may be shifted by $\mathbf{y} \in \mathbb{T}^d := [-\pi, \pi)^d$
- scaling factor j replaced by a matrix \mathbf{J} [MS03]
- for fixed $|\det \mathbf{J}| = 2$: several matrices \mathbf{J} available [LP10]

⇒ preference of direction

- handle “curse of dimension”

The pattern and the generating set



The pattern $\mathcal{P}(\mathbf{M})$,

$$\mathbf{M} = \begin{pmatrix} 28 & -12 \\ 12 & 4 \end{pmatrix}$$


Let $\mathbf{M} \in \mathbb{Z}^{d \times d}$ be regular.

- pattern
 $\mathcal{P}(\mathbf{M}) := [-\frac{1}{2}, \frac{1}{2})^d \cap \mathbf{M}^{-1}\mathbb{Z}^d$
- generating set
 $\mathcal{G}(\mathbf{M}) := \mathbf{M}\mathcal{P}(\mathbf{M}) = \mathbf{M}[-\frac{1}{2}, \frac{1}{2})^d \cap \mathbb{Z}^d$

We have

- $m := |\mathcal{P}(\mathbf{M})| = |\mathcal{G}(\mathbf{M})| = |\det \mathbf{M}|$
- the group $(\mathcal{P}(\mathbf{M}), + \text{ mod } 1)$
- subpatterns $\mathcal{P}(\mathbf{N})$, für
 $\mathbf{M} = \mathbf{JN}$, $\mathbf{J}, \mathbf{N} \in \mathbb{Z}^{d \times d}$

The pattern and the generating set



subpattern $\mathcal{P}(\mathbf{N}) \subset \mathcal{P}(\mathbf{M})$,

$$\mathbf{M} = \begin{pmatrix} 28 & -12 \\ 12 & 4 \end{pmatrix} = \mathbf{J}_Y \begin{pmatrix} 28 & -12 \\ 6 & 2 \end{pmatrix}$$

$$\mathbf{J}_Y = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$$

Let $\mathbf{M} \in \mathbb{Z}^{d \times d}$ be regular.

- pattern
 $\mathcal{P}(\mathbf{M}) := \left[-\frac{1}{2}, \frac{1}{2}\right)^d \cap \mathbf{M}^{-1}\mathbb{Z}^d$
- generating set
 $\mathcal{G}(\mathbf{M}) := \mathbf{M}\mathcal{P}(\mathbf{M}) = \mathbf{M}\left[-\frac{1}{2}, \frac{1}{2}\right)^d \cap \mathbb{Z}^d$

We have

- $m := |\mathcal{P}(\mathbf{M})| = |\mathcal{G}(\mathbf{M})| = |\det \mathbf{M}|$
- the group $(\mathcal{P}(\mathbf{M}), + \text{ mod } 1)$
- subpatterns $\mathcal{P}(\mathbf{N})$, für
 $\mathbf{M} = \mathbf{JN}$, $\mathbf{J}, \mathbf{N} \in \mathbb{Z}^{d \times d}$

The Fourier transform

Let $\mathbf{M} \in \mathbb{Z}^{d \times d}$ be regular.

For fixed orderings of $\mathcal{G}(\mathbf{M}^T)$ and $\mathcal{P}(\mathbf{M})$:

- Fourier matrix

$$\mathcal{F}(\mathbf{M}) := \frac{1}{\sqrt{m}} \left(e^{-2\pi i \mathbf{h}^T \mathbf{y}} \right)_{\mathbf{h} \in \mathcal{G}(\mathbf{M}^T), \mathbf{y} \in \mathcal{P}(\mathbf{M})} \in \mathbb{C}^{m \times m}$$

- discrete Fourier transform for $\mathbf{a} = (a_{\mathbf{y}})_{\mathbf{y} \in \mathcal{P}(\mathbf{M})} \in \mathbb{C}^m$

$$\hat{\mathbf{a}} = (\hat{a}_{\mathbf{h}})_{\mathbf{h} \in \mathcal{G}(\mathbf{M}^T)} := \sqrt{m} \mathcal{F}(\mathbf{M}) \mathbf{a} \in \mathbb{C}^m$$

\Rightarrow **fast Fourier transform (B., 2013)**

We further need for $f \in L_2(\mathbb{T}^d)$ its Fourier coefficients

$$c_{\mathbf{k}}(f) := \langle f, e^{i\mathbf{k}^T \cdot} \rangle = \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} f(\mathbf{x}) e^{-i\mathbf{k}^T \mathbf{x}} d\mathbf{x}, \quad \mathbf{k} \in \mathbb{Z}^d.$$

Translation invariant space

The space $V_{\mathbf{M}}^{\xi} := \text{span}\{T_{\mathbf{y}}\xi, \mathbf{y} \in \mathcal{P}(\mathbf{M})\} \subset L_2(\mathbb{T}^d)$ is \mathbf{M} -invariant, i.e. for all $\mathbf{y} \in \mathcal{P}(\mathbf{M})$, $\varphi \in V_{\mathbf{M}}^{\xi}$ we have

$$T_{\mathbf{y}}\varphi := \varphi(\circ - 2\pi\mathbf{y}) \in V_{\mathbf{M}}^{\xi}.$$

Further

$$\varphi = \sum_{\mathbf{y} \in \mathcal{P}(\mathbf{M})} a_{\varphi, \mathbf{y}} T_{\mathbf{y}}\xi, \quad a_{\mathbf{y}} \in \mathbb{C},$$

Translation invariant space

The space $V_{\mathbf{M}}^{\xi} := \text{span}\{T_{\mathbf{y}}\xi, \mathbf{y} \in \mathcal{P}(\mathbf{M})\} \subset L_2(\mathbb{T}^d)$ is \mathbf{M} -invariant, i.e. for all $\mathbf{y} \in \mathcal{P}(\mathbf{M})$, $\varphi \in V_{\mathbf{M}}^{\xi}$ we have

$$T_{\mathbf{y}}\varphi := \varphi(\circ - 2\pi\mathbf{y}) \in V_{\mathbf{M}}^{\xi}.$$

Further

$$\varphi = \sum_{\mathbf{y} \in \mathcal{P}(\mathbf{M})} a_{\varphi, \mathbf{y}} T_{\mathbf{y}}\xi, \quad a_{\mathbf{y}} \in \mathbb{C},$$

in Fourier coefficients: For all $\mathbf{h} \in \mathcal{G}(\mathbf{M}^T)$, $\mathbf{z} \in \mathbb{Z}^d$

$$c_{\mathbf{h} + \mathbf{M}^T \mathbf{z}}(\varphi) = \sum_{\mathbf{y} \in \mathcal{P}(\mathbf{M})} a_{\varphi, \mathbf{y}} e^{-2\pi i \mathbf{h}^T \mathbf{y}} c_{\mathbf{h} + \mathbf{M}^T \mathbf{z}}(\xi) = \hat{a}_{\varphi, \mathbf{h}} c_{\mathbf{h} + \mathbf{M}^T \mathbf{z}}(\xi),$$

where $\hat{\mathbf{a}} = \sqrt{m} \mathcal{F}(\mathbf{M}) \mathbf{a}$.

Especially: $V_{\mathbf{N}}^{\varphi} \subseteq V_{\mathbf{M}}^{\xi}$, $\mathbf{M} = \mathbf{JN}$

The wavelet transform

Given $\mathbf{M} = \mathbf{JN}$, $|\det \mathbf{J}| = 2$, and $\xi, \varphi \in L_2(\mathbb{T}^d)$ such that

- $T_{\mathbf{y}}\xi$, $\mathbf{y} \in \mathcal{P}(\mathbf{M})$, linear independent
- $T_{\mathbf{x}}\varphi$, $\mathbf{x} \in \mathcal{P}(\mathbf{N})$, linear independent, i.e. $\dim V_{\mathbf{N}}^{\varphi} = \frac{1}{2} \dim V_{\mathbf{M}}^{\xi} = \frac{|\det \mathbf{M}|}{2}$

$\Rightarrow \exists$ Wavelet $\psi \in L_2(\mathbb{T}^d)$ s.t. $V_{\mathbf{M}}^{\xi} = V_{\mathbf{N}}^{\varphi} \oplus V_{\mathbf{N}}^{\psi}$

The wavelet transform

Given $\mathbf{M} = \mathbf{JN}$, $|\det \mathbf{J}| = 2$, and $\xi, \varphi \in L_2(\mathbb{T}^d)$ such that

- $T_y \xi$, $\mathbf{y} \in \mathcal{P}(\mathbf{M})$, linear independent
- $T_x \varphi$, $\mathbf{x} \in \mathcal{P}(\mathbf{N})$, linear independent, i.e. $\dim V_{\mathbf{N}}^{\varphi} = \frac{1}{2} \dim V_{\mathbf{M}}^{\xi} = \frac{|\det \mathbf{M}|}{2}$

$\Rightarrow \exists$ Wavelet $\psi \in L_2(\mathbb{T}^d)$ s.t. $V_{\mathbf{M}}^{\xi} = V_{\mathbf{N}}^{\varphi} \oplus V_{\mathbf{N}}^{\psi}$

$$\text{Decompose } f = \sum_{\mathbf{y} \in \mathcal{P}(\mathbf{M})} a_{f,\mathbf{y}} T_{\mathbf{y}} \xi = g + h = \sum_{\mathbf{x} \in \mathcal{P}(\mathbf{N})} a_{g,\mathbf{x}} T_{\mathbf{x}} \varphi + \sum_{\mathbf{x} \in \mathcal{P}(\mathbf{N})} a_{h,\mathbf{x}} T_{\mathbf{x}} \psi$$

by computing

$$\hat{a}_{g,\mathbf{k}} = \frac{1}{\sqrt{|\det \mathbf{J}|}} \sum_{\mathbf{l} \in \mathcal{G}(\mathbf{J}^T)} \overline{\hat{a}_{\varphi,\mathbf{k}+\mathbf{N}^T \mathbf{l}}} \hat{a}_{f,\mathbf{k}+\mathbf{N}^T \mathbf{l}}, \quad \mathbf{k} \in \mathcal{G}(\mathbf{N}^T),$$

Analogously for $\hat{a}_{h,\mathbf{k}}$ using the coefficients $a_{\psi,\mathbf{y}}$ of $\psi \in V_{\mathbf{M}}^{\xi}$

\Rightarrow **Fast wavelet transform in $\mathcal{O}(m)$ (B., 2013)**

Constructing wavelets

Let's take

- a nonnegative function $g : \mathbb{R}^d \rightarrow \mathbb{R}_+$ with
 - $\sum_{\mathbf{z} \in \mathbb{Z}^d} g(\mathbf{x} + \mathbf{z}) = 1$
 - $g(\mathbf{x}) > 0, \mathbf{x} \in [-\frac{1}{2}, \frac{1}{2})^d$
- matrices $\mathbf{J}_1, \dots, \mathbf{J}_n, \mathbf{M}_0 \in \mathbb{Z}^{d \times d}, \mathcal{J}_{l,k} = (\mathbf{J}_l, \dots, \mathbf{J}_k), |\det \mathbf{J}_l| = 2$
 $\Rightarrow \mathbf{M}_l := \mathbf{J}_l \cdot \dots \cdot \mathbf{J}_1 \mathbf{M}_0, m_l := 2^l |\det \mathbf{M}_0|$

and helping functions

- $B_{\mathcal{J}_{l,n}}(\mathbf{x}) := \begin{cases} \left(\sum_{\mathbf{z} \in \mathbb{Z}^d} g(\mathbf{x} + \mathbf{J}_l^T \mathbf{z}) \right) B_{\mathcal{J}_{l+1,n}}(\mathbf{J}_l^{-T} \mathbf{x}) & l \leq n \\ g(\mathbf{x}) & \text{else.} \end{cases}$
- $\tilde{B}_{\mathcal{J}_{l,n}}(\mathbf{x}) := e^{-2\pi i \mathbf{x}^T \mathbf{w}_l} \left(\sum_{\mathbf{z} \in \mathbb{Z}^d} g(\mathbf{x} + \mathbf{J}_l^T \mathbf{z} - \mathbf{v}_l) \right) B_{\mathcal{J}_{l+1,n}}(\mathbf{J}_l^{-T} \mathbf{x}), \quad l \leq n,$
 where $\mathbf{v}_l \in \mathcal{P}(\mathbf{J}_l^T) \setminus \{\mathbf{0}\}$ and $\mathbf{w}_l \in \mathcal{P}(\mathbf{J}_l) \setminus \{\mathbf{0}\}$ (both unique.)

Constructing wavelets II

Definition

Define scaling functions $\varphi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}}$ and wavelets $\psi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}}$ in Fourier coefficients

$$\mathbf{c}_{\mathbf{k}}(\varphi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}}) := \frac{1}{\sqrt{m_l}} B_{\mathcal{J}_{l+1,n}}(\mathbf{M}_l^{-\mathbf{T}} \mathbf{k}), \quad \mathbf{k} \in \mathbb{Z}^d, \quad l = 0, \dots, n$$

$$\mathbf{c}_{\mathbf{k}}(\psi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}}) = \frac{1}{\sqrt{m_l}} \tilde{B}_{\mathcal{J}_{l+1,n}}(\mathbf{M}_l^{-\mathbf{T}} \mathbf{k}), \quad \mathbf{k} \in \mathbb{Z}^{d \times d}, \quad l = 0, \dots, n-1.$$

If g arbitrary smooth $\Rightarrow B_{\mathcal{J}_{l,n}}$ smooth \Rightarrow localization

Theorem (B., 2013)

For $l = 0, \dots, n-1$

$$a) \varphi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}} \in \text{span} \left\{ T_{\mathbf{y}} \varphi_{\mathbf{M}_{l+1}}^{\mathcal{J}_{l+2,n}} ; \mathbf{y} \in \mathcal{P}(\mathbf{M}_{l+1}) \right\} =: V_{l+1}$$



$$b) V_{l+1} = V_l \oplus \text{span} \left\{ T_{\mathbf{y}} \psi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}} ; \mathbf{y} \in \mathcal{P}(\mathbf{M}_l) \right\}.$$

With slight restriction on $g \Rightarrow \psi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}} = \psi_{\mathbf{M}_l}^{(\mathbf{J}_{l+1})}$ and $\varphi_{\mathbf{M}_l}^{\mathcal{J}_{l+1,n}} = \varphi_{\mathbf{M}_l}^{(\mathbf{J}_{l+1})}$.

Example of a wavelet

De la Vallée Poussin-type scaling functions and wavelets



- based on $g = B_{\Xi}$, $\Xi = \begin{pmatrix} 1 & 0 & \frac{1}{10} & 0 \\ 0 & 1 & 0 & \frac{1}{10} \end{pmatrix}$
- $\mathbf{M} = \begin{pmatrix} 28 & -12 \\ 12 & 4 \end{pmatrix} = \mathbf{J}_X \begin{pmatrix} 14 & -6 \\ 12 & 4 \end{pmatrix}$
- $c_{\mathbf{k}}(\varphi_{\mathbf{M}}) = g(\mathbf{M}^{-T} \mathbf{k})$



Example of a wavelet

De la Vallée Poussin-type scaling functions and wavelets


- based on $g = B_{\Xi}$, $\Xi = \begin{pmatrix} 1 & 0 & \frac{1}{10} & 0 \\ 0 & 1 & 0 & \frac{1}{10} \end{pmatrix}$
- $\mathbf{M} = \begin{pmatrix} 28 & -12 \\ 12 & 4 \end{pmatrix} = \mathbf{J}_X \begin{pmatrix} 14 & -6 \\ 12 & 4 \end{pmatrix}$
- $c_{\mathbf{k}}(\varphi_{\mathbf{M}}) = g(\mathbf{M}^{-T}\mathbf{k})$, $c_{\mathbf{k}}(\varphi_{\mathbf{N}}^{(\mathbf{J}_X)}) = c_{\mathbf{k}}(\varphi_{\mathbf{N}}) = g(\mathbf{N}^{-T}\mathbf{k})$



Example of a wavelet

De la Vallée Poussin-type scaling functions and wavelets



- based on $g = B_{\Xi}$, $\Xi = \begin{pmatrix} 1 & 0 & \frac{1}{10} & 0 \\ 0 & 1 & 0 & \frac{1}{10} \end{pmatrix}$
- $\mathbf{M} = \begin{pmatrix} 28 & -12 \\ 12 & 4 \end{pmatrix} = \mathbf{J}_X \begin{pmatrix} 14 & -6 \\ 12 & 4 \end{pmatrix}$
- $c_{\mathbf{k}}(\varphi_{\mathbf{M}}) = g(\mathbf{M}^{-T}\mathbf{k})$, $c_{\mathbf{k}}(\varphi_{\mathbf{N}}^{(\mathbf{J}_X)}) = c_{\mathbf{k}}(\varphi_{\mathbf{N}}) = g(\mathbf{N}^{-T}\mathbf{k})$



Example of a wavelet


De la Vallée Poussin-type scaling functions and wavelets

- based on $g = B_{\Xi}$, $\Xi = \begin{pmatrix} 1 & 0 & \frac{1}{10} & 0 \\ 0 & 1 & 0 & \frac{1}{10} \end{pmatrix}$
- $\mathbf{M} = \begin{pmatrix} 28 & -12 \\ 12 & 4 \end{pmatrix} = \mathbf{J}_X \begin{pmatrix} 14 & -6 \\ 12 & 4 \end{pmatrix}$
- $c_{\mathbf{k}}(\varphi_{\mathbf{M}}) = g(\mathbf{M}^{-T}\mathbf{k})$, $c_{\mathbf{k}}(\varphi_{\mathbf{N}}^{(\mathbf{J}_X)}) = c_{\mathbf{k}}(\varphi_{\mathbf{N}}) = g(\mathbf{N}^{-T}\mathbf{k})$





Example of a decomposition

- radial function based on a piecewise quadratic function
- jump in second directional derivative on a circle
- sampling with $\mathbf{M} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \Rightarrow 512 \times 512$ pixel image.




Example of a decomposition




Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_Y \mathbf{N}_2$$




Wavelet $\psi_{\mathbf{N}_2}^{(\mathbf{J}_Y)}$, $\mathbf{N}_2 = \begin{pmatrix} 512 & 0 \\ 0 & 128 \end{pmatrix}$




Corresp. wavelet part

Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_Y \mathbf{J}_Y \mathbf{N}_3$$




Wavelet $\psi_{\mathbf{N}_3}^{(\mathbf{J}_Y)}$, $\mathbf{N}_3 = \begin{pmatrix} 512 & 0 \\ 0 & 64 \end{pmatrix}$




Corresp. wavelet part

Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_Y \mathbf{J}_Y \mathbf{J}_X \mathbf{N}_4$$




Wavelet $\psi_{\mathbf{N}_4}^{(\mathbf{J}_X)}$, $\mathbf{N}_4 = \begin{pmatrix} 256 & 0 \\ 0 & 64 \end{pmatrix}$




Corresp. wavelet part

Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_Y \mathbf{J}_Y - \mathbf{J}_X \mathbf{N}_5$$




Wavelet $\psi_{\mathbf{N}_5}^{(\mathbf{J}_X)}$, $\mathbf{N}_5 = \begin{pmatrix} 256 & 32 \\ 0 & 64 \end{pmatrix}$




Corresp. wavelet part

Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_{Y-X} \mathbf{J}_Y \mathbf{J}_X \mathbf{N}_6$$




Wavelet $\psi_{\mathbf{N}_6}^{(\mathbf{J}_X)}$, $\mathbf{N}_6 = \begin{pmatrix} 256 & 64 \\ 0 & 64 \end{pmatrix}$




Corresp. wavelet part

Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_{Y-} \mathbf{J}_{Y-} \mathbf{J}_X \mathbf{N}_7$$




Wavelet $\psi_{\mathbf{N}_7}^{(\mathbf{J}_X)}$, $\mathbf{N}_7 = \begin{pmatrix} 256 & 96 \\ 0 & 64 \end{pmatrix}$




Corresp. wavelet part

Example of a decomposition

$$\mathbf{M} = \mathbf{J}_Y \mathbf{J}_{Y-} \mathbf{J}_{Y-} \mathbf{J}_X \mathbf{N}_7$$



Wavelet $\psi_{\mathbf{N}_7}^{(\mathbf{J}_X)}$, $\mathbf{N}_7 = \begin{pmatrix} 256 & 96 \\ 0 & 64 \end{pmatrix}$



Corresp. wavelet part

numbering: binary in matrices and directly in multiples of 32 in shear

Conclusion

Patterns $\mathcal{P}(\mathbf{M})$ and generating sets $\mathcal{G}(\mathbf{M}^T)$

- generalize equally spaced points
- still resemble an FFT
- fast wavelet transform based with corresponding TI spaces

The constructed wavelets generalize the onedimensional
de la Vallée Poussin wavelets

- to arbitrary dyadic scaling matrices
 - based on arbitrary smooth functions g
- ⇒ localization

Literature

B., *The fast Fourier transform and fast wavelet transform for patterns on the torus*, ACHA 35 (2013) 39–51.

B., *Translationsinvariante Räume multivariater anisotroper Funktionen auf dem Torus*, Dissertation, Universität zu Lübeck, 2013.

[LP10] D. Langemann, J. Prestin, *Multivariate periodic wavelet analysis*, ACHA 28 (2010) 46–66.

[MS03] I. E. Maximenko, M. A. Skopina, *Multivariate periodic wavelets*, St. Petersburg. Math. J. 15 (2003) 165–190.

[PT95] G. Plonka, M. Tasche, *On the computation of periodic spline wavelets*, ACHA 2 (1995) 1–14.

[Se95] K. Selig, *periodische Wavelet-Packets und eine gradoptimale Schauderbasis*, Dissertation, Universität Rostock, 1998.

Literature

B., *The fast Fourier transform and fast wavelet transform for patterns on the torus*, ACHA 35 (2013) 39–51.

B., *Translationsinvariante Räume multivariater anisotroper Funktionen auf dem Torus*, Dissertation, Universität zu Lübeck, 2013.

[LP10] D. Langemann, J. Prestin, *Multivariate periodic wavelet analysis*, ACHA 28 (2010) 46–66.

[MS03] I. E. Maximenko, M. A. Skopina, *Multivariate periodic wavelets*, St. Petersburg. Math. J. 15 (2003) 165–190.

[PT95] G. Plonka, M. Tasche, *On the computation of periodic spline wavelets*, ACHA 2 (1995) 1–14.

[Se95] K. Selig, *periodische Wavelet-Packets und eine gradoptimale Schauderbasis*, Dissertation, Universität Rostock, 1998.

Thank you for your attention.