



USER GUIDE

Access

This Maricultura Brazilian Aquaculture Planning tool was developed utilizing R (version). The application is coded with the “Shiny” package. It is accessible through RStudio and directly from a web browser. To run the Shiny App in RStudio, fork, clone, and download the `maricultura-app` repository on GitHub:

(<https://github.com/maricultura>).

To access from a web browser, use the url:

(<https://maricultura-gp.shinyapps.io/maricultura-app/>)

Tool Navigation

This tool is equipped with a Navigation bar. Upon opening the application, only the “About,” “Site Suitability,” “Metadata,” and “User Guide” tabs are available. These tabs can be accessed on the toolbar. Productivity and Profitability analyses are available, but only after running the primary site suitability analysis. This is because the productivity and profitability analyses are functions of the site suitability analysis outcomes.


Site Suitability

The site suitability analysis considers physical and infrastructural constraints, as well as biological thresholds, to determine which parcels within Brazil’s EEZ are available for marine aquaculture development. As a user, you can select different constraints and thresholds for your analysis. Some inputs are variable, and some are binary. Variable inputs that you can manipulate include: Sea Surface Temperature, Current Velocity, Minimum Dissolved Oxygen, Depth, and Maximum Distance to Shore. Inputs that can be selected to include or omit are: Marine Protected Areas, Artificial Reefs, Natural Reefs, Oil Pipelines, Oil Production, and Shipping Lanes.

You can navigate through the site suitability constraints within the two tabs in the left side panel of the “Site Suitability” tab. Once you have selected your inputs for variable and fixed constraints, press the “Run” button. This will load a map of suitable sites for marine aquaculture development in Brazil. You may also choose to download the map as a .tif image file, which can be uploaded for further analyses and visualization in spatial mapping programs, such as RStudio and ArcGIS.

Site Productivity

Once you have run the site suitability analysis, the productivity analysis will become available to you on the Navigation Bar. The tab is titled “Biomass.” This application feature will utilize the output from your site suitability analysis to determine which of



your generated sites will produce the most fish biomass. A piecewise linear function is used to estimate. The analysis uses a piecewise linear function to estimate how a species' growth rate varies spatially with temperature. The growth rates for the species of interest is then fused as a proxy for the Brody Growth Coefficient (K) in the von Bertalanffy model to generate an estimated length at harvest of individual fish for each parcel. These lengths at harvest are then converted to weights using an allometric growth conversion. The final biomass at harvest is estimated by multiplying the estimated weight for one individual by the stocking density, then subtracting the expected mortality during the grow out period.

Currently this application only features two species: cobia and Atlantic Salmon. Our hope is to integrate other presets of commonly cultured species that might be suitable for Brazil. For now, however, you may choose either of the species, run the analyses, and download the map. The output depicts the total possible mass of fish that could be produced in a cell per year.

Site Profitability

Here we put a general description like the two above

Map Features

- ❖ Coordinates
 - Latitude and Longitude coordinates are displayed in the top left corner of the map to identify location of specific parcels
 - Coordinates are dependent on cursor location
- ❖ Box zoom (press the `Shift` key while dragging the mouse)
 - This enables you to zoom into a particular region of interest
- ❖ Full screen
 - This generates a full-screen version of the output map
 - Click `Escape` to exit full-screen mode
- ❖ Reset zoom and initial view
 - Click to return to original output
- ❖ Zoom in/out buttons
 - Click to for general magnification/reduction of map
- ❖ Basemaps
 - There are two available basemaps
 - The first is a simple ESRI map showing country borders and names
 - The second displays major cities and regions boundaries



Equations

Include Equations that are used for analyses : Economic and Productivity

Variables

Species	a	b	Min SST	Max SST
Cobia				
Atlantic Salmon				
Gilted Seabream				

Implications

We hope that this tool can assist with planning strategies for the development of aquaculture. It can estimate the location of suitable sites, productivity, and profitability of mariculture in Brazil. A challenge in planning for aquaculture development is that one strategy does not fit all situations. Our tool addresses this challenge by allowing for the consideration of various scenarios. For instance, it can be used to simulate scenarios with different farmed species and cage specifications. In addition, strategic planning needs to be adaptive to be able to survive changing conditions. Our tool makes it possible to explore how varying different parameters affects the overall suitability of a location. For example, it can estimate how changes in the market price of a species can affect the profitability of farming it. Lastly, detailed planning efforts can benefit from developing their own customized tools. Our tool was designed to be easily adaptable to other regions and to include additional input parameters. The code used to build the tool is open source and can be used as a starting point for the creation of new tools that expand on our results. Researchers from any country can integrate their own physical constraints and EEZ geographic shapefiles into our published code and estimate where most suitable sites to implement aquaculture might be.