

ICS663: Pattern Recognition

Department of Information and Computer Sciences
University of Hawai'i at Manoa

Kyungim Baek

ICS663 (Fall 2015)

1

Announcement

- Homework assignment #1
 - Due: **Monday September 21, by 5:00 PM**
- Project proposal
 - Due: **Wednesday September 23, by 5:00 PM**
 - Presentation (~ 10 minutes)
 - Monday (9/28)
 - BJ, Thomas, Tyson, Sharif, Danny, Jeremy
 - Wednesday (9/30)
 - Tetsuya, Kelly, Nurit

ICS663 (Fall 2015)

2

Previously...

- Discriminant function for the minimum-error-rate classification

$$g_i(\mathbf{x}) = \ln p(\mathbf{x} | \omega_i) + \ln P(\omega_i)$$

- Case of multivariate normal, $p(\mathbf{x} | \omega_i) \sim N(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$:

$$p(\mathbf{x} | \omega_i) = \frac{1}{(2\pi)^{d/2} |\boldsymbol{\Sigma}_i|^{1/2}} \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu}_i)' \boldsymbol{\Sigma}_i^{-1} (\mathbf{x} - \boldsymbol{\mu}_i) \right]$$

$$g_i(\mathbf{x}) = -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu}_i)' \boldsymbol{\Sigma}_i^{-1} (\mathbf{x} - \boldsymbol{\mu}_i) - \frac{d}{2} \ln 2\pi - \frac{1}{2} \ln |\boldsymbol{\Sigma}_i| + \ln P(\omega_i)$$

ICS663 (Fall 2015)

3

Previously...

- Case 1: $\boldsymbol{\Sigma}_i = \sigma^2 \mathbf{I}$

$$g_i(\mathbf{x}) = -\frac{\|\mathbf{x} - \boldsymbol{\mu}_i\|^2}{2\sigma^2} + \ln P(\omega_i) = \mathbf{w}_i' \mathbf{x} + w_{i0}$$

$$\text{where } \mathbf{w}_i = \frac{\boldsymbol{\mu}_i}{\sigma^2}; \quad w_{i0} = -\frac{1}{2\sigma^2} \boldsymbol{\mu}_i' \boldsymbol{\mu}_i + \ln P(\omega_i)$$

- The hyperplane separating R_i and R_j (decision surface between R_i and R_j)

$$g_i(\mathbf{x}) = g_j(\mathbf{x}) \Rightarrow \mathbf{w}^i (\mathbf{x} - \mathbf{x}_0) = 0$$

$$\text{where } \mathbf{w} = \boldsymbol{\mu}_i - \boldsymbol{\mu}_j \text{ and } \mathbf{x}_0 = \frac{1}{2} (\boldsymbol{\mu}_i + \boldsymbol{\mu}_j) - \frac{\sigma^2}{\|\boldsymbol{\mu}_i - \boldsymbol{\mu}_j\|^2} \ln \frac{P(\omega_i)}{P(\omega_j)} (\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)$$

ICS663 (Fall 2015)

4

Previously...

- Case 2: $\Sigma_i = \Sigma$

$$g_i(\mathbf{x}) = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_i)' \Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu}_i) + \ln P(\omega_i) = \mathbf{w}_i' \mathbf{x} + w_{i0}$$

where $\mathbf{w}_i = \Sigma^{-1} \boldsymbol{\mu}_i$; $w_{i0} = -\frac{1}{2} \boldsymbol{\mu}_i' \Sigma^{-1} \boldsymbol{\mu}_i + \ln P(\omega_i)$

- The hyperplane separating R_i and R_j (decision surface between R_i and R_j)

$$g_i(\mathbf{x}) = g_j(\mathbf{x}) \Rightarrow \mathbf{w}'(\mathbf{x} - \mathbf{x}_0) = 0$$

where $\mathbf{w} = \Sigma^{-1}(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)$ and

$$\mathbf{x}_0 = \frac{1}{2}(\boldsymbol{\mu}_i + \boldsymbol{\mu}_j) - \frac{1}{(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)' \Sigma^{-1}(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)} \ln \frac{P(\omega_i)}{P(\omega_j)} (\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)$$

ICS663 (Fall 2015)

5

Lecture 5

- Discriminant Functions for the Normal Density
 - Case 3: $\Sigma_i = \text{Arbitrary}$
- Bayes Decision Theory for Discrete Features
 - Example: Independent Binary Features
- Parametric Density Estimation
 - Introduction

ICS663 (Fall 2015)

6

Case 3: $\Sigma_i = \text{Arbitrary}$

$$g_i(\mathbf{x}) = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_i)' \Sigma_i^{-1}(\mathbf{x} - \boldsymbol{\mu}_i) - \frac{d}{2} \ln 2\pi - \frac{1}{2} \ln |\Sigma_i| + \ln P(\omega_i)$$

independent of i

→ $g_i(\mathbf{x}) = \mathbf{x}' \mathbf{W}_i \mathbf{x} + \mathbf{w}_i' \mathbf{x} + w_{i0}$

where

$$\mathbf{W}_i = -\frac{1}{2} \Sigma_i^{-1}$$


$$\mathbf{w}_i = \Sigma_i^{-1} \boldsymbol{\mu}_i$$

$$w_{i0} = -\frac{1}{2} \boldsymbol{\mu}_i' \Sigma_i^{-1} \boldsymbol{\mu}_i - \frac{1}{2} \ln |\Sigma_i| + \ln P(\omega_i)$$

ICS663 (Fall 2015)

7

$\Sigma_i = \text{Arbitrary}$: Decision Surfaces




ICS663 (Fall 2015)

8

$\Sigma_i = \text{Arbitrary: Decision Surfaces}$

- Decision regions for four normal distributions



- See Example 1 in the textbook

ICS663 (Fall 2015)

9

$\Sigma_i = \text{Arbitrary: Example}$

- Samples:

$$\omega_1 : (1,2)^t, (3,1)^t, (5,2)^t, (3,3)^t$$

$$\omega_2 : (7,6)^t, (8,4)^t, (9,6)^t, (8,8)^t$$

- Compute sample mean and covariance for each class:

$$\mu = \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i \quad \Sigma = \frac{1}{N-1} \sum_{i=1}^N (\mathbf{x}_i - \mu)(\mathbf{x}_i - \mu)^t$$

$$\mu_1 = \frac{1}{4} \left(\begin{pmatrix} 1 \\ 2 \end{pmatrix} + \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} 5 \\ 2 \end{pmatrix} + \begin{pmatrix} 3 \\ 3 \end{pmatrix} \right) = \begin{pmatrix} 3 \\ 2 \end{pmatrix}, \quad \Sigma_1 = \begin{pmatrix} 8/3 & 0 \\ 0 & 2/3 \end{pmatrix}$$

$$\mu_2 = \frac{1}{4} \left(\begin{pmatrix} 7 \\ 6 \end{pmatrix} + \begin{pmatrix} 8 \\ 4 \end{pmatrix} + \begin{pmatrix} 9 \\ 6 \end{pmatrix} + \begin{pmatrix} 8 \\ 8 \end{pmatrix} \right) = \begin{pmatrix} 8 \\ 6 \end{pmatrix}, \quad \Sigma_2 = \begin{pmatrix} 2/3 & 0 \\ 0 & 8/3 \end{pmatrix}$$

ICS663 (Fall 2015)

10

$\Sigma_i = \text{Arbitrary: Example (cont'd)}$

- Discriminant functions:

$$\begin{aligned} g_1(\mathbf{x}) &= -\frac{1}{2} \mathbf{x}^t \Sigma_1^{-1} \mathbf{x} + \mu_1^t \Sigma_1^{-1} \mathbf{x} - \frac{1}{2} \mu_1^t \Sigma_1^{-1} \mu_1 - \frac{1}{2} \ln |\Sigma_1| + \ln P(\omega_1) \\ &= -\frac{1}{2} \begin{pmatrix} x_1 & x_2 \end{pmatrix} \begin{pmatrix} 3/8 & 0 \\ 0 & 3/2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 3 & 2 \end{pmatrix} \begin{pmatrix} 3/8 & 0 \\ 0 & 3/2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &\quad - \frac{1}{2} \begin{pmatrix} 3 & 2 \end{pmatrix} \begin{pmatrix} 3/8 & 0 \\ 0 & 3/2 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \end{pmatrix} - \frac{1}{2} \ln \begin{vmatrix} 8/3 & 0 \\ 0 & 2/3 \end{vmatrix} + \ln P(\omega_1) \\ &= -(3/16)x_1^2 - (3/4)x_2^2 + (9/8)x_1 + 3x_2 - 75/16 - (1/2)\ln(16/9) + \ln P(\omega_1) \end{aligned}$$

$$\begin{aligned} g_2(\mathbf{x}) &= -\frac{1}{2} \begin{pmatrix} x_1 & x_2 \end{pmatrix} \begin{pmatrix} 3/2 & 0 \\ 0 & 3/8 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 8 & 6 \end{pmatrix} \begin{pmatrix} 3/2 & 0 \\ 0 & 3/8 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &\quad - \frac{1}{2} \begin{pmatrix} 8 & 6 \end{pmatrix} \begin{pmatrix} 3/2 & 0 \\ 0 & 3/8 \end{pmatrix} \begin{pmatrix} 8 \\ 6 \end{pmatrix} - \frac{1}{2} \ln \begin{vmatrix} 2/3 & 0 \\ 0 & 8/3 \end{vmatrix} + \ln P(\omega_2) \\ &= -(3/4)x_1^2 - (3/16)x_2^2 + 12x_1 + (18/8)x_2 - 219/4 - (1/2)\ln(16/9) + \ln P(\omega_2) \end{aligned}$$

ICS663 (Fall 2015)

11

$\Sigma_i = \text{Arbitrary: Example (cont'd)}$


$$g_{12}(\mathbf{x}) = g_1(\mathbf{x}) - g_2(\mathbf{x}) = \frac{9}{16}x_1^2 - \frac{9}{16}x_2^2 - \frac{87}{8}x_1 + \frac{3}{4}x_2 + \frac{801}{16} + \ln P(\omega_1) - \ln P(\omega_2)$$

- Decision boundaries: $g_{12}(\mathbf{x}) = 0$

$$- P(\omega_1) = 0.8, P(\omega_2) = 0.2: 3x_1^2 - 3x_2^2 - 58x_1 + 4x_2 + 274.3936 = 0$$

$$- P(\omega_1) = 0.5, P(\omega_2) = 0.5: 3x_1^2 - 3x_2^2 - 58x_1 + 4x_2 - 267 = 0$$

$$- P(\omega_1) = 0.2, P(\omega_2) = 0.8: 3x_1^2 - 3x_2^2 - 58x_1 + 4x_2 + 259.6064 = 0$$



ICS663 (Fall 2015)

12

Bayes Decision Theory: Discrete Features

- Components of \mathbf{x} can take only one of m discrete values, v_1, v_2, \dots, v_m
- Case of *independent binary features* in two category problem:
 - Let $\mathbf{x} = (x_1, x_2, \dots, x_d)^t$ where each x_i is either 0 or 1, with probabilities:

$$p_i = P(x_i = 1 | \omega_1) \quad \text{and} \quad q_i = P(x_i = 1 | \omega_2)$$

$$P(\mathbf{x} | \omega_1) = P(x_1, x_2, \dots, x_d | \omega_1) = \prod_{i=1}^d P(x_i | \omega_1) = \prod_{i=1}^d p_i^{x_i} (1 - p_i)^{1-x_i}$$

$$P(\mathbf{x} | \omega_2) = P(x_1, x_2, \dots, x_d | \omega_2) = \prod_{i=1}^d P(x_i | \omega_2) = \prod_{i=1}^d q_i^{x_i} (1 - q_i)^{1-x_i}$$

ICS663 (Fall 2015)

13

Independent Binary Features

- Likelihood ratio:

$$\frac{P(\mathbf{x} | \omega_1)}{P(\mathbf{x} | \omega_2)} = \prod_{i=1}^d \left(\frac{p_i}{q_i} \right)^{x_i} \left(\frac{1-p_i}{1-q_i} \right)^{1-x_i}$$

- For minimum error-rate decision, the discriminant function in this case is:

$$\begin{aligned} g(\mathbf{x}) &= \ln \frac{P(\mathbf{x} | \omega_1)}{P(\mathbf{x} | \omega_2)} + \ln \frac{P(\omega_1)}{P(\omega_2)} \\ &= \sum_{i=1}^d \left[x_i \ln \frac{p_i}{q_i} + (1-x_i) \ln \frac{1-p_i}{1-q_i} \right] + \ln \frac{P(\omega_1)}{P(\omega_2)} \end{aligned}$$

ICS663 (Fall 2015)

14

Independent Binary Features

- The discriminant function is linear in the x_i :

$$g(\mathbf{x}) = \sum_{i=1}^d w_i x_i + w_0$$

where:

$$w_i = \ln \frac{p_i(1-q_i)}{q_i(1-p_i)} \quad i = 1, \dots, d$$

$$w_0 = \sum_{i=1}^d \ln \frac{1-p_i}{1-q_i} + \ln \frac{P(\omega_1)}{P(\omega_2)}$$

decide ω_1 if $g(\mathbf{x}) > 0$ and ω_2 if $g(\mathbf{x}) \leq 0$

ICS663 (Fall 2015)

15

Parametric Density Estimation

ICS663 (Fall 2015)

16

Recall: Bayes' Theorem

$$P(\omega_j | \mathbf{x}) = \frac{p(\mathbf{x} | \omega_j)P(\omega_j)}{p(\mathbf{x})}$$

- Interpretation: $\text{posterior} = \frac{\text{likelihood} \cdot \text{prior}}{\text{evidence}}$
- Requirements for Bayesian classification
 - $P(\omega_i)$: prior probability distributions
 - $p(\mathbf{x} | \omega_i)$: class-conditional densities (likelihoods)

ICS663 (Fall 2015)

17

Why Learn?

- Bayesian decision theory assumes complete knowledge about the probabilistic structure of the problem
- But, in practice we have access only to *partial information*
- Forms of partial information
 - **Training data** D_n : often noisy, partially missing
 - **Prior knowledge** about probability distributions
 - **Dependence structure** in data
- Objective: **Estimate** $P(\omega_i | \mathbf{x}) \leftarrow P_n(\omega_i | \mathbf{x})$

ICS663 (Fall 2015)

from R. Meir
18

Estimation Problem

- **Data**: collection of samples to be classified into c classes $\omega_1, \omega_2, \dots, \omega_c$ and divided into data sets (according to class) as follows D_1, D_2, \dots, D_c (assume supervised learning)
- **Needed**:
 - Prior probabilities $P(\omega_i)$
 - Class-conditional probabilities $p(\mathbf{x} | \omega_i)$

ICS663 (Fall 2015)

19

Estimation Problem (cont'd)


- **Estimation problem**:
 - Estimating $P(\omega_i)$ is not very difficult in the context of supervised learning. For example, assume

$$P(\omega_i) = \frac{\# \text{ of times class } i \text{ appears in sample}}{\text{sample size}}$$
 - Estimating $p(\mathbf{x} | \omega_i)$ is hard, especially in high-dimensional feature spaces and when the number of training samples available seems to be too small
 - Assumption: $p(\mathbf{x} | \omega_i) = p(\mathbf{x} | \omega_i, \theta) - \text{known}$ parametric form, \exists a 'true' θ_0

ICS663 (Fall 2015)

20

Parametric Classification



$$p(\mathbf{x} | \omega_i, \theta) + D_n \xrightarrow{(1)} \theta_n \xrightarrow{(2)} P_n(\omega_i | \mathbf{x}) \xrightarrow{(3)} \alpha_n(\mathbf{x})$$

- (1) θ_n parameter estimate
 (2) $P_n(\omega_i | \mathbf{x})$ posterior estimate
 (3) $\alpha_n(\mathbf{x})$ empirical classifier

ICS663 (Fall 2015)

from R. Meir
21

Parameter Estimation

- Simplifying assumptions:
 - Feature independence
 - Samples have been drawn independently (i.e. samples follow the i.i.d. model – independent and identically distributed random variables)

ICS663 (Fall 2015)

22

Estimation Methods

- **Maximum-Likelihood (ML) estimation:** *parameters of probabilistic distributions are fixed but unknown values*
 - Parameters are unknown constants to be identified through training
 - Best estimate of parameter values is achieved by maximizing the probability of obtaining the samples observed
- **Bayesian estimation:** *parameters are random variables that follow a known (e.g. Gaussian) distribution*
 - Parameters as random variables having some known prior distribution
 - By observing the behavior of the training samples, the posterior probabilities inferred help revising the parameter values
 - The more the training samples, the better the chances of refining the posterior probabilities, and subsequently, to peak the parameter values

ICS663 (Fall 2015)

from S. Iliescu
23

Estimation Methods (cont'd)

- Differences between ML and Bayesian estimation
 - Conceptual (philosophical): are parameters constants or random variables?
 - Different approaches, but usually leading to the same results (when sufficient training samples are available)

ICS663 (Fall 2015)

from S. Iliescu
24