| KEEN | GOM | BENTHIC | MONITORING | DATA |
|------|-----|---------|------------|------|

FISH COUNTS

| Scanned by: | |
|---------------|------------|
| Date scanned: | <u>\\\</u> |

| SITE: | Pema | auid |
|-------|------|------|
| TR: | L | 4 |

DATE: 8 / 8 / 16
Observer: BM

Depth (m) at 0m:_ Depth (m) at 40m:_ Visibility (m): _ Temperature (C):__

| | | | YOY | 0-10cm | 10-50cm | 50-100cm | >100cm | TOTAL |
|------------------------------------|---|------|---------|--------|---------|----------|---|-----------------|
| Cunner | (Tautogolabrus adsperus) | TAAD | 1-0.093 | | 2 | | | |
| Tautog | (Tautoga onitis) | TAON | | | | | | |
| Tautog Black Sea Bass Striped Bass | (Centropristis striata) | CEST | | | | | | |
| Striped Bass | (Morone saxatalis) | MOSA | | | | | | |
| Pollock | (Pollachius virens) | POVI | | | | | | |
| Spotted Hake | (Urophyicus regia) | URRE | | | | | | |
| Summer Flounder | (Paralichthrus denatus) L-eyed | PADE | | | | | | |
| Winter Flounder | (Pseudopleuronectes americanus) R-eyed. | PSAM | | | | | | |
| Windowpane Flound | leı (Scopthalmus aquosus) | SCAQ | | | | | | |
| Fourspot Flounder | (Paralichthys oblongus) | PAOB | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | 14 11 11 11 11 | |
| | | | * | | | | | |
| | | | | | | | | To City Service |
| | | | | | | | | |
| | | | - | | | | | |
| | | | | | | | | |
| | | | | | | | 450000000000000000000000000000000000000 | \$21/9090000 |
| | | | | | | | | |
| | | | | | | | | |

*no cryptic fish counted ie: gunnels, grubbys, sculpins, sea raven

Wolffish ANLU GAMO Cod (barbel) PRCA Northern Sea Robin Acadian Redfish SEFA SYFU Northern Pipefish MEAE Haddock MEBI Silver hake MAAM Ocean Pout Checked by Date Checked: _____

STCH Scup/Porgy ANRO American Eel POSA Bluefish SPMA Northern Puffer **LEER** Little Skate **LEOC** Winter Skate SQAC Spiny Dogfish

CATA Sand Tiger Shark