

Subsurface Kelp  
MorphometricsScanned by: Kelsey Schultz  
Date scanned: Kelsey SchultzSITE: CANOE  
BEACHDATE 8 / 27 / 15Observer: ETHAN EDSON

Collect 1 kelp of each sp. Every 4 m

Kelps

AGCL Agarum clathratum (shotgun kelp)

ALES Alaria esculenta (distinctive midrib)

LADI Laminaria digitata

SL Saccharina spp. (lasagne)

#	Transect	SPECIES	Blade Length (cm)	Widest Width (cm)	Stipe Length (cm)
1	2	SL	96.5	14	3.5
2		SL	27	8.5	1.6
3		SL	34.5	15.5	6.0
4		SL	32	8.5	—
5		SL	19	7	—
6		SL	23	7	1.0
7		SL	41	14	1.0
8		SL	30.5	8	1.0
9		SL	19	5.5	0.5
10		SL	12.5	5	0.5
11					
12					
13					
14					
15					
16					
17					
18					
19					
20					
21					
22					
23					
24					
25					
26					
27					
28					
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					

LADI width is taken with all blades placed together

Data entry by: \_\_\_\_\_  
Date entered: \_\_\_\_\_Checked by: \_\_\_\_\_  
Date checked: \_\_\_\_\_