

✓ ¡Felicitaciones! ¡Aprobaste!  
PARA APROBAR 75 % o más

Continúa aprendiendo

CALIFICACIÓN  
100 %

## Practice quiz onTangent Lines to Functions

PUNTOS TOTALES DE 2

1. Suppose that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a function. Which of the following expressions corresponds to  $f'(2)$ , the slope of the tangent line to the graph of  $f(x)$  at  $x = 2$ ?

1 / 1 punto


- ☒  $f'(2) = \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h}$
- ☐  $f'(2) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$
- ☐  $f'(2) = mx + b$
- ☐  $f'(2) = 2$

✓ Correcto

This expression can be obtained from the first screen of our video by plugging in 2 for  $a$ .

2. Suppose that  $h : \mathbb{R} \rightarrow \mathbb{R}$  is a function whose graph is shown as the blue curve in the figure. For how many values of  $a$  is  $h'(a) = 0$ ?

1 / 1 punto



- ☐ 3
- ☐ Never
- ☐ Always
- ☒ 2

✓ Correcto

$h'(a)$  gives the slope of the tangent line to the graph of  $h$  at the point  $x = a$ .



¡Felicitaciones! ¡Aprobaste!

PARA APROBAR 75 % o más

Continúa aprendiendo

CALIFICACIÓN

91,66 %

## Practice quiz on Exponents and Logarithms

PUNTOS TOTALES DE 12

1. Re write the number  $784 = 2 \times 2 \times 2 \times 2 \times 7 \times 7$  using exponents.

1 / 1 punto

- ☐  $(2 \times 7)^6$
- ☐  $(2^6)(7^6)$
- ☐  $(16^4)(49^2)$
- ☒  $(2^4)(7^2)$

✓ Correcto

For this type of problem, count the number of times each relevant factor appears in the product. That number is the exponent for that factor.



2. What is  $(x^2 - 5)^0$ ?

1 / 1 punto

- ☐  $(x^2)$
- ☐  $(x^2) - 5$
- ☒ 1
- ☐  $-4$

✓ Correcto

Any real number (except zero) raised to the "zeroith" power = 1.

3. Simplify  $((x - 5)^2)^{-3}$

1 / 1 punto

- ☒  $(x - 5)^{-6}$
- ☐  $(x - 5)$
- ☐  $(x - 5)^{-1}$
- ☐  $(x - 5)^{-5}$



## Prueba de práctica sobre exponentes y logaritmos

Cuestionario Práctico • 40 min

4. Simplify  $(\frac{8^2}{8^7})^2$

1 / 1 punto

- ☐  $8^{-1}$
- ☐  $8^{-5}$
- ☐  $8^{-4}$
- ☒  $8^{-10}$

✓ **Correcto**

We can first simplify what is inside the parenthesis to  $8^{-5}$  using the Division and Negative Powers Rule.

Then apply division and negative powers-- the result is the same.  $\frac{8^4}{8^{14}} = 8^{-10}$

5.  $\log 35 = \log 7 + \log x$

1 / 1 punto

Solve for  $x$



## Prueba de práctica sobre exponentes y logaritmos

Cuestionario Práctico • 40 min

5.  $\log 35 = \log 7 + \log x$

1 / 1 punto

Solve for  $x$

- ☐ 4
- ☒ 5
- ☐ 28
- ☐ 7

✓ **Correcto**

$$\log(x) = \log 35 - \log 7$$

$$\log(x) = \log\left(\frac{35}{7}\right)$$

By the Quotient Rule  $\log x = \log 5$

6.  $\log_2(x^2 + 5x + 7) = 0$

0 / 1 punto



## Prueba de práctica sobre exponentes y logaritmos

Cuestionario Práctico • 40 min

6.  $\log_2(x^2 + 5x + 7) = 0$

0 / 1 punto

Solve for  $x$

- ☒  $x = 2$  or  $x = 3$
- ☐  $x = 2$
- ☐  $x = -2$  or  $x = -3$
- ☐  $x = 3$

✗ Incorrecto

We use the property that  $b^{\log_b a} = a$

Use both sides as exponent for 2.

$$2^{\log_2 x^2 + 5x + 7} = 2^0$$

$$x^2 + 5x + 7 = 1$$

$$x^2 + 5x + 6 = 0$$



## Prueba de práctica sobre exponentes y logaritmos

Cuestionario Práctico • 40 min

7. Simplify  $\log_2 72 - \log_2 9$

1 / 1 punto

- ☐  $\log_2 4$
- ☐ 4
- ☐  $\log_2 63$
- ☒ 3

✓ Correcto

By the quotient rule, this is  $\log_2 \frac{72}{9} = \log_2 2^3 = 3$

8. Simplify  $\log_3 9 - \log_3 3 + \log_3 5$

1 / 1 punto

- ☐ 8
- ☐ 15
- ☒  $\log_3 15$

9. Simplify  $\log_2(3^8 \times 5^7)$

1 / 1 punto

- ☐  $(5 \times \log_2 3) + (8 \times \log_2 5)$
- ☒  $(8 \times \log_2 3) + (7 \times \log_2 5)$
- ☐  $56 \times \log_2 15$
- ☐  $15 \times \log_2 56$

✓ **Correcto**

We first apply the Product Rule to convert to the sum:  $\log_2(3^8) + \log_2(5^7)$ . Then apply the power and root rule.

10. If  $\log_{10} y = 100$ , what is  $\log_2 y = ?$

1 / 1 punto

- ☒ 332.19
- ☐ 301.03
- ☐ 20
- ☐ 500

11. A tree is growing taller at a continuous rate. In the past 12 years it has grown from 3 meters to 15 meters. What is its rate of growth per year?

1 / 1 punto

- ☒ 13.41%
- ☐ 11.41%
- ☐ 10.41%
- ☐ 12.41%

✓ **Correcto**

$$\frac{\ln \frac{15}{3}}{12} = 0.1341$$

12. Bacteria can reproduce exponentially if not constrained. Assume a colony grows at a continually compounded rate of 400% per day. How many days before a colony with initial mass of  $6.25 \times 10^{-10}$  grams weights 1000 Kilograms?

1 / 1 punto

- ☐ 875 days
- ☒ 8.75 days



¡Felicitaciones! ¡Aprobaste!

PARA APROBAR 75 % o más

Continúa aprendiendo

CALIFICACIÓN

100 %

# Graded quiz on Tangent Lines to Functions, Exponents and Logarithms

CALIFICACIÓN DEL ÚLTIMO ENVÍO

100%

1. Convert  $\frac{1}{49}$  to exponential form, using 7 as the factor.

1 / 1 punto

☒  $7^{-2}$

☐  $(7^2)$

☐  $49^{-1}$

☐  $\frac{7}{7^3}$



## Prueba de práctica sobre exponentes y logaritmos

Cuestionario Práctico • 40 min

12. Bacteria can reproduce exponentially if not constrained. Assume a colony grows at a continually compounded rate of 400% per day. How many days before a colony with initial mass of  $6.25 \times 10^{-10}$  grams weights 1000 Kilograms?

1 / 1 punto

☐ 875 days

☒ 8.75 days

☐ 0.875 days

☐ 87.5 days



Correcto

$$6.25 \times 10^{-10} \times e^{4t} = 10^6$$

$$4t = \ln \left( \frac{10^6}{(6.25 \times 10^{-10})} \right) = 35.00878$$

$$t = \ln \frac{10^6}{6.25 \times 10^{-10}} = 8.752195$$





4. Simplify  $[(x^4)(y^{-6})]^{-1}$

1 / 1 punto

- ☐  $(x^3)(y^{-7})$
- ☒  $(x^{-4})(y^6)$
- ☐  $\frac{(x^4)}{(y^{-6})}$
- ☐  $\frac{(x-4)}{(y^6)}$

✓ Correcto

By the Power to a Power Rule, each of the exponents is multiplied by  $(-1)$

5. Solve for  $x$ :

1 / 1 punto

$\log_2(39x) - \log_2(x - 5) = 4$

- ☐  $\frac{80}{38}$
- ☐ 23



2. A light-year (the distance light travels in a vacuum in one year) is 9, 460 trillion meters. Express in scientific notation.

1 / 1 punto

- ☐  $0.946 \times 10^{16}$
- ☐  $9460 \times 10^{12}$  meters
- ☒  $9.46 \times 10^{15}$  meters.
- ☐  $9.46 \times 10^{15}$  kilometers

✓ Correcto

9, 460 is  $(9.4 \times 10^3)$  meters and one trillion meters is  $10^{12}$  meters.  $(9.4 \times 10^3)(10^{12}) = 9.4 \times 10^{15}$ . A kilometer is 1000 meters.

3. Simplify  $(x^8)(y^3)(x^{-10})(y^{-2})$

1 / 1 punto

- ☐  $(x^2)(y)$
- ☐  $(x)(y^{-2})$
- ☐  $(x^{-80})(y^{-6})$
- ☒  $(x^{-2})(y)$



5. Solve for  $x$ :

1 / 1 punto

$$\log_2(39x) - \log_2(x - 5) = 4$$

- ☐  $\frac{80}{38}$
- ☐  $\frac{23}{80}$
- ☒  $\frac{-80}{23}$
- ☐  $\frac{39}{23}$



Correcto

$\log_2 \frac{39x}{(x - 5)} = 4$  by the Quotient Rule.

Since both sides are equal, we can use them as exponents in an equation.

$$2^{\log_2 \frac{39x}{(x - 5)}} = 2^4$$

$39x$



6. Simplify this expression:

1 / 1 punto

$$\left(x^{\frac{1}{2}}\right)^{\frac{-3}{2}}$$

- ☒  $x^{\frac{-3}{4}}$
- ☐  $x^{\frac{1}{3}}$
- ☐  $x^{-1}$
- ☐  $x^{\frac{4}{3}}$



Correcto

We use the Power to a Power Rule -- multiply exponents:

$$x^{\frac{1}{2} \times \frac{-3}{2}} = x^{\frac{-3}{4}}$$

7. Simplify  $\log_2 8 - \log_2 4 - (\log_3 4.5 + \log_3 2)$

1 / 1 punto





7. Simplify  $\log_2 8 - \log_2 4 - (\log_3 4.5 + \log_3 2)$

1 / 1 punto

- ☐ 1
- ☐ 0
- ☒ -1
- ☐ 2

✓ Correcto

This is equivalent to:

$$\log_2\left(\frac{8}{4}\right) - \log_3(4.5 \times 2) = 1 - 2 = -1$$

8. If  $\log_3 19 = 2.680$ , what is  $\log_9 19$ ?

1 / 1 punto

- ☐ 0.8934
- ☐ 5.216
- ☐ 1.304



8. If  $\log_3 19 = 2.680$ , what is  $\log_9 19$ ?

1 / 1 punto

- ☐ 0.8934
- ☐ 5.216
- ☒ 1.304
- ☐ 0.4347

✓ Correcto

To convert from  $\log_3$  to  $\log_9$ , divide by  $\log_3 9$ . Which is equal to 2, so the answer is 1.34

9. If  $\log_{10} b = 1.8$  and  $\log_a b = 2.5752$ , what is  $a$ ?

1 / 1 punto

- ☐ 6
- ☒ 5
- ☐ 3



9. If  $\log_{10} b = 1.8$  and  $\log_a b = 2.5752$ , what is  $a$ ?

1 / 1 punto

- ☐ 6
- ☒ 5
- ☐ 3
- ☐ 4

✓ Correcto

To solve for  $a$  in the formula;

$$\log_a b = \frac{\log_x b}{\log_x a}$$

$$\log_a b = 2.5752 \text{ and } \log_{10} b = 1.8$$

$$\text{Therefore, } \log_{10} a \text{ must equal to } \frac{1.8}{2.5752} = 0.69897$$



10. An investment of 1,600 is worth 7,400 after 8.5 years. What is the continuously compounded rate of return of this investment?

1 / 1 punto

- ☐ 17.01%
- ☐ 19.01%
- ☐ 20.01
- ☒ 18.02%

✓ Correcto

$$\frac{\ln \frac{7400}{1600}}{8.5} = 0.18017$$

11. A pearl grows in an oyster at a continuously compounded rate of .24 per year. If a 25-year old pearl weighs 1 gram, what did it weigh when it began to form?

1 / 1 punto

- ☒ 0.002478



11. A pearl grows in an oyster at a continuously compounded rate of .24 per year. If a 25-year old pearl weighs 1 gram, what did it weigh when it began to form?

1 / 1 punto

- ☒ 0.002478
- ☐ 0.0002478
- ☐ 0.02478
- ☐ 0.2478

✓ Correcto

$$e^{(0.24 \times 25)} = \frac{1}{x}$$

$$x = \frac{1}{(e^{0.24 \times 25})}$$

1



12.  $\log_2 z = 6.754$ . What is  $\log_{10}(z)$ ?

1 / 1 punto

- ☐ 0.49185
- ☐ 1.3508
- ☐ 0.82956
- ☒ 2.03316

✓ Correcto

$$\frac{\log_2 z}{\log_2 10} =$$

$$(\log_{10} z) \times (\log_2 10) = 3.321928$$

$$\text{Therefore, } \log_{10} z = \frac{6.754}{3.321928} = 2.03316$$

13. Suppose that  $a : \mathbb{R} \rightarrow \mathbb{R}$  is a function and that  $a(1) = 10$ . Suppose that  $a'(a)$  is negative for every single

1 / 1 punto



$$(\log_{10} z) \times (\log_2 10) = 3.321928$$

$$\text{Therefore, } \log_{10} z = \frac{6.754}{3.321928} = 2.03316$$

13. Suppose that  $g : \mathbb{R} \rightarrow \mathbb{R}$  is a function, and that  $g(1) = 10$ . Suppose that  $g'(a)$  is negative for every single value of  $a$ . Which of the following could possibly be  $g(1.5)$ ?

1 / 1 punto

- ☐  $g(1.5) = 10.1$
- ☐  $g(1.5) = 103.4$
- ☒  $g(1.5) = 9.7$
- ☐  $g(1.5) = 11$

✓ **Correcto**

Since the slope of the tangent line to the graph of  $g$  is negative everywhere on the graph, we know that  $g$  is *decreasing* function! And therefore we must have  $g(1.5) < g(1)$ . That is the case here, so this value is at least possible.