# A Topography of Climate Change Research

Max Callaghan



February 5, 2018

1 / 25





- To contribute evidence-based policy-making on climate change, the IPCC aims to comprehensively assess
- These assessments should be aim to balance legitimacy, credibility and relevance (?)

### Motivation



#### Motivation - Update 2018




Now that 2017 has ended, I have update the query to include papers from 2017 as well.

These extra 36,000 papers take the total number of documents to 309,697.

### Approach - Words, words, words





- Topic modelling is a way of reducing the dimensionality of a corpus of documents
- A large matrix of documents x words is factorised by a matrix of topics x words and a matrix of topics x documents (?)
- Topics describe the latent structure of the document corpus






Figure: A topic model of 3495 documents on climate change from the year 2000



$$V_{i\mu} \approx (WH)_{i\mu} = \sum_{a=1}^{r} W_{ia} H_{a\mu}$$

V: 8769 x 3495




Figure: A topic model of 3495 documents on climate change from the year 2000

Max Callaghan (MCC)

Climate Topography

February 5, 2018 7 / 25

W: 8769 x 50

#### Research Questions



- What is the thematic structure of the literature on climate change, and how has this changed over the five assessment periods of the IPCC
- What can this modelled thematic structure tell us about the past and future relationship between the IPCC and scientific literature on climate change?



The topic models above assume that the topics, and the words that make them up, are stable over time. Two approaches to better model dynamic topics:



The topic models above assume that the topics, and the words that make them up, are stable over time. Two approaches to better model dynamic topics:

 Dynamic Topic Modelling (DTM) (?) assume that a constant number of topics exists over all topic models, but allows the words in the topics to evolve from one time period to another



The topic models above assume that the topics, and the words that make them up, are stable over time. Two approaches to better model dynamic topics:

- Dynamic Topic Modelling (DTM) (?) assume that a constant number of topics
  exists over all topic models, but allows the words in the topics to evolve from one
  time period to another
- Dynamic Non-negative Matrix Factorisation (?) has varying numbers of topics in each window and allows for topics to emerge and/or disappear.



The topic models above assume that the topics, and the words that make them up, are stable over time. Two approaches to better model dynamic topics:

- Dynamic Topic Modelling (DTM) (?) assume that a constant number of topics
  exists over all topic models, but allows the words in the topics to evolve from one
  time period to another
- Dynamic Non-negative Matrix Factorisation (?) has varying numbers of topics in each window and allows for topics to emerge and/or disappear.

Where the size and variety of the literature we want to model has increased exponentially, we need an approach that allows for the emergence of new topics.



#### Window Topics: 1991






#### Window Topics: 1992











### Window Topics: 1993






83		100	F. 188. 9	
	200	27 30	4	
34	S Street	3.35	130	
4	34.076			



















### Dynamic NMF - application to climate change




- Choosing the number of window topics is non-trivial. Data-driven approaches are limited (see below), and human selection is time consuming.
- To facilitate the description of trends over the assessment periods of the IPCC, and to minimize the number of modelling decisions, I consider each IPCC assessment period as a time window.

# Dynamic NMF - application to climate change



- Choosing the number of window topics is non-trivial. Data-driven approaches are limited (see below), and human selection is time consuming.
- To facilitate the description of trends over the assessment periods of the IPCC, and to minimize the number of modelling decisions, I consider each IPCC assessment period as a time window.

 Starting from a logarithmic relationship between the number of documents and the ideal topic number, I compare 5 runs with varying numbers of topics for each window



## Dynamic NMF - number of topics



#### Human topic number criteria

Intelligibility

#### Data-driven topic number criteria

- Reconstruction accuracy
- Predictive capacity

## Dynamic NMF - number of topics





#### Human topic number criteria

Intelligibility

#### Data-driven topic number criteria


- Reconstruction accuracy
- Predictive capacity





### Preliminary results - structure





- A network of comprehensible topics is generated with 100 topics
- Topics can be matched to the IPCC working group from which the majority of the topic documents are referenced in
- Topics from the same working group are significantly more likely to be correlated with each other than those which are not

# Preliminary results - Inter-working Group topics - WG 1 and 3 $\,$



IPCC Coverage	Primary WG	Topic Title	WG 1	WG 2	WG 3
0.16%	1	{rainfal, monsoon, rain}	0.50%	0.50%	0.00%
0.10%	2	{veget, ndvi, cover}	0.41%	0.59%	0.00%
0.16%	1	{snow, cover, winter}	0.59%	0.41%	0.00%
0.17%	2	{region, local, scale}	0.41%	0.59%	0.00%
0.16%	1	{coastal, mangrov, rise}	0.57%	0.42%	0.01%

# Preliminary results - Inter-working Group topics - WG 1 and 3



IPCC Coverage	Primary WG	Topic Title	WG 1	WG 2	WG 3
0.09%	3	{gas, coal, greenhous}	0.30%	0.15%	0.56%
0.10%	3	{transport, vehicl, road}	0.24%	0.12%	0.64%
0.13%	1	{emiss, reduct, reduc}	0.45%	0.21%	0.34%
0.09%	1	{methan, oxid, methanotroph}	0.63%	0.16%	0.20%
0.13%	3	{ghg, greenhous, gas}	0.15%	0.09%	0.75%

# Preliminary results - Inter-working Group topics - WG 2 and 3 $\,$




IPCC Coverage	Primary WG	Topic Title	WG 1	WG 2	WG 3
0.11%	2	{sustain, develop, resourc}	0.04%	0.51%	0.46%
0.08%	3	{build, construct, design}	0.03%	0.38%	0.59%
0.11%	2	{environment, impact, life}	0.06%	0.58%	0.36%
0.19%	3	{polici, tax, govern}	0.02%	0.32%	0.66%
0.16%	2	{urban, citi, plan}	0.07%	0.55%	0.38%



How can we get a sense of which topics are better covered in IPCC reports?


- Each document d either matches or does not match an IPCC reference
- ullet For each topic h, we can sum the scores for each category of document
- The "IPCC proportion" of each topic is the proportion of the sum of the document score accounted for by documents which match IPCC references.






- The physical science aspects of climate change, as well topics on impacts, adaptation and scenarios are well covered by the IPCC
- Topics on specific technological solutions (particularly NETs), as well as soils, are less well covered









#### Next steps



- Adjusting second stage of model to give more weight to topics with many documents (to better capture emergence)
- Further analysis of new model
- (Manual) comparison of (AR6) topics with AR6 outline

### Frame Title

