

Prop. Let U, V, W be f.d.v.s. over F . Let $m = \dim(U)$, $n = \dim(V)$, $p = \dim(W)$, and let α, β, γ be ordered bases for U, V, W respectively.

(i) If $\sigma \in \mathcal{L}(V, U)$ has matrix $A = {}_{\alpha}[\sigma]_{\beta}$ and $\tau \in \mathcal{L}(W, V)$ has matrix $B = {}_{\beta}[\tau]_{\gamma}$, then $\sigma\tau \in \mathcal{L}(W, U)$ has matrix $C = {}_{\alpha}[\sigma\tau]_{\gamma}$ given by the matrix product AB .

(ii) If $\sigma \in \mathcal{L}(V, U)$ has matrix $A = {}_{\alpha}[\sigma]_{\beta}$ and $y \in V$ has β -coordinate vector

$$[y]_{\beta} = \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{bmatrix},$$

then $\sigma(y) \in U$ has α -coordinate vector given by

$$[\sigma(y)]_{\alpha} = A [y]_{\beta}.$$

Proof. (i) Follows from previous discussion.

(ii) To find $[\sigma(y)]_{\alpha}$, we write $\sigma(y)$ as a linear combination of the elements in the basis $\alpha = (e_1, \dots, e_m)$. Say $\beta = (f_1, \dots, f_n)$ and $A = (a_{ij})$. Then

$$\begin{aligned} \sigma(y) &= \sigma\left(\sum_{j=1}^n \lambda_j f_j\right) \\ &= \sum_{j=1}^n \lambda_j \sigma(f_j) \\ &= \sum_{j=1}^n \lambda_j \left(\sum_{i=1}^m a_{ij} e_i\right) \\ &= \sum_{j=1}^n \sum_{i=1}^m a_{ij} \lambda_j e_i \\ &= \sum_{i=1}^m \left(\sum_{j=1}^n a_{ij} \lambda_j\right) e_i \end{aligned}$$

$$So \quad [Ay]_{\alpha} = \begin{bmatrix} \sum a_{1j} \lambda_j \\ \vdots \\ \sum a_{mj} \lambda_j \end{bmatrix} = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{bmatrix} = A[y]_{\beta}.$$

□

Properties of matrix multiplication

Prop.

- (i) Matrix multiplication is associative; i.e., if $A \in M_{m \times n}(F)$, $B \in M_{n \times p}(F)$, $C \in M_{p \times q}(F)$, then

$$A(BC) = (AB)C.$$

- (ii) Matrix multiplication is distributive; i.e., if $A, B \in M_{m \times n}(F)$, $C, D \in M_{n \times p}(F)$, then

$$A(C+D) = AC + AD$$

$$(A+B)C = AC + BC.$$

- (iii) For each positive integer r , define the $r \times r$ identity matrix as the matrix $I_r \in M_r(F)$ with all diagonal entries equal to 1 and all other entries equal to zero (i.e., $I_r = (\delta_{ij})$). Then, for all $A \in M_{m \times n}(F)$,

$$I_m A = A = A I_n.$$

Proof 1 (By calculation) (i) Say $A = (a_{ij})$, $B = (b_{jk})$, $C = (c_{ke})$.

Then

$$\begin{aligned}(A(BC))_{i,e} &= \sum_{j=1}^n a_{ij} (BC)_{j,e} \\&= \sum_{j=1}^n a_{ij} \left(\sum_{k=1}^p b_{jk} c_{ke} \right) \\&= \sum_{j=1}^n \sum_{k=1}^p a_{ij} b_{jk} c_{ke} \\&= \sum_{k=1}^p \sum_{j=1}^n a_{ij} b_{jk} c_{ke} \\&= \sum_{k=1}^p \left(\sum_{j=1}^n a_{ij} b_{jk} \right) c_{ke} \\&= \sum_{k=1}^p (AB)_{i,k} c_{ke} \\&= ((AB)C)_{i,e}\end{aligned}$$

(ii), (iii) Exercise.

□

Proof 2 (Using structure of linear maps)

(i) Consider the diagram

$$F^q \xrightarrow{I} F^p \xrightarrow{\sigma} F^n \xrightarrow{\rho} F^m$$

where ρ, σ, I have matrices A, B, C for the standard bases. Then

$$\begin{aligned} A(BC) &= \begin{pmatrix} [\rho]_{E_m} & [\sigma]_{E_p} & [I]_{E_q} \end{pmatrix} \begin{pmatrix} [\sigma]_{E_p} & [I]_{E_q} \end{pmatrix} \\ &= \begin{pmatrix} [\rho]_{E_m} & [\sigma]_{E_p} \end{pmatrix} \begin{pmatrix} [\sigma]_{E_p} & [I]_{E_q} \end{pmatrix} \\ &= \begin{pmatrix} [\rho(\sigma)]_{E_q} \end{pmatrix} \\ &= \begin{pmatrix} [(\rho\sigma)]_{E_q} \end{pmatrix} \\ &= \begin{pmatrix} [\rho\sigma]_{E_p} & [I]_{E_q} \end{pmatrix} \\ &= \begin{pmatrix} [\rho]_{E_m} & [\sigma]_{E_p} \end{pmatrix} \begin{pmatrix} [I]_{E_q} \end{pmatrix} \\ &= (AB)C. \end{aligned} \quad \left| \begin{array}{l} E_r = \text{standard ordered} \\ \text{basis of } F^r \end{array} \right.$$

(ii) Let $\sigma_1: F^n \rightarrow F^m$, $\sigma_2: F^n \rightarrow F^m$, $I_1: F^p \rightarrow F^n$, $I_2: F^p \rightarrow F^n$ be the linear maps with matrices A, B, C, D respectively in the standard bases. Then

$$(A+B)C = ([\sigma_1] + [\sigma_2]) [\tau_1]$$

$$= [\sigma_1 + \sigma_2] \cdot [\tau_1]$$

$$= [(\sigma_1 + \sigma_2)\tau_1]$$

$$= [\sigma_1\tau_1 + \sigma_2\tau_1]$$

$$= [\sigma_1\tau_1] + [\sigma_2\tau_1]$$

$$= [\sigma_1][\tau_1] + [\sigma_2][\tau_1]$$

$$= AC + BC.$$

Similarly, $A(C+D) = AC + AD.$

(iii) I_n represents the identity map in standard basis $e_n = (e_1, \dots, e_n)$

$$\begin{aligned} \mathcal{A}_{F^n} : F^n &\longrightarrow F^n \\ e_j &\longmapsto \sum_{i=1}^n \delta_{ij} e_i = e_j. \end{aligned}$$

Let $\sigma : F^n \longrightarrow F^m$ be the linear map with matrix A in the standard bases. Then

$$A I_n = [\sigma][\mathcal{A}_{F^n}]$$

$$= [\sigma \mathcal{A}_{F^n}]$$

$$= [\sigma]$$

$$= A.$$

Similarly $I_m A = A.$

Examples

Def. Let $n \in \mathbb{Z}_{\geq 0}$ and let F be a field. Let $P_n(F)$ denote the set of all polynomials in the formal variable X of degree at most n ; i.e.,

$$P_n(F) = \{a_0 + a_1 X + \dots + a_n X^n \mid a_0, a_1, \dots, a_n \in F\}.$$

Then $P_n(F)$ is an F -v.s. isomorphic to F^{n+1} .

Example Consider the differentiation map

$$D: P_2(F) \longrightarrow P_1(F)$$

$$a_0 + a_1 X + a_2 X^2 \longmapsto a_1 + 2a_2 X$$

Let $\alpha = (1, X)$, $\beta = (1, X, X^2)$ be the standard ordered bases for $P_1(F)$ and $P_2(F)$, respectively. Then

$$D(1) = 0 \cdot 1 + 0 \cdot X$$

$$D(X) = 1 \cdot 1 + 0 \cdot X$$

$$D(X^2) = 0 \cdot 1 + 2 \cdot X$$

$$\text{So } {}_{\alpha}[D]_{\beta} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

Change of basis

Given a linear map $T: U \rightarrow V$ between f.d.v.s. over F with $m = \dim V, n = \dim U$, and given ordered bases α, β of U, V respectively we can construct a matrix

$$A = {}_{\beta}[T]_{\alpha} \in M_{m \times n}(F)$$

such that the following diagram commutes.

$$\begin{array}{ccccc} x & U & \xrightarrow{T} & V & y \\ \downarrow & \downarrow \iota & & \downarrow \iota & \downarrow \\ [x]_{\alpha} & F^{n \times 1} & \xrightarrow{I_A} & F^{m \times 1} & [y]_{\beta} \end{array}$$
$$\begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{bmatrix} \mapsto A \cdot \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{bmatrix}$$

Q1 How does changing α and/or β affect the matrix A ?

Q2 What can we deduce about matrices from this diagram?

Def. Let V be a f.d.v.s over F with $\dim(V) = m$. Let β, β' be ordered bases for V .

We call

$${}_{\beta'}[I_V]_{\beta} \in M_m(F)$$

the change of basis matrix from β to β' . (Here $I_V: V \rightarrow V$ is the identity map.)