

# CS70 Modular Arithmetic

Kelvin Lee

*kelvinlee@berkeley.edu*

September 22, 2020

# Overview

- 1 Basic Definitions
- 2 Multiplicative Inverse
- 3 Euclid's Algorithm
- 4 Extended Euclid's algorithm
- 5 Functions
- 6 Bijection
- 7 Fermat's Little Theorem

# Basic Definitions

## Definition (Congruence)

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
- $x$  and  $y$  have the same remainder w.r.t.  $m$

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
- $x$  and  $y$  have the same remainder w.r.t.  $m$
- $x = y + km$  for some integer  $k$



## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
  - $x$  and  $y$  have the same remainder w.r.t.  $m$
  - $x = y + km$  for some integer  $k$
- 
- In modulo  $m$ , only the numbers  $\{0, 1, 2, \dots, m - 1\}$  exist.

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
  - $x$  and  $y$  have the same remainder w.r.t.  $m$
  - $x = y + km$  for some integer  $k$
- 
- In modulo  $m$ , only the numbers  $\{0, 1, 2, \dots, m - 1\}$  exist.
  - Division is not well-defined.

# Basic Definitions

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
  - $x$  and  $y$  have the same remainder w.r.t.  $m$
  - $x = y + km$  for some integer  $k$
- 
- In modulo  $m$ , only the numbers  $\{0, 1, 2, \dots, m - 1\}$  exist.
  - Division is not well-defined.

## Definition (Multiplicative Inverse)

# Basic Definitions

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y \pmod{m}$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
  - $x$  and  $y$  have the same remainder w.r.t.  $m$
  - $x = y + km$  for some integer  $k$
- 
- In modulo  $m$ , only the numbers  $\{0, 1, 2, \dots, m - 1\}$  exist.
  - Division is not well-defined.

## Definition (Multiplicative Inverse)

Normally we say that the **multiplicative inverse** of  $x$  is  $y$  if  $xy = 1$ .

# Basic Definitions

## Definition (Congruence)

$x$  is **congruent** to  $y$  modulo  $m$  or  $x \equiv y(\text{mod } m)$  if and only if any one of the following is true:

- $(x - y)$  is divisible by  $m$
  - $x$  and  $y$  have the same remainder w.r.t.  $m$
  - $x = y + km$  for some integer  $k$
- 
- In modulo  $m$ , only the numbers  $\{0, 1, 2, \dots, m - 1\}$  exist.
  - Division is not well-defined.

## Definition (Multiplicative Inverse)

Normally we say that the **multiplicative inverse** of  $x$  is  $y$  if  $xy = 1$ . In the modular space, the **multiplicative inverse** of  $x \text{ mod } m$  is  $y$  if  $xy \equiv 1(\text{mod } m)$ .

# Theorems

## Theorem (Modular operations)

## Theorem (Modular operations)

$a \equiv c \pmod{m}$  and  $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$  and  $a \cdot b \equiv c \cdot d \pmod{m}$ .



# Theorems

## Theorem (Modular operations)

$a \equiv c \pmod{m}$  and  $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$  and  $a \cdot b \equiv c \cdot d \pmod{m}$ .

## Theorem (Existence of multiplicative inverse)

# Theorems

## Theorem (Modular operations)

$a \equiv c \pmod{m}$  and  $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$  and  $a \cdot b \equiv c \cdot d \pmod{m}$ .

## Theorem (Existence of multiplicative inverse)

$\gcd(x, m) = 1 \implies x$  has a multiplicative inverse modulo  $m$  and it is **unique**.

# Euclid's Algorithm

# Euclid's Algorithm

How do we compute  $\gcd$  of two numbers  $x$  and  $y$ ?

# Euclid's Algorithm

How do we compute  $\gcd$  of two numbers  $x$  and  $y$ ?

Theorem (Euclid's Algorithm)

# Euclid's Algorithm

How do we compute gcd of two numbers  $x$  and  $y$ ?

## Theorem (Euclid's Algorithm)

*Let  $x \geq y > 0$ . Then*

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

# Euclid's Algorithm

How do we compute gcd of two numbers  $x$  and  $y$ ?

## Theorem (Euclid's Algorithm)

*Let  $x \geq y > 0$ . Then*

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

## Example

# Euclid's Algorithm

How do we compute gcd of two numbers  $x$  and  $y$ ?

## Theorem (Euclid's Algorithm)

*Let  $x \geq y > 0$ . Then*

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

## Example

Compute  $\gcd(16, 10)$ :



# Euclid's Algorithm

How do we compute gcd of two numbers  $x$  and  $y$ ?

## Theorem (Euclid's Algorithm)

Let  $x \geq y > 0$ . Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

## Example

Compute  $\gcd(16, 10)$ :

$$\begin{aligned}\gcd(16, 10) &= \gcd(10, 6) \\ &= \gcd(6, 4) \\ &= \gcd(4, 2) \\ &= \gcd(2, 0) \\ &= 2.\end{aligned}$$

# Extended Euclid's algorithm

# Extended Euclid's algorithm

How to compute the multiplicative inverse?

# Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers  $a$  and  $b$  such that:

$$\gcd(x, y) = ax + by.$$

# Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers  $a$  and  $b$  such that:

$$\gcd(x, y) = ax + by.$$

## Theorem (Bézout's Identity)

# Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers  $a$  and  $b$  such that:

$$\gcd(x, y) = ax + by.$$

## Theorem (Bézout's Identity)

*For nonzero integers  $x$  and  $y$ , let  $d$  be the greatest common divisor such that  $d = \gcd(x, y)$ . Then, there exist integers  $a$  and  $b$  such that*

$$ax + by = d.$$

# Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers  $a$  and  $b$  such that:

$$\gcd(x, y) = ax + by.$$

## Theorem (Bézout's Identity)

*For nonzero integers  $x$  and  $y$ , let  $d$  be the greatest common divisor such that  $d = \gcd(x, y)$ . Then, there exist integers  $a$  and  $b$  such that*

$$ax + by = d.$$

- When  $\gcd(x, y) = 1$ , we can deduce that  $b$  is an inverse of  $y \bmod x$ .

# Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers  $a$  and  $b$  such that:

$$\gcd(x, y) = ax + by.$$

## Theorem (Bézout's Identity)

*For nonzero integers  $x$  and  $y$ , let  $d$  be the greatest common divisor such that  $d = \gcd(x, y)$ . Then, there exist integers  $a$  and  $b$  such that*

$$ax + by = d.$$

- When  $\gcd(x, y) = 1$ , we can deduce that  $b$  is an inverse of  $y \bmod x$ .
- This uses back substitutions repetitively so that the final expression is in terms of  $x$  and  $y$ .



# Functions

## Definition (Function)

## Definition (Function)

Let  $A$  and  $B$  be nonempty sets. A **function**  $f$  from  $A$  to  $B$  is an assignment of exactly one element of  $B$  to each element of  $A$ . (vertical line test)

## Definition (Function)

Let  $A$  and  $B$  be nonempty sets. A **function**  $f$  from  $A$  to  $B$  is an assignment of exactly one element of  $B$  to each element of  $A$ . (vertical line test)

- To denote such a function, we write  $f : A \rightarrow B$  ( $f$  maps  $A$  to  $B$ ).

## Definition (Function)

Let  $A$  and  $B$  be nonempty sets. A **function**  $f$  from  $A$  to  $B$  is an assignment of exactly one element of  $B$  to each element of  $A$ . (vertical line test)

- To denote such a function, we write  $f : A \rightarrow B$  ( $f$  maps  $A$  to  $B$ ).
- $A$  is the **domain** and  $B$  is the **co-domain**.

## Definition (Function)

Let  $A$  and  $B$  be nonempty sets. A **function**  $f$  from  $A$  to  $B$  is an assignment of exactly one element of  $B$  to each element of  $A$ . (vertical line test)

- To denote such a function, we write  $f : A \rightarrow B$  ( $f$  maps  $A$  to  $B$ ).
- $A$  is the **domain** and  $B$  is the **co-domain**.
- Pre-image is a **subset** of domain, and the image/range is the **subset** of co-domain.

## Definition (Function)

Let  $A$  and  $B$  be nonempty sets. A **function**  $f$  from  $A$  to  $B$  is an assignment of exactly one element of  $B$  to each element of  $A$ . (vertical line test)

- To denote such a function, we write  $f : A \rightarrow B$  ( $f$  maps  $A$  to  $B$ ).
- $A$  is the **domain** and  $B$  is the **co-domain**.
- Pre-image is a **subset** of domain, and the image/range is the **subset** of co-domain.
  - If  $f(a) = b$ , where  $a \in A$  and  $b \in B$ , then we say that  $b$  is the image of  $a$  and  $a$  is the pre-image of  $b$ .

# Bijection



## Definition (One-to-one)

## Definition (One-to-one)

A function  $f$  is said to be **one-to-one** if and only if  $f(a) = f(a')$  implies that  $a = a'$  for all  $a, a' \in A$ . A function is said to be **injective** if it is **one-to-one**.

## Definition (One-to-one)

A function  $f$  is said to be **one-to-one** if and only if  $f(a) = f(a')$  implies that  $a = a'$  for all  $a, a' \in A$ . A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that  $a \neq a' \implies f(a) \neq f(a')$ . (Why?)

# Bijection

## Definition (One-to-one)

A function  $f$  is said to be **one-to-one** if and only if  $f(a) = f(a')$  implies that  $a = a'$  for all  $a, a' \in A$ . A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that  $a \neq a' \implies f(a) \neq f(a')$ . (Why?)

## Definition (Onto)

## Definition (One-to-one)

A function  $f$  is said to be **one-to-one** if and only if  $f(a) = f(a')$  implies that  $a = a'$  for all  $a, a' \in A$ . A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that  $a \neq a' \implies f(a) \neq f(a')$ . (Why?)

## Definition (Onto)

A function  $f$  is called **onto**, or a surjection, if and only if for every element  $b \in B$  there is an element  $a \in A$  such that  $f(a) = b$ . We also say that  $f$  is **surjective** if it's onto.

# Bijection

## Definition (One-to-one)

A function  $f$  is said to be **one-to-one** if and only if  $f(a) = f(a')$  implies that  $a = a'$  for all  $a, a' \in A$ . A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that  $a \neq a' \implies f(a) \neq f(a')$ . (Why?)

## Definition (Onto)

A function  $f$  is called **onto**, or a surjection, if and only if for every element  $b \in B$  there is an element  $a \in A$  such that  $f(a) = b$ . We also say that  $f$  is **surjective** if it's onto.

- To show that a function is *onto*, choose  $a = f^{-1}(b)$  and so  $f(f^{-1}(b)) = b$ .

## Definition (Bijection)

## Definition (Bijection)

A function  $f$  is a **bijection** if and only if it is both *one-to-one* and *onto*. We also say that  $f$  is bijective.



## Definition (Bijection)

A function  $f$  is a **bijection** if and only if it is both *one-to-one* and *onto*. We also say that  $f$  is bijective.

- If  $f : A \rightarrow B$  is a bijection, it will have an **inverse** function (a lemma from notes), and  $|A| = |B|$ .

# Fermat's Little Theorem

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

*For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have*

$$a^{p-1} \equiv 1 \pmod{p}.$$

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

*For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have*

$$a^{p-1} \equiv 1 \pmod{p}.$$

***Proof:***

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

*For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have*

$$a^{p-1} \equiv 1 \pmod{p}.$$

### ***Proof:***

- Consider  $S = \{1, 2, \dots, p-1\}$  and  $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$ .

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

*For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have*

$$a^{p-1} \equiv 1 \pmod{p}.$$

### ***Proof:***

- Consider  $S = \{1, 2, \dots, p-1\}$  and  $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$ .
- They are the same set under mod  $p$  (different order).

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

### **Proof:**

- Consider  $S = \{1, 2, \dots, p-1\}$  and  $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$ .
- They are the same set under mod  $p$  (different order).

$$\prod_{k=1}^{p-1} k \equiv \prod_{k=1}^{p-1} ka \pmod{p}$$



# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

### **Proof:**

- Consider  $S = \{1, 2, \dots, p-1\}$  and  $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$ .
- They are the same set under mod  $p$  (different order).

$$\prod_{k=1}^{p-1} k \equiv \prod_{k=1}^{p-1} ka \pmod{p}$$

$$(p-1)! \equiv a^{p-1}(p-1)! \pmod{p}$$

# Fermat's Little Theorem

## Theorem (Fermat's Little Theorem)

For any prime  $p$  and any  $a \in \{1, 2, \dots, p-1\}$ , we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

### **Proof:**

- Consider  $S = \{1, 2, \dots, p-1\}$  and  $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$ .
- They are the same set under mod  $p$  (different order).

$$\prod_{k=1}^{p-1} k \equiv \prod_{k=1}^{p-1} ka \pmod{p}$$

$$(p-1)! \equiv a^{p-1}(p-1)! \pmod{p}$$

$$a^{p-1} \equiv 1 \pmod{p}$$

# Problem Time!