

# CS70 Countability

Kelvin Lee

*kelvinlee@berkeley.edu*

October 21, 2020

# Overview

- 1 Bijection
- 2 Cardinality
- 3 Cantor-Bernstein's Theorem
- 4 Cantor's Diagonalization

How do we determine if two sets have the same **cardinality**, or size?

**How do we determine if two sets have the same **cardinality**, or size?**

- This is obvious for finite sets, but trickier for infinite sets.

**How do we determine if two sets have the same **cardinality**, or size?**

- This is obvious for finite sets, but trickier for infinite sets.
- Two finite sets have the same size if and only if their elements can be paired up, so that each element of one set has a unique partner in the other set, and vice versa.

**How do we determine if two sets have the same **cardinality**, or size?**

- This is obvious for finite sets, but trickier for infinite sets.
- Two finite sets have the same size if and only if their elements can be paired up, so that each element of one set has a unique partner in the other set, and vice versa.
- We formalize this through the concept of a **bijection**, which you should have already learned about.

**How do we determine if two sets have the same **cardinality**, or size?**

- This is obvious for finite sets, but trickier for infinite sets.
- Two finite sets have the same size if and only if their elements can be paired up, so that each element of one set has a unique partner in the other set, and vice versa.
- We formalize this through the concept of a **bijection**, which you should have already learned about.
- To show that two infinite sets have the same **cardinality**, we need to establish a bijection (one-to-one correspondence) between the two sets.

**Are there more natural numbers  $\mathbb{N}$  than there are positive integers  $\mathbb{Z}^+$ ?**



**Are there more natural numbers  $\mathbb{N}$  than there are positive integers  $\mathbb{Z}^+$ ?**


- We can actually define a mapping between the natural numbers and the positive integers.

**Are there more natural numbers  $\mathbb{N}$  than there are positive integers  $\mathbb{Z}^+$ ?**

- We can actually define a mapping between the natural numbers and the positive integers.
- Define  $f : \mathbb{N} \rightarrow \mathbb{Z}^+$  such that  $f(n) = n + 1$ . We can see there's a one-to-one correspondence (why?).


**Are there more natural numbers  $\mathbb{N}$  than there are positive integers  $\mathbb{Z}^+$ ?**

- We can actually define a mapping between the natural numbers and the positive integers.
- Define  $f : \mathbb{N} \rightarrow \mathbb{Z}^+$  such that  $f(n) = n + 1$ . We can see there's a one-to-one correspondence (why?).



**Are there more natural numbers  $\mathbb{N}$  than there are positive integers  $\mathbb{Z}^+$ ?**


- We can actually define a mapping between the natural numbers and the positive integers.
- Define  $f : \mathbb{N} \rightarrow \mathbb{Z}^+$  such that  $f(n) = n + 1$ . We can see there's a one-to-one correspondence (why?).



- This tells us that  $|\mathbb{N}| = |\mathbb{Z}^+|$ .

**Are there more natural numbers  $\mathbb{N}$  than there are positive integers  $\mathbb{Z}^+$ ?**

- We can actually define a mapping between the natural numbers and the positive integers.
- Define  $f : \mathbb{N} \rightarrow \mathbb{Z}^+$  such that  $f(n) = n + 1$ . We can see there's a one-to-one correspondence (why?).



- This tells us that  $|\mathbb{N}| = |\mathbb{Z}^+|$ .
- We have just shown that  $\infty + 1 = \infty$ !

# Examples

**What about  $\mathbb{N}$  and  $\mathbb{Z}$ ?**

## What about $\mathbb{N}$ and $\mathbb{Z}$ ?

- They both actually have the same size! Let's see why, consider the following function  $f$ :

## What about $\mathbb{N}$ and $\mathbb{Z}$ ?

- They both actually have the same size! Let's see why, consider the following function  $f$ :

$$f(x) = \begin{cases} \frac{x}{2}, & \text{if } x \text{ is even} \\ \frac{-(x+1)}{2}, & \text{if } x \text{ is odd} \end{cases}$$



## What about $\mathbb{N}$ and $\mathbb{Z}$ ?

- They both actually have the same size! Let's see why, consider the following function  $f$ :

$$f(x) = \begin{cases} \frac{x}{2}, & \text{if } x \text{ is even} \\ \frac{-(x+1)}{2}, & \text{if } x \text{ is odd} \end{cases}$$

This function is in fact a bijection. Thus, the two sets have the same size.

# Countable Sets

# Countable Sets

- A set  $S$  is **countable** if there is a bijection between  $S$  and  $\mathbb{N}$  or some subset of  $\mathbb{N}$ .

# Countable Sets

- A set  $S$  is **countable** if there is a bijection between  $S$  and  $\mathbb{N}$  or some subset of  $\mathbb{N}$ .
- Intuitively, any finite set  $S$  is clearly **countable**.

# Countable Sets

- A set  $S$  is **countable** if there is a bijection between  $S$  and  $\mathbb{N}$  or some subset of  $\mathbb{N}$ .
- Intuitively, any finite set  $S$  is clearly **countable**.
- The examples we did earlier are countable because they are subsets of  $\mathbb{N}$ , which is a **countable** set.

# Examples

# Examples

**Now consider the set of rational numbers  $\mathbb{Q}$ , is it larger than  $\mathbb{N}$ ?**

**Recall that  $\mathbb{Q} = \left\{ \frac{x}{y} \mid x, y \in \mathbb{Z}, y \neq 0 \right\}$ .**

# Examples

**Now consider the set of rational numbers  $\mathbb{Q}$ , is it larger than  $\mathbb{N}$ ?**

**Recall that  $\mathbb{Q} = \left\{ \frac{x}{y} \mid x, y \in \mathbb{Z}, y \neq 0 \right\}$ .**

- The two sets actually have the same cardinality! Let's first learn some new definitions and an important theorem.



# Examples

**Now consider the set of rational numbers  $\mathbb{Q}$ , is it larger than  $\mathbb{N}$ ?**

**Recall that  $\mathbb{Q} = \left\{ \frac{x}{y} \mid x, y \in \mathbb{Z}, y \neq 0 \right\}$ .**

- The two sets actually have the same cardinality! Let's first learn some new definitions and an important theorem.

## Definition 1

# Examples

**Now consider the set of rational numbers  $\mathbb{Q}$ , is it larger than  $\mathbb{N}$ ?**

**Recall that  $\mathbb{Q} = \left\{ \frac{x}{y} \mid x, y \in \mathbb{Z}, y \neq 0 \right\}$ .**

- The two sets actually have the same cardinality! Let's first learn some new definitions and an important theorem.

## **Definition 1**

If there is a injective function  $f : A \rightarrow B$ , then  $|A| \leq |B|$ .

# Examples

**Now consider the set of rational numbers  $\mathbb{Q}$ , is it larger than  $\mathbb{N}$ ?**

**Recall that  $\mathbb{Q} = \left\{ \frac{x}{y} \mid x, y \in \mathbb{Z}, y \neq 0 \right\}$ .**

- The two sets actually have the same cardinality! Let's first learn some new definitions and an important theorem.

## **Definition 1**

If there is a injective function  $f : A \rightarrow B$ , then  $|A| \leq |B|$ .

## **Definition 2**

# Examples

**Now consider the set of rational numbers  $\mathbb{Q}$ , is it larger than  $\mathbb{N}$ ?**

**Recall that  $\mathbb{Q} = \left\{ \frac{x}{y} \mid x, y \in \mathbb{Z}, y \neq 0 \right\}$ .**

- The two sets actually have the same cardinality! Let's first learn some new definitions and an important theorem.

## **Definition 1**

If there is a injective function  $f : A \rightarrow B$ , then  $|A| \leq |B|$ .

## **Definition 2**

If there is a surjective function  $f : A \rightarrow B$ , then  $|A| \geq |B|$ .

# Cantor-Bernstein's Theorem

# Cantor-Bernstein's Theorem

## Schröder–Bernstein Theorem (Cantor–Bernstein)

# Cantor-Bernstein's Theorem

## Schröder–Bernstein Theorem (Cantor–Bernstein)

If  $A$  and  $B$  are sets with  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ . In other words, if there are injective functions  $f : A \rightarrow B$  and  $g : B \rightarrow A$ , then there is a bijection  $h$  between  $A$  and  $B$ .

# Cantor-Bernstein's Theorem

## Schröder–Bernstein Theorem (Cantor–Bernstein)

If  $A$  and  $B$  are sets with  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ . In other words, if there are injective functions  $f : A \rightarrow B$  and  $g : B \rightarrow A$ , then there is a bijection  $h$  between  $A$  and  $B$ .

***Proof:***



# Cantor-Bernstein's Theorem

## Schröder–Bernstein Theorem (Cantor–Bernstein)

If  $A$  and  $B$  are sets with  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ . In other words, if there are injective functions  $f : A \rightarrow B$  and  $g : B \rightarrow A$ , then there is a bijection  $h$  between  $A$  and  $B$ .

### ***Proof:***

The proof of this theorem is out of scope for this class. We'll skip that for now. □

# Cantor-Bernstein's Theorem

## Schröder–Bernstein Theorem (Cantor–Bernstein)

If  $A$  and  $B$  are sets with  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ . In other words, if there are injective functions  $f : A \rightarrow B$  and  $g : B \rightarrow A$ , then there is a bijection  $h$  between  $A$  and  $B$ .

### ***Proof:***

The proof of this theorem is out of scope for this class. We'll skip that for now. □

- This theorem will be very useful when showing a set  $S$  is countable. We can give separate injections  $f : S \rightarrow \mathbb{N}$  and  $g : \mathbb{N} \rightarrow S$ , instead of designing a bijection (which is trickier).

# Examples

# Examples

- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .

# Examples

- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .
- Now the theorem comes in handy and all we need to do now is to prove  $|\mathbb{Q}| \leq |\mathbb{N}|$ .

# Examples

- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .
- Now the theorem comes in handy and all we need to do now is to prove  $|\mathbb{Q}| \leq |\mathbb{N}|$ .
- Recall the definition, we must exhibit an injection  $f : \mathbb{Q} \rightarrow \mathbb{N}$ .

# Examples

- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .
- Now the theorem comes in handy and all we need to do now is to prove  $|\mathbb{Q}| \leq |\mathbb{N}|$ .
- Recall the definition, we must exhibit an injection  $f : \mathbb{Q} \rightarrow \mathbb{N}$ .
- Notice that each rational number  $\frac{a}{b}$  ( $\gcd(a, b) = 1$ ) can be represented by the point  $(a, b) \in \mathbb{Z} \times \mathbb{Z}$  (the set of all pairs of integers).

# Examples

- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .
- Now the theorem comes in handy and all we need to do now is to prove  $|\mathbb{Q}| \leq |\mathbb{N}|$ .
- Recall the definition, we must exhibit an injection  $f : \mathbb{Q} \rightarrow \mathbb{N}$ .
- Notice that each rational number  $\frac{a}{b}$  ( $\gcd(a, b) = 1$ ) can be represented by the point  $(a, b) \in \mathbb{Z} \times \mathbb{Z}$  (the set of all pairs of integers).
- However, not all points are valid.



# Examples


- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .
- Now the theorem comes in handy and all we need to do now is to prove  $|\mathbb{Q}| \leq |\mathbb{N}|$ .
- Recall the definition, we must exhibit an injection  $f : \mathbb{Q} \rightarrow \mathbb{N}$ .
- Notice that each rational number  $\frac{a}{b}$  ( $\gcd(a, b) = 1$ ) can be represented by the point  $(a, b) \in \mathbb{Z} \times \mathbb{Z}$  (the set of all pairs of integers).
- However, not all points are valid.
- Thus, we can actually tell that  $|\mathbb{Z} \times \mathbb{Z}| \geq |\mathbb{Q}|$ .

# Examples


- Now back to our problem. First it is obvious that  $|\mathbb{N}| \leq |\mathbb{Q}|$  because  $\mathbb{N} \subseteq \mathbb{Q}$ .
- Now the theorem comes in handy and all we need to do now is to prove  $|\mathbb{Q}| \leq |\mathbb{N}|$ .
- Recall the definition, we must exhibit an injection  $f : \mathbb{Q} \rightarrow \mathbb{N}$ .
- Notice that each rational number  $\frac{a}{b}$  ( $\gcd(a, b) = 1$ ) can be represented by the point  $(a, b) \in \mathbb{Z} \times \mathbb{Z}$  (the set of all pairs of integers).
- However, not all points are valid.
- Thus, we can actually tell that  $|\mathbb{Z} \times \mathbb{Z}| \geq |\mathbb{Q}|$ .
- If we are able to come up with an injection from  $\mathbb{Z} \times \mathbb{Z}$  to  $\mathbb{N}$ , then this will also be an injection from  $\mathbb{Q}$  to  $\mathbb{N}$  (why?).

# Examples

## Examples




# Examples




- Map each pair  $(a, b)$  to its position along the spiral, starting at the origin (indexing through each point starting from index 0).

# Examples



- Map each pair  $(a, b)$  to its position along the spiral, starting at the origin (indexing through each point starting from index 0).
- This mapping maps every pair of integers injectively to a natural number.

## Examples



- Map each pair  $(a, b)$  to its position along the spiral, starting at the origin (indexing through each point starting from index 0).
- This mapping maps every pair of integers injectively to a natural number.
- Thus we have  $|\mathbb{Q}| \leq |\mathbb{Z} \times \mathbb{Z}| \leq |\mathbb{N}|$ . Remember that  $|\mathbb{N}| \leq |\mathbb{Q}|$ , then by the Cantor-Bernstein Theorem  $|\mathbb{N}| = |\mathbb{Q}|$ .

# Cantor's Diagonalization



# Cantor's Diagonalization

- An set  $S$  is **uncountable** if it is not **countable**, or cannot be enumerated. This introduces the concept of **diagonalization**.

# Cantor's Diagonalization

- An set  $S$  is **uncountable** if it is not **countable**, or cannot be enumerated. This introduces the concept of **diagonalization**.
- The idea of **diagonalization** is that we can always create an element that does not belong to the set, i.e., enumeration is impossible.

# Cantor's Diagonalization

- An set  $S$  is **uncountable** if it is not **countable**, or cannot be enumerated. This introduces the concept of **diagonalization**.
- The idea of **diagonalization** is that we can always create an element that does not belong to the set, i.e., enumeration is impossible.
- One example would be  $\mathbb{R}[0, 1]$  demonstrated in lecture.

# Cantor's Diagonalization

- An set  $S$  is **uncountable** if it is not **countable**, or cannot be enumerated. This introduces the concept of **diagonalization**.
- The idea of **diagonalization** is that we can always create an element that does not belong to the set, i.e., enumeration is impossible.
- One example would be  $\mathbb{R}[0, 1]$  demonstrated in lecture.
- We can create a real number where each of its  $i$ th digit differs from the  $i$ th digit of the  $i$ th element.

# Cantor's Diagonalization

- An set  $S$  is **uncountable** if it is not **countable**, or cannot be enumerated. This introduces the concept of **diagonalization**.
- The idea of **diagonalization** is that we can always create an element that does not belong to the set, i.e., enumeration is impossible.
- One example would be  $\mathbb{R}[0, 1]$  demonstrated in lecture.
- We can create a real number where each of its  $i$ th digit differs from the  $i$ th digit of the  $i$ th element.
- Thus the real interval  $\mathbb{R}[0, 1]$  is uncountable, so do its supersets.

# Problem Time!